



Cochrane
Library

Cochrane Database of Systematic Reviews

Interventions for preventing high altitude illness: Part 1. Commonly-used classes of drugs (Review)

Nieto Estrada VH, Molano Franco D, Medina RD, Gonzalez Garay AG, Martí-Carvajal AJ, Arevalo-Rodriguez I

Nieto Estrada VH, Molano Franco D, Medina RD, Gonzalez Garay AG, Martí-Carvajal AJ, Arevalo-Rodriguez I.

Interventions for preventing high altitude illness: Part 1. Commonly-used classes of drugs.

Cochrane Database of Systematic Reviews 2017, Issue 6. Art. No.: CD009761.

DOI: 10.1002/14651858.CD009761.pub2.

www.cochranelibrary.com

TABLE OF CONTENTS

HEADER	1
ABSTRACT	1
PLAIN LANGUAGE SUMMARY	2
SUMMARY OF FINDINGS FOR THE MAIN COMPARISON	4
BACKGROUND	6
OBJECTIVES	8
METHODS	8
RESULTS	11
Figure 1.	12
Figure 2.	14
Figure 3.	15
Figure 4.	18
Figure 5.	19
Figure 6.	21
ADDITIONAL SUMMARY OF FINDINGS	31
DISCUSSION	36
AUTHORS' CONCLUSIONS	37
ACKNOWLEDGEMENTS	37
REFERENCES	38
CHARACTERISTICS OF STUDIES	48
DATA AND ANALYSES	168
Analysis 1.1. Comparison 1 Carbonic anhydrase inhibitors: acetazolamide versus placebo, Outcome 1 Incidence of acute mountain sickness.	170
Analysis 1.2. Comparison 1 Carbonic anhydrase inhibitors: acetazolamide versus placebo, Outcome 2 Incidence of high altitude pulmonary oedema.	172
Analysis 1.3. Comparison 1 Carbonic anhydrase inhibitors: acetazolamide versus placebo, Outcome 3 Incidence of high altitude cerebral oedema.	173
Analysis 1.4. Comparison 1 Carbonic anhydrase inhibitors: acetazolamide versus placebo, Outcome 4 Incidence of adverse events: Paraesthesia.	174
Analysis 1.5. Comparison 1 Carbonic anhydrase inhibitors: acetazolamide versus placebo, Outcome 5 Differences in HAI/AMS scores.	175
Analysis 2.1. Comparison 2 Steroids: budesonide vs. placebo, Outcome 1 Incidence of acute mountain sickness.	176
Analysis 3.1. Comparison 3 Steroids: dexamethasone vs. placebo, Outcome 1 Incidence of acute mountain sickness.	177
Analysis 3.2. Comparison 3 Steroids: dexamethasone vs. placebo, Outcome 2 Differences in HAI/AMS scores.	178
Analysis 4.1. Comparison 4 Calcium modulators: nifedipine vs. placebo, Outcome 1 Differences in HAI/AMS scores.	179
Analysis 5.1. Comparison 5 NSAIDs and other analgesic: aspirin vs. placebo, Outcome 1 Incidence of AMS.	179
Analysis 6.1. Comparison 6 NSAIDs and other analgesic: ibuprofen vs. placebo, Outcome 1 Incidence of acute mountain sickness.	180
ADDITIONAL TABLES	180
APPENDICES	190
HISTORY	205
CONTRIBUTIONS OF AUTHORS	206
DECLARATIONS OF INTEREST	206
SOURCES OF SUPPORT	207
DIFFERENCES BETWEEN PROTOCOL AND REVIEW	207

[Intervention Review]

Interventions for preventing high altitude illness: Part I. Commonly-used classes of drugs

Víctor H Nieto Estrada¹, Daniel Molano Franco², Roger David Medina³, Alejandro G Gonzalez Garay⁴, Arturo J Martí-Carvajal⁵, Ingrid Arevalo-Rodriguez⁶

¹Department of Critical Care Medicine, Hospital de San José, Fundación Universitaria de Ciencias de la Salud, Bogotá, Colombia.

²Department of Critical Care, Fundación Universitaria de Ciencias de la Salud, Hospital de San Jose, Clinica universitaria Colombia UNISANITAS, Bogota, Colombia. ³Division of Research, Fundación Universitaria de Ciencias de la Salud, Bogotá D.C., Colombia.

⁴Methodology Research Unit/Neonatology, National Institute of Pediatrics, Mexico City, Mexico. ⁵Iberoamerican Cochrane Network, Valencia, Venezuela. ⁶Cochrane Ecuador. Centro de Investigación en Salud Pública y Epidemiología Clínica (CISPEC). Facultad de Ciencias de la Salud Eugenio Espejo, Universidad Tecnológica Equinoccial, Quito, Ecuador

Contact address: Ingrid Arevalo-Rodriguez, Division of Research, Fundación Universitaria de Ciencias de la Salud - Hospital de San Jose/Hospital Infantil de San Jose, Bogotá D.C., Colombia. inarev7@yahoo.com, iarevalo@fucsalud.edu.co.

Editorial group: Cochrane Anaesthesia, Critical and Emergency Care Group.

Publication status and date: New, published in Issue 6, 2017.

Citation: Nieto Estrada VH, Molano Franco D, Medina RD, Gonzalez Garay AG, Martí-Carvajal AJ, Arevalo-Rodriguez I. Interventions for preventing high altitude illness: Part 1. Commonly-used classes of drugs. *Cochrane Database of Systematic Reviews* 2017, Issue 6. Art. No.: CD009761. DOI: 10.1002/14651858.CD009761.pub2.

Copyright © 2017 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

ABSTRACT

Background

High altitude illness (HAI) is a term used to describe a group of cerebral and pulmonary syndromes that can occur during travel to elevations above 2500 metres (8202 feet). Acute hypoxia, acute mountain sickness (AMS), high altitude cerebral oedema (HACE) and high altitude pulmonary oedema (HAPE) are reported as potential medical problems associated with high altitude. In this review, the first in a series of three about preventive strategies for HAI, we assess the effectiveness of six of the most recommended classes of pharmacological interventions.

Objectives

To assess the clinical effectiveness and adverse events of commonly-used pharmacological interventions for preventing acute HAI.

Search methods

We searched the Cochrane Central Register of Controlled Trials (CENTRAL), MEDLINE (OVID), Embase (OVID), LILACS and trial registries in January 2017. We adapted the MEDLINE strategy for searching the other databases. We used a combination of thesaurus-based and free-text terms to search.

Selection criteria

We included randomized-controlled and cross-over trials conducted in any setting where commonly-used classes of drugs were used to prevent acute HAI.

Data collection and analysis

We used standard methodological procedures as expected by Cochrane.

Interventions for preventing high altitude illness: Part I. Commonly-used classes of drugs (Review)

Copyright © 2017 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

Main results

We included 64 studies (78 references) and 4547 participants in this review, and classified 12 additional studies as ongoing. A further 12 studies await classification, as we were unable to obtain the full texts. Most of the studies were conducted in high altitude mountain areas, while the rest used low pressure (hypobaric) chambers to simulate altitude exposure. Twenty-four trials provided the intervention between three and five days prior to the ascent, and 23 trials, between one and two days beforehand. Most of the included studies reached a final altitude of between 4001 and 5000 metres above sea level. Risks of bias were unclear for several domains, and a considerable number of studies did not report adverse events of the evaluated interventions. We found 26 comparisons, 15 of them comparing commonly-used drugs versus placebo. We report results for the three most important comparisons:

Acetazolamide versus placebo (28 parallel studies; 2345 participants)

The risk of AMS was reduced with acetazolamide (risk ratio (RR) 0.47, 95% confidence interval (CI) 0.39 to 0.56; $I^2 = 0\%$; 16 studies; 2301 participants; moderate quality of evidence). No events of HAPE were reported and only one event of HACE (RR 0.32, 95% CI 0.01 to 7.48; 6 parallel studies; 1126 participants; moderate quality of evidence). Few studies reported side effects for this comparison, and they showed an increase in the risk of paraesthesia with the intake of acetazolamide (RR 5.53, 95% CI 2.81 to 10.88, $I^2 = 60\%$; 5 studies, 789 participants; low quality of evidence).

Budenoside versus placebo (2 parallel studies; 132 participants)

Data on budenoside showed a reduction in the incidence of AMS compared with placebo (RR 0.37, 95% CI 0.23 to 0.61; $I^2 = 0\%$; 2 studies, 132 participants; low quality of evidence). Studies included did not report events of HAPE or HACE, and they did not find side effects (low quality of evidence).

Dexamethasone versus placebo (7 parallel studies; 205 participants)

For dexamethasone, the data did not show benefits at any dosage (RR 0.60, 95% CI 0.36 to 1.00; $I^2 = 39\%$; 4 trials, 176 participants; low quality of evidence). Included studies did not report events of HAPE or HACE, and we rated the evidence about adverse events as of very low quality.

Authors' conclusions

Our assessment of the most commonly-used pharmacological interventions suggests that acetazolamide is an effective pharmacological agent to prevent acute HAI in dosages of 250 to 750 mg/day. This information is based on evidence of moderate quality. Acetazolamide is associated with an increased risk of paraesthesia, although there are few reports about other adverse events from the available evidence. The clinical benefits and harms of other pharmacological interventions such as ibuprofen, budenoside and dexamethasone are unclear. Large multicentre studies are needed for most of the pharmacological agents evaluated in this review, to evaluate their effectiveness and safety.

PLAIN LANGUAGE SUMMARY

Drugs commonly-used for preventing high altitude illness

Background

High altitude illness (HAI) is a term used to describe a group of brain and breathing conditions that can occur while travelling to altitudes above 2500 metres (8202 feet). HAI is generally characterized by headache, nausea, vomiting and tiredness (often called acute mountain sickness), but may affect the brain or the lungs in different individuals. In this review, we assessed the most commonly-used drugs to prevent the onset of this illness.

Study characteristics

The evidence is current to January 2017. We included 64 studies related to six different types of drugs recommended for HAI prevention. Most of the studies were conducted in high altitude mountain areas, while the rest used low pressure (hypobaric) chambers to simulate altitude exposure. The participants' ages ranged between 16 and 65 years. Eleven studies included people at a high risk of this condition due to their history of HAI or other illnesses such as asthma. Twenty-four trials provided the intervention between three and five days prior to the ascent, and 23 trials, between one and two days beforehand. Most of the included studies reached a final altitude of between 4001 and 5000 metres above sea level. In 23 of the included studies, the source of funding was unclear. Only 18 studies declared their possible conflicts of interests. We classed 24 more studies as still ongoing or waiting for assessment.

Key results

Our findings suggest that acetazolamide is an effective treatment for the prevention of acute HAI in dosages of 250 to 750 mg/day, when this drug is compared to a placebo (i.e. a pill with no active agent). Most of the available information relates to the prevention of uncomplicated HAI (headache, nausea, vomiting and tiredness) rather than to more serious brain or lung problems. We also found that acetazolamide is associated with an increased risk of paraesthesia in the fingers (i.e. a sensation of tingling, tickling, pricking, or burning of the skin), although this outcome is not well reported in the available evidence. The benefits and harms of other drugs such as ibuprofen, budenoside and dexamethasone are unclear, due to the small number of studies.

Quality of the evidence

We rated the quality of the evidence as moderate to very low. Several studies had quality shortcomings, including their use of small numbers of participants and a lack of reporting of important outcomes such as side effects. For most of the drugs covered by the studies, additional research is required to clarify their effectiveness and safety.

SUMMARY OF FINDINGS FOR THE MAIN COMPARISON *[Explanation]*

Acetazolamide compared with placebo for preventing high altitude illness						
Patient or population: people at risk of high altitude illness Setting: High altitude; studies undertaken in India, South America and USA. Intervention: acetazolamide Comparison: placebo						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Placebo	Acetazolamide				
Incidence of acute mountain sickness (AMS)- Follow-up: From arrival to 24 hours later	241 per 1000	113 per 1000 (94 to 135)	RR 0.47 (0.39 to 0.56)	2301 (16 studies)	⊕⊕⊕○ moderate ¹	
Incidence of high altitude pulmonary oedema (HAPE)- Follow-up: From arrival to 24 hours later	See comment	See comment	Not estimable	1138 (7 studies)	⊕⊕⊕○ moderate ²	These trials reported no event
Incidence of high altitude cerebral oedema (HACE)- Follow-up: From arrival to 24 hours later	2 per 1000	1 per 1000 (0 to 14)	RR 0.32 (0.01 to 7.48)	1126 (6 studies)	⊕⊕⊕○ moderate ²	
Adverse events: Pares-thesias- Follow-up: From arrival to 24 hours later	91 per 1000	504 per 1000 (256 to 992)	RR 5.53 (2.81 to 10.88)	789 (5 studies)	⊕⊕○○ Low ³	

Adverse events: side effects- Follow-up: From arrival to 24 hours later	106 per 1000	232 per 1000 (144 to 374)	RR 2.19 (1.36 to 3.53)	400 (1 study)	⊕⊕○○ Low ⁴
--	---------------------	-------------------------------------	----------------------------------	------------------	---------------------------------

The **corresponding risk** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).
CI: Confidence interval; **RR:** Risk ratio;

GRADE Working Group grades of evidence

High quality: Further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: We are very uncertain about the estimate.

¹Risk of bias downgraded (-1) due to unclear selection, performance and detection bias in most of included studies. High risk of attrition bias in five studies.

²Risk of bias downgraded (-1) due to unclear selection, performance and detection bias.

³ Risk of bias downgraded (-2) due to unclear selection, performance and detection bias, as well as considerable heterogeneity (60%)

⁴Risk of bias downgraded (-2) due to high levels of attrition bias.

BACKGROUND

High altitude illness (HAI) is a term used to describe a group of cerebral and pulmonary syndromes that can occur during travel to elevations above 2500 metres (8202 feet). HAI is arbitrarily classified as high (1500 to 3500 metres or 4921 to 11,842 feet), very high (3500 to 5500 metres or 11,842 to 18,044 feet) and extreme (above 5500 metres or 18,044 feet) (Paralíkar 2010). Because of the large number of people who ascend rapidly to between 2500 and 3500 m (8202 to 11,842 ft), high altitude illness is common in this height range (Paralíkar 2010). Although the proportion of oxygen remains unchanged at 20.93%, increases in altitude result in a lower partial pressure of oxygen in the inspired air (Anonymous 1892; Wilson 2009). This reduction in the driving pressure of oxygen, along the oxygen cascade from the lungs to the tissues, can compromise the supply of oxygen to the tissues (Wilson 2009), especially the cardiovascular and pulmonary systems (Leissner 2009). The physiological responses to hypoxia and acclimatization related to HAI include hyperventilation (increased depth and rate of breathing), elevation of systemic blood pressure and tachycardia (elevations of heart rate) (Leissner 2009; Naeije 2010). However, in many instances these physiologic changes may be inadequate, such that the sojourn to altitude and the concomitant hypoxia are complicated by altitude-associated medical illness (Palmer 2010), which is also known as high altitude illness.

Description of the condition

High altitude illness (HAI)

There are two types of mountain sicknesses: acute mountain sickness (AMS) and chronic mountain sickness (CMS), also called Monge's disease (Monge 1942). Acute hypoxia, acute mountain sickness, high altitude cerebral oedema (HACE), high altitude pulmonary oedema (HAPE), cerebrovascular syndromes, peripheral oedema, retinopathy, thromboembolism, sleep disorders and periodic breathing, high altitude pharyngitis and bronchitis, ultraviolet exposure and keratitis (snow blindness), and exacerbation of pre-existing illness are reported as medical problems potentially associated with high altitude ascent (CATMAT 2007; Palmer 2010; Schoene 2008). Factors such as the rate of ascent, the absolute change in altitude, and individual physiology are the primary determinants of whether HAI will develop or not (Leissner 2009; Palmer 2010). The risk categories for acute mountain sickness are shown in Appendix 1 (Luks 2010).

In the 19th century, Dr Daniel Vergara, a Mexican physiologist, pioneered the studies on high altitude physiology and the physiological and anatomical mechanisms of adaptation to high elevations. Forty years later, Dr Carlos Monge, a Peruvian physiologist, reported his ideas on this issue. The work of these pioneers was summarized early this century (Rodríguez de Romo 2002).

Both the physiology and pathophysiology of high altitude have recently been widely reviewed (Bärtsch 2007; Leissner 2009; Palmer 2010; Paralíkar 2010). In brief, these reviews confirm both the increase in respiratory rate and increase in haemoglobin concentration on exposure to a low oxygen pressure, and that such changes are often inadequate. They identify the rate of ascent, the absolute change in altitude and individual variation in physiology as the primary determinants of whether HAI will develop or not (Palmer 2010). HAI is considered an important cause of mountain mortality (Windsor 2009).

Acute mountain sickness (AMS) or high altitude cerebral oedema (HACE)

AMS is a multisystem disorder with prominent neurological features characterized by headache, anorexia, nausea and sometimes vomiting, light-headedness, insomnia, and fatigue (Bailey 2009a; Leissner 2009; Palmer 2010). Headache is the most prevalent symptom of acute mountain sickness. In contrast, HACE is a potentially fatal neurologic disorder and it is characterized by altered consciousness or ataxia (Bailey 2009a; Hackett 2004; Imray 2010), or both, in an individual with AMS or high altitude pulmonary oedema (HAPE). If left untreated, HACE can result in death due to cerebral oedema (Bailey 2009a). HACE is widely viewed as the end stage of AMS and is normally preceded by symptoms of AMS (Basnyat 2003), which suggest a similar pathophysiologic process (Bailey 2009a; Imray 2010; Palmer 2010). Both syndromes share a common pathophysiology linked by intracranial hypertension (Bailey 2009a; Kallenberg 2007; Schoonman 2008; Wilson 2009). The severity of AMS can be scored using the Lake Louise Questionnaire, Environmental Symptoms Questionnaire, or by the use of a simple analogue scale (Imray 2010). Headache is a very common symptom at altitude and some authors have suggested it could be viewed as a distinct clinical entity.

The definition of AMS seems to be problematic, as it will vary greatly between studies. A Lake Louise Score higher than two (including headache) is not equivalent to a criterion score of 0.70 with AMS-C (cerebral) from the Environmental Symptoms Questionnaire (Maggiorini 1998). It has been suggested that a previous review came to an erroneous conclusion because they included a study which used the AMS-R (respiratory) score for diagnosis of AMS. The value of the AMS-R score is questionable for diagnosing AMS (Dumont 2000). Pathophysiology with a focus on the molecular basis of AMS and HACE has been widely described by Bailey 2009a, and advances in the genetics, molecular mechanisms, and physiology that underpin them have been extensively described by Wilson 2009.

This review treats headache as a common and early symptom of AMS. Indeed, the exact definition of what constitutes AMS will vary when using different scoring systems and when interpreted by different authors. In this review we have taken care not to pool data inappropriately where the scoring systems used cannot be directly compared.

High altitude pulmonary oedema (HAPE)

HAPE is a non-cardiogenic pulmonary oedema (Luks 2008a; Schoene 2004; Stream 2008). It is characterized by cough, progressive dyspnoea with exertion, and decreased exercise tolerance, generally developing within two to four days after arrival at high altitude (Palmer 2010; Stream 2008). It is rare after one week of acclimatization at a particular altitude (Maggiorini 2010; Palmer 2010). Hypoxia is the trigger that results in a complex cascade of events leading to HAPE (Stream 2008). Essentially, HAPE is due to a “persistent imbalance between the forces that drive water into the airspace and the biologic mechanisms for its removal” (Scherrer 2010), with the hallmark of this condition being hypoxic pulmonary hypertension. The hypertension may be mediated by at least four mechanisms: defective pulmonary nitric oxide synthesis, exaggerated endothelin-1 synthesis, exaggerated sympathetic activation, and a defect in alveolar transepithelial sodium transport (Scherrer 2010). An extensive review of pulmonary hypertension induced by HAI is reported by Pasha 2010.

Epidemiology of acute HAI

It has been estimated that 84% of people who fly directly to 3860 m are affected by AMS (Basnyat 2003). The incidence of HACE and HAPE is much lower than for AMS, with estimates in the range of 0.1% to 4.0% (Basnyat 2003). The rate of ascent, altitude reached (especially the sleeping altitude), and individual susceptibility are the most important risk factors for the development of HAI (Basnyat 2003; Schneider 2002). Other risk factors are a history of HAI and permanent residence lower than 900 metres, exertion in children and adults (Basnyat 2003), obesity (Ri-Li 2003), and coronary heart disease (Dehnert 2010). It is advisable that those with asthma be sure that their condition is well controlled before they undertake exertion at altitude (CATMAT 2007). See Appendix 2 for other medical terms.

Description of the intervention

The risk of high altitude illness (HAI) begins with a non-acclimatized person ascending to an altitude higher than 2500 metres (Paralihar 2010). However, a susceptible individual may develop AMS at an intermediate altitude such as 2000 metres (Montgomery 1989). Several interventions to prevent HAI have been described, compiled, and published in guidelines and consensus statements (CATMAT 2007; Luks 2010). Interventions for HAI prevention can be classified as pharmacological and non-pharmacological (Bärtsch 1992; Luks 2010; Luks 2008b; Wright 2008). The Committee to Advise on Tropical Medicine and Travel proposed a consensus for HAI in 2007, describing prevention and treatment approaches among several topics regarding this medical condition (CATMAT 2007).

In 2014, the Wilderness Medical Society (WMS) published an update of their 2010 guidelines (Luks 2010), detailing preven-

tion and treatment directives for HAI (AMS, HACE, HAPE). This guideline was developed by an expert panel that compiled and classified all available evidence on HAI prevention and treatment. Recommendations based on evidence, using American College of Chest Physicians strategies, were agreed upon. For AMS and HACE, the experts proposed a risk classification where low-risk people are discarded for prevention interventions. For HAPE, pharmacological prophylaxis is recommended for those with a previous diagnosis of HAI (Luks 2014). However, the document does not include all of the most frequent and broadly-described pharmacological interventions for prevention and treatment of HAI. The most commonly suggested interventions are summarized below.

1. **Carbonic anhydrase inhibitors:** acetazolamide and methazolamide (Bernhard 1998; Carlsten 2004; Hussain 2004; Swenson 2007; Van Patot 2008; Wright 1983; Wright 2008).
2. **Steroids:** budesonide, prednisolone and dexamethasone (Basu 2002a; Basu 2002b; Ellsworth 1991; Hackett 1988; Johnson 1984; Rock 1989a).
3. **Bronchodilator drugs:** Include salmeterol, theophylline and montelukast (Sartori 2002; Kleinsasser 2002; Wright 2008).
4. **Selective inhibitor of phosphodiesterase type 5 (PDE5):** tadalafil (Maggiorini 2006) and sildenafil (Bates 2007; Kleinsasser 2002; Richalet 2005).
5. **Calcium modulators:** Include nifedipine and flunarizine (Bartsch 1991; Hohenhaus 1994).
6. **Non-steroidal anti-inflammatory drugs (NSAIDs)** and other analgesic: aspirin, carbasalate and ibuprofen (Burtscher 1998; Burtscher 2001).

How the intervention might work

Extensive reviews of the pharmacotherapy of HAI have recently been published (Maggiorini 2010; Wright 2008). Below is a list and brief description of the common agents that have so far been suggested. Appendix 3 provides more detail, and discusses the potential adverse effects of each agent.

1. Carbonic anhydrase (CA) inhibitors (acetazolamide and methazolamide) generate inhibition of CA in the kidneys, resulting in increased bicarbonate excretion in the urine and metabolic acidoses. The result is an offsetting of hyperventilation-induced respiratory alkalosis, allowing chemoreceptors to respond more fully to hypoxic stimuli at altitude (Leaf 2007). Acetazolamide can also cause pulmonary vasodilation unrelated to carbonic anhydrase inhibition (Höhne 2007; Swenson 2006).
2. Steroids (dexamethasone, budesonide and prednisolone): Hypoxia-induced vasogenic oedema has been suggested as one of the major mechanisms responsible for development of AMS (Hackett 1999). Glucocorticoids blocks hypoxia-induced endothelial dysfunction (Murata 2004; Murata 2005).

3. Bronchodilators (salmeterol, theophylline or aminophylline, montelukast). The human beta-2 adrenergic receptor (B2AR) has been found to play a very important role in the pathogenesis of HAPE, and salmeterol was found to have a high binding affinity with human B2AR (Chandramoorthi 2008). Furthermore, salmeterol enhances alveolar clearance by stimulating amiloride-sensitive sodium (Na) channels (Maggiolini 2010). Non-selective phosphodiesterase inhibitor (theophylline or aminophylline): anti-hypoxia and antioxidation effects of aminophylline (Yang 2007) could be responsible for reducing periodic breathing, cerebral and pulmonary microvascular permeability, and pulmonary artery pressure (Wright 2008). Montelukast is a leukotriene receptor antagonist (LTRA) that reduces the bronchoconstriction (Tintinger 2010).

4. Selective inhibitors of phosphodiesterase type 5 (tadalafil and sildenafil) induce overproduction of nitric oxide, which attenuates pulmonary vasoconstriction during acute hypoxia (Ozaki 2001; Zhao 2001). It causes a reduction in pulmonary hypertension.

5. Calcium channel blockers (CCBs): calcium channel antagonists or calcium antagonists (nifedipine, flunarizine) are a group of medications that disrupt the movement of calcium (Ca²⁺) through calcium channels and reduce pulmonary vascular resistance (Hackett 1992), leading to a reduction of the pulmonary hypertension.

6. Non-steroidal anti-inflammatory drugs (NSAIDs) and other analgesics (aspirin, ibuprofen, carbasalate): it is postulated that prostaglandin-mediated increases in cerebral microvascular permeability may contribute to the pathophysiology of AMS, and treatment with prostaglandin synthesis inhibitors could reduce this response (CATMAT 2007).

See Appendix 3 for adverse events of the pharmacological interventions.

Why it is important to do this review

It is important to conduct this systematic review for many reasons. First, many people all over the world travel to recreational areas located at high altitudes, putting themselves at increased risk of developing acute HAI. HAI may be severe and life-threatening, so effective prevention is likely to be of great value both to these visitors to high-altitude areas, and to those responsible for their treatment and rescue when required. At the other end of the spectrum, reliable prevention of minor degrees of AMS would greatly enhance the experience of many travellers. Travel to high altitudes may also aggravate underlying illnesses, particularly cardiopulmonary diseases (CATMAT 2007). Second, the true role of the many approaches for preventing acute HAI is uncertain (Adams 2004; Bärtsch 2004; CATMAT 2007; Elphick 2004), meaning that their clinical effectiveness and safety must be assessed. Third, it is necessary to answer questions such as: Are all of these interventions equally useful regardless of the type of HAI? and Is there

a reason to believe that some forms are more appropriate for some persons at risk than others?. Four, an updated meta-analysis on AMS prevention needs to be produced (Dumont 2000).

A systematic review, including a rigorous assessment of the risks of bias, of the most up-to-date evidence, will help clinicians make informed decisions about the use of non-pharmacological and pharmacological interventions for preventing acute HAI. The protocol for this review included all agents to prevent high altitude illness (Martí-Carvajal 2012), but we have decided to split the review into a series of three publications about the prevention of this condition (Part 1: Commonly-used drugs. Part 2: Less commonly-used drugs. Part 3: Miscellaneous and non-pharmacological interventions). This review includes six groups of the most highly recommended agents to prevent acute HAI.

OBJECTIVES

To assess the clinical effectiveness and adverse events of commonly-used interventions for preventing acute HAI.

METHODS

Criteria for considering studies for this review

Types of studies

We include randomized controlled trials (RCTs) irrespective of publication status (trials may be unpublished or published as articles, abstracts, or letters), language (no language limitation) or country. We applied no restrictions by length of follow-up. We also included cross-over trials (See Differences between protocol and review and section).

We excluded quasi-randomized studies and prospective observational studies for evaluating clinical effectiveness.

Types of participants

We include trials involving participants who are at risk of developing high altitude illness (AMS or HACE, HAPE). We include participants with and without a history of high altitude illness. We applied no age or gender restrictions.

Types of interventions

The published protocol for this review included all agents to prevent high altitude illness (Martí-Carvajal 2012). However we decided to split the topic into a series of three publications about the prevention of this condition (See Differences between protocol and review section). This is the first of the three and includes the

following six groups of the most widely recommended agents to prevent acute HAI:

1. Carbonic anhydrase inhibitors: Including acetazolamide and methazolamide.
 2. Steroids: Including budesonide, prednisolone and dexamethasone.
 3. Bronchodilator drugs: Including salmeterol, theophylline and montelukast.
 4. Selective inhibitor of phosphodiesterase type 5 (PDE5): Including taladafil and sildenafil.
 5. Calcium channel modulators: Including nifedipine and flunarizine.
 6. Non-steroidal anti-inflammatory drugs (NSAIDs) and other analgesics: Including aspirin, carbamazepine and ibuprofen.
- We include trials where the relevant medication was administered before beginning the ascent. We exclude trials using these drugs during or after the ascent.

Types of outcome measures

We modified the following outcome measures from the published protocol (Marti-Carvajal 2012). This is a departure from the protocol and it is explained in the [Differences between protocol and review](#) section.

Primary outcomes

1. Incidence of acute mountain sickness (AMS - as defined by each study) at any time.

Secondary outcomes

1. Incidence of high altitude pulmonary oedema (HAPE - as defined by each study) at any time.
2. Incidence of high altitude cerebral oedema (HACE - as defined by each study), at any time.
3. Incidence of adverse events in general, including paraesthesia, at any time.
4. Differences in HAI/AMS scores at high altitude. We analysed the differences between groups by any measure of AMS severity and between 0 and 48 hours at high altitude.

Search methods for identification of studies

We used the same search methods for the identification of studies, which are common to the three reviews included in this series.

Electronic searches

We searched the Cochrane Central Register of Controlled Trials (CENTRAL) (the Cochrane Library, January 2017); MEDLINE (OVID, 1966 to January 2017); Embase (OVID, 1980 to January 2017); LILACS (1982 to January 2017). We used the specific search terms listed below in combination with the Cochrane

highly sensitive search strategy for identifying randomized controlled trials (RCTs) in the *Cochrane Handbook for Systematic Reviews of Interventions* (Higgins 2011). [Appendix 4](#) to [Appendix 7](#) show the search strategies used in this set of reviews. We undertook the most recent search in January 2017.

Searching other resources

We also searched trials registries through the World Health Organization International Clinical Trials Registry Platform Search Portal (ICTRP) (see [Appendix 8](#)). We looked through the reference lists of the retrieved publications and review articles. We undertook the most recent search in January 2017.

Data collection and analysis

Data collection and analysis methods were common to the three reviews included in this series.

Selection of studies

Two review authors independently assessed each reference identified by the search against the inclusion criteria. We resolved any disagreements by discussion, and by consultation with a third review author as an arbiter if we could not reach agreement. We retrieved in full those references which appeared to meet the inclusion criteria for further independent assessment by the same three review authors.

Data extraction and management

We used a predefined form to extract the following data: eligibility criteria, demographics (age, gender, country), rate of ascent (metres/hour), final altitude reached (metres), AMS scale, design study, history of HAI, type of HAI, proposed intervention, and main outcomes, among others. See [Appendix 9](#) for details of the data extraction form. For eligible studies, two review authors extracted the data using the selected form. We resolved discrepancies through discussion or, if required, we involved a third review author. We entered data into Review Manager 5 software and checked them for accuracy.

Assessment of risk of bias in included studies

Three review authors independently assessed risks of bias for each study, using the criteria outlined in the *Cochrane Handbook for Systematic Reviews of Interventions* (Higgins 2011). We resolved any disagreement by discussion. We judged the methodological quality of each study using Cochrane's process for assessing risk of bias, a two-part tool that addresses the six specific domains: random sequence generation; allocation concealment; blinding of participants, personnel, and outcome assessors; incomplete outcome data; selective reporting; and other potential biases (Higgins

2011). The first part describes the risk of bias; the second part provides criteria for making judgements about the risk of bias from each of the six domains (Appendix 9). Based on this process we implemented a 'Risk of bias' worksheet to be filled out for each study. Two review authors independently assessed the risks of bias, resolving any disagreement through consultation with an additional review author. We display the results by creating a 'Risk of bias' graph and a 'Risk of bias' summary figure using RevMan 5.3 software, if appropriate. We present the risks of bias in the Results section. We also provided summary assessments of the risks of bias for each outcome within and across studies.

Measures of treatment effect

For dichotomous outcomes (such as incidence of AMS or HAPE), we show results as summary risk ratios (RRs) with 95% confidence intervals (CIs). For continuous outcomes (such as differences in AMS scores), we present the results as summary mean differences (MDs) or standardized mean differences (SMDs) as appropriate, with a 95% CI. Because we identified a considerable number of cross-over trials, we have included these studies separately and analysed this information using the criteria outlined in the *Cochrane Handbook for Systematic Reviews of Interventions-Chapter 16.4* (Elbourne 2002; Higgins 2011; Stedman 2011), specifically related to estimation of the Mantel-Haenzel odds ratio (OR) for paired outcomes.

Unit of analysis issues

The published protocol did not include consideration of any unit of analysis issues. However, our searches identified 12 cross-over studies and we included them in the analyses, but separately from the parallel studies. In brief, we used the methods recommended by Elbourne (Elbourne 2002; Stedman 2011). This is a departure from the protocol (Martí-Carvajal 2012) and is explained in the [Differences between protocol and review](#) section.

Dealing with missing data

For all outcomes, we carried out analyses, as far as possible, on an intention-to-treat (ITT) basis (i.e. we attempted to include all randomized participants in the denominator of the assessed groups in the analyses). Due to the fact that we included studies with missing information (especially standard deviations) or data not suitable for planned analyses, we followed the methods recommended by the *Cochrane Handbook for Systematic Reviews of Interventions-Chapter 16.1.3*. (Higgins 2011). In brief, we transformed median values and their interquartile ranges or range extracted from included studies to means and standard deviations according to Wan and colleagues (Hozo 2005; Wan 2014). This is a departure from the protocol (Martí-Carvajal 2012) and it is explained in the [Differences between protocol and review](#) section.

Assessment of heterogeneity

We used the I^2 statistic to measure statistical heterogeneity among the trials in each analysis. When we identified substantial heterogeneity, we explored it by prespecified subgroup analysis. The I^2 statistic describes the percentage of total variation across trials due to heterogeneity rather than sampling error (Higgins 2003). We considered a value for I^2 greater than 50% (Higgins 2011) to be statistically significant. We assessed the clinical and methodological diversity of the included studies in a comparison for sufficient homogeneity before choosing to estimate summary effect sizes.

Assessment of reporting biases

We assessed whether the review was subject to publication bias by using a funnel plot to graphically illustrate variability between trials. If we detected asymmetry, we planned to explore causes other than publication bias. We produced a funnel plot if we could include 10 or more RCTs in a comparison.

Data synthesis

We summarized the findings using the random-effects model (DerSimonian 1986). We carried out statistical analyses using Review Manager 5 (RevMan 5.3). We interpreted differences as important where the 95% confidence interval did not cross the value of no difference between groups. We also applied trial sequential analysis, as cumulative meta-analyses are at risk of producing random errors due to sparse data and repetitive testing of the accumulating data (Brok 2009; Wetterslev 2008). To minimize random errors, we calculated the required information size (i.e. the number of participants needed in a meta-analysis to detect or reject a certain intervention effect) (Wetterslev 2008). The required information size calculation also accounted for the heterogeneity or diversity present in the meta-analysis (Wetterslev 2008). In our meta-analysis, we based the diversity-adjusted required information size on the event proportion in the control group; assumption of a plausible risk ratio reduction (RRR) of 20% on the RR reduction observed in the included trials with low risk of bias; a risk of type I error of 5%; a risk of type II error of 20%; and the assumed diversity of the meta-analysis. We added the trials according to the year of publication, and if more than one trial had been published in a year, we added trials alphabetically according to the last name of the first trial author. On the basis of the required information size, we constructed trial sequential monitoring boundaries (Lan 1983; Thorlund 2009; Wetterslev 2008). These boundaries determine the statistical inference one may draw regarding the cumulative meta-analysis that has not reached the required information size; if the trial sequential monitoring boundary is crossed before the required information size is reached, firm evidence may perhaps be established and further trials may turn out to be superfluous. On the other hand, if the boundary is not crossed, it is most probably necessary to continue doing trials in order to detect or reject a certain intervention effect. This can be determined by assessing

if the cumulative Z-curve crosses the trial sequential boundaries. Furthermore, trial sequential analysis can test the futility before the required information size has been reached, i.e. trial sequential analysis provides an area of futility. If futility boundaries are crossed, then further trials may be unnecessary (CTU 2011). We conducted TSA using software from the Copenhagen Trial Unit (CTU 2011). This is a departure from the published protocol (Martí-Carvajal 2012). See the details in the [Differences between protocol and review](#) section.

Subgroup analysis and investigation of heterogeneity

We investigated heterogeneity by an informed clinical evaluation of each outcome, combining data only when clinically appropriate. We also investigated statistical heterogeneity using the I^2 statistic, as described above. For the primary outcomes, we considered subgroup analysis for the following factors, as appropriate:

1. Extreme altitude exposure versus high or very high exposure (high: 1500 to 3500 metres or 4921 to 11,842 feet; very high: 3500 to 5500 metres or 11,842 to 18,044 feet; and extreme: above 5500 metres or 18,044 feet) (Paralíkar 2010).
2. Presence or absence of people at high risk of HAI.
3. The presence or absence of significant pre-existing disease: cardiovascular diseases, chronic obstructive pulmonary disease (COPD), diabetes mellitus.

Sensitivity analysis

We performed a sensitivity analysis comparing the general results versus RCTs of high methodological quality (studies classified as having a 'low risk of bias' (Higgins 2011)). We chose only three core domains: generation of allocation sequence, incomplete outcome data, and selective reporting bias.

Summary of findings tables

We used the GRADE system (Guyatt 2008) to assess the quality of the body of evidence associated with primary outcomes (incidence of AMS, HAPE, HACE and adverse events), and we con-

structed three 'Summary of findings' tables using the GRADE profiler software for the three major comparisons in this review (acetazolamide versus placebo, budenoside versus placebo and dexamethasone versus placebo). The outcomes covered in these tables are the incidence of AMS, the incidence of HAPE, the incidence of HACE and adverse events ([Summary of findings for the main comparison](#); [Summary of findings 2](#); [Summary of findings 3](#)). The GRADE approach appraises the quality of a body of evidence based on the extent to which one can be confident that an estimate of effect or association reflects the item being assessed. Evaluation of the quality of a body of evidence considers within-study risk of bias, directness of the evidence, heterogeneity of the data, precision of effect estimates and risk of publication bias (Balslem 2011; Guyatt 2011a; Guyatt 2011b; Guyatt 2011c; Guyatt 2011d; Guyatt 2011e; Guyatt 2011f; Guyatt 2011g; Guyatt 2011h).

RESULTS


Description of studies

See [Characteristics of included studies](#) and [Characteristics of excluded studies](#).

Results of the search

We carried out the latest search strategies in January 2017 and identified 1280 references. After reviewing the references by title and abstract, we selected 173 of the citations to review as full texts (see [Figure 1](#)). After reading the articles, we included 64 studies and 4547 participants (distributed across 78 references), excluded 38 studies (distributed in 40 references), classified 12 as ongoing studies, and 12 as studies awaiting assessment (most of them due to full text not yet available). We also identified 31 additional studies focusing on other interventions not covered by this review. We will incorporate these in subsequent reviews in this series.

Figure 1. Study flow diagram.



Included studies

Twelve of 64 included studies are cross-over trials (18.7%) that we analysed separately (Fischer 2000a; Fischer 2004; Fulco 2006; Greene 1981; Johnson 1984; Larson 1982b; Luks 2007; Muza 2004; Rock 1989a; Rock 1989b; Rock 1989c; Subudhi 2011). Fifty-four trials were developed at high altitude (84%), and the remaining 11 were performed in hypobaric chambers (17.1%; Table 1; Baumgartner 2003; Fischer 2000a; Fischer 2004; Fulco 2006; Johnson 1984; Luks 2007; Muza 2004; Subudhi 2011; Rock 1989a; Rock 1989b; Rock 1989c).

Participants

The participants' ages ranged between 16 and 65 years. Nineteen of the studies included only men (29.6%; Table 1. Anonymous 1981; Basu 2002a; Basu 2002b; Baumgartner 2003; Fischer 2000a; Fischer 2000b; Fischer 2004; Hackett 1988; Hillenbrand 2006; Hussain 2001; Jain 1986; Johnson 1984; Ke 2013; Küpper 2008; Moraga 2007; Rock 1989a; Rock 1989b; Rock 1989c; Zheng 2014).

Eleven out of 64 studies included people at high risk of AMS, HAPE or HACE, due to a history of these conditions or comorbidities such as asthma (17.1%; Bartsch 1991; Bernhard 1994; Bernhard 1998; Burtcher 1998; Burtcher 2001; Burtcher 2014; Hohenhaus 1994; Maggiorini 2006; Mirrakhimov 1993; Sartori 2002; Wright 1983).

Setting

Nineteen of the studies were undertaken in the USA (29.6%); 17 were carried out in India (26.1%); and six out of 65 studies were carried out in South America (9.2%; Anonymous 1981; Bates 2011; Bernhard 1994; Bernhard 1998; Moraga 2007; Wang 2013). The remaining studies were carried out in other countries (Table 1).

Administration of intervention to prevent AMS

Twenty-four out of 64 studies provided the intervention between three and five days prior to the ascent (37.5%; Table 1), and 22 between one and two days prior (34.3%; Table 1). The remaining studies provided the intervention in other time intervals. Four trials did not provide information about this issue (ASCENT 2012; Hillenbrand 2006; SPACE 2011; Wright 2004). In 25% of the trials, the participants hiked to endpoint altitude (trekking), and 12 studies used a combination of means of transportation, including cars, trains, and cable-cars (18.7%; Table 1).

Altitude

Most of the included studies reached a final altitude of between 4001 and 5000 metres above sea level (59.3%; Table 1). The most frequent difference between the endpoint and the baseline altitude was 3001 to 4000 metres (35.9%; Table 1), followed by a difference of more than 4000 metres (28.1%). The most frequent durations for ascent were of less than five hours (14 studies, 21.8%; Table 1) and three days or more (14 studies, 21.8%; Table 1). Eighteen studies did not provide information about these issues (28.1%; ASCENT 2012; Burtcher 1998; Burtcher 2001; Basu 2002a; Faull 2015; Fulco 2006; HEAT 2010; Jain 1986; Johnson 1984; Luks 2007; Montgomery 1989; Muza 2004; PACE 2006; PHAIT 2004; Rock 1989a; Rock 1989b; Rock 1989c; Van Patot 2008).

Scale used to assess AMS

The most commonly-used scale used was the Lake Louise Score (23 trials, 35.9%), and the criterion to define AMS onset was a score three or more points in eight trials (12.5%; Table 1. Bates 2011; Burtcher 2014; Chen 2015; Muza 2004; PACE 2006; Parati 2013; Subudhi 2011; Wright 2004). In 19 studies, the criteria used to define the onset of AMS were unclear (29.6%; Anonymous 1981; Banderet 1977; Bartsch 1991; Basu 2002a; Bradwell 1986; Burki 1992; Faull 2015; Fischer 2000a; Fischer 2000b; Greene 1981; HEAT 2010; Hochapfel 1986; Jain 1986; Luks 2007; Mirrakhimov 1993; Sartori 2002; Wright 1983; Wang 2013; Zell 1988).

Funding

In 23 of the included studies, the source of funding was unclear (35.9%; Table 1), and only 19 of 64 studies declared their possible conflicts of interests (29.6%; Basnyat 2003; Basnyat 2008; Burtcher 1998; Burtcher 2014; Carlsten 2004; HEAT 2010; Hillenbrand 2006; Hohenhaus 1994; Ke 2013; Lipman 2012; Luks 2007; Maggiorini 2006; Muza 2004; Bernhard 1994; Parati 2013; PHAIT 2004; Subudhi 2011; Van Patot 2008; Wang 2013).

Excluded studies

We excluded 38 studies (40 references) from the review. Twenty-eight out of 38 were excluded for not focusing on HAI or AMS prevention (73.6%), but reported instead physiological or laboratory results related to altitude ascent. In eight studies, authors reported results for the treatment of HAI or AMS (21%). We excluded the remaining references for other reasons. Readers can find more information about this aspect in the [Characteristics of excluded studies](#).

Studies awaiting classification

We classified 12 studies (Dugas 1995; Ellsworth 1987; Furian 2016; Hefri 2014; Kasic 1991; Lee 2011; Pun 2014; Roncin 1996; Swenson 1997; Utz 1970; Wang 1998; Xiangjun 2014) as awaiting assessment. We were unable to obtain the full texts from the authors, the Anaesthesia, Critical and Emergency Care Cochrane Group (ACE) or the Iberoamerican Cochrane Centre. See [Characteristics of studies awaiting classification](#).

NCT01606527; NCT01682551; NCT01794078; NCT01993667; NCT02244437; NCT02450968; NCT02604173; NCT02811016; NCT02941510), given that we were only able to find them on trial registers, but we considered that they could be published shortly. See [Characteristics of ongoing studies](#).

Ongoing studies

We considered 12 additional studies to be ongoing (ChiCTR-TRC-13003319; ChiCTR-TRC-13003590; NCT00886912;

Risk of bias in included studies

We assessed the risks of bias for the studies across six domains. We provide a summary of our assessment of the methodological quality of included studies in [Figure 2](#) and [Figure 3](#).

Figure 2. Risk of bias graph: review authors' judgements about each risk of bias item presented as percentages across all included studies.




Figure 3. Risk of bias summary: review authors' judgements about each risk of bias item for each included study.

	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding of participants and personnel (performance bias)	Blinding of outcome assessment (detection bias)	Incomplete outcome data (attrition bias)	Selective reporting (reporting bias)	Other bias
Anonimus 1961	?	?	?	?	?	?	?
ASCENT 2012	?	?	?	?	?	?	?
Banderet 1977	?	?	?	?	?	?	?
Bartsch 1991	?	?	?	?	?	?	?
Basnyat 2003	?	?	?	?	?	?	?
Basnyat 2009	?	?	?	?	?	?	?
Blau 2003a	?	?	?	?	?	?	?
Blau 2002b	?	?	?	?	?	?	?
Bates 2011	?	?	?	?	?	?	?
Baumgartner 2003	?	?	?	?	?	?	?
Bernhard 1994	?	?	?	?	?	?	?
Bernhard 1996	?	?	?	?	?	?	?
Bradwell 1998	?	?	?	?	?	?	?
Burki 1992	?	?	?	?	?	?	?
Burtscher 1999	?	?	?	?	?	?	?
Burtscher 2001	?	?	?	?	?	?	?
Burtscher 2014	?	?	?	?	?	?	?
Carlsten 2004	?	?	?	?	?	?	?
Chen 2015	?	?	?	?	?	?	?
Chow 2005	?	?	?	?	?	?	?
Ellsworth 1991	?	?	?	?	?	?	?
Fault 2015	?	?	?	?	?	?	?
Fischer 2006a	?	?	?	?	?	?	?
Fischer 2006b	?	?	?	?	?	?	?
Fischer 2004	?	?	?	?	?	?	?
Fulco 2006	?	?	?	?	?	?	?
Oreene 1991	?	?	?	?	?	?	?
Hackett 1976	?	?	?	?	?	?	?
Hackett 1989	?	?	?	?	?	?	?
HEAT 2010	?	?	?	?	?	?	?
Hillebrand 2008	?	?	?	?	?	?	?
Hochapfel 1996	?	?	?	?	?	?	?
Hohenhaus 1994	?	?	?	?	?	?	?
Hussain 2001	?	?	?	?	?	?	?
Jain 1998	?	?	?	?	?	?	?
Johnson 1984	?	?	?	?	?	?	?
Kayser 2008	?	?	?	?	?	?	?
Ke 2013	?	?	?	?	?	?	?
Kupper 2008	?	?	?	?	?	?	?
Larson 1982a	?	?	?	?	?	?	?
Larson 1982b	?	?	?	?	?	?	?
Lipman 2012	?	?	?	?	?	?	?
Luke 2007	?	?	?	?	?	?	?
Maggiorni 2006	?	?	?	?	?	?	?
Mirakhimov 1993	?	?	?	?	?	?	?
Mordgomery 1999	?	?	?	?	?	?	?
Moraga 2007	?	?	?	?	?	?	?
Muca 2004	?	?	?	?	?	?	?
PACE 2008	?	?	?	?	?	?	?
Parab 2013	?	?	?	?	?	?	?
PHAT 2004	?	?	?	?	?	?	?
Rock 1987	?	?	?	?	?	?	?
Rock 1989a	?	?	?	?	?	?	?
Rock 1989b	?	?	?	?	?	?	?
Rock 1989c	?	?	?	?	?	?	?
Sartori 2002	?	?	?	?	?	?	?
SPACE 2011	?	?	?	?	?	?	?
Subudhi 2011	?	?	?	?	?	?	?
Van Pelt 2008	?	?	?	?	?	?	?
Wang 2013	?	?	?	?	?	?	?
Wright 1983	?	?	?	?	?	?	?
Wright 2004	?	?	?	?	?	?	?
Zell 1988	?	?	?	?	?	?	?
Zheng 2014	?	?	?	?	?	?	?

Allocation

The authors reported a valid method of randomization in 19 studies, (ASCENT 2012; Basnyat 2008; Bates 2011; Bernhard 1998; Chen 2015; Chow 2005; Ellsworth 1991; Faull 2015; HEAT 2010; Hillenbrand 2006; Jain 1986; Ke 2013; Larson 1982a; Lipman 2012; Maggiorini 2006; Moraga 2007; PHAIT 2004; Van Patot 2008; Zheng 2014), whereas this information was not clearly reported in the remaining studies (70.3%). Similarly, 14 studies undertook and reported random allocation concealment (Basnyat 2003; Basnyat 2008; Chen 2015; Chow 2005; Hillenbrand 2006; Lipman 2012; Maggiorini 2006; PACE 2006; PHAIT 2004; Rock 1989a; Rock 1989b; Rock 1989c; Wright 2004; Zheng 2014), and the information was absent from the remaining included studies (78.1%).

Blinding

Twenty-two studies reported blinding of participants and personnel (Basnyat 2003; Basnyat 2008; Bates 2011; Bernhard 1998; Bradwell 1986; Burtscher 2014; Chow 2005; Ellsworth 1991; Fulco 2006; Hackett 1976; Hillenbrand 2006; Hochapfel 1986; Ke 2013; Larson 1982a; Larson 1982b; Luks 2007; PACE 2006; Rock 1989a; Rock 1989b; Rock 1989c; Wang 2013; Zheng 2014). In two studies, we classified this domain as high risk (Banderet 1977; Chen 2015).

We considered the risk of detection bias to be low in 12 studies (Bartsch 1991; Chow 2005; Fulco 2006; Hackett 1976; Hillenbrand 2006; Maggiorini 2006; Rock 1987; Rock 1989a; Rock 1989b; Rock 1989c; Wright 1983; Zheng 2014), and unclear in the remaining studies (81.2%). In eight studies, we rated the risk of bias as low for both performance and detection bias (Chow 2005; Fulco 2006; Hackett 1976; Hillenbrand 2006; Rock 1989a; Rock 1989b; Rock 1989c; Zheng 2014).

Incomplete outcome data

Significant numbers of participants were lost or excluded from the final analysis of eight studies (Bartsch 1991; Basnyat 2003; HEAT 2010; Hillenbrand 2006; Johnson 1984; Luks 2007; PHAIT 2004; Subudhi 2011). Nine further studies presented unclear data (ASCENT 2012; Basu 2002a; Basu 2002b; Bradwell 1986; Faull 2015; Fischer 2000a; Hackett 1976; Hochapfel 1986; Jain 1986). In the studies with minimal attrition bias, we often found that the data analyses were undertaken on a per protocol basis, and we took this into account for data collection, including all the randomized participants in the denominators of the assessed groups.

Selective reporting

Reporting adverse events associated with the different types of interventions is fundamental to a complete assessment of their usefulness in clinical practice. We found that the majority of the studies did not report on adverse events associated with the classes of drugs commonly-used for prevention of AMS (such as paraesthesia) (73.4%; Banderet 1977; Bartsch 1991; Basnyat 2008; Basu 2002a; Basu 2002b; Bates 2011; Baumgartner 2003; Bernhard 1994; Bernhard 1998; Burki 1992; Burtscher 1998; Burtscher 2014; Carlsten 2004; Chen 2015; Ellsworth 1991; Faull 2015; Fischer 2000b; Fischer 2004; Fulco 2006; Hackett 1976; Hackett 1988; Hochapfel 1986; Hohenhaus 1994; Jain 1986; Kayser 2008; Küpper 2008; Larson 1982a; Larson 1982b; Lipman 2012; Luks 2007; Maggiorini 2006; Mirrakhimov 1993; Montgomery 1989; Moraga 2007; Muza 2004; Parati 2013; Rock 1987; Rock 1989a; Rock 1989b; Rock 1989c; Sartori 2002; SPACE 2011; Subudhi 2011; Van Patot 2008; Wang 2013; Wright 1983; Wright 2004). The remaining studies reported at least one adverse event related to the assessed intervention.

Other potential sources of bias

We found a possibility of industry bias in 29 studies, mainly related to the unclear role of the sponsors in the development of the study and the unknown effect of the first phase on cross-over trials in final results (Anonymous 1981; Basu 2002b; Bernhard 1994; Bradwell 1986; Burtscher 1998; Burtscher 2001; Fischer 2000a; Fischer 2000b; Fischer 2004; Fulco 2006; Greene 1981; HEAT 2010; Johnson 1984; Küpper 2008; Larson 1982a; Larson 1982b; Luks 2007; Mirrakhimov 1993; Montgomery 1989; Muza 2004; PACE 2006; PHAIT 2004; Rock 1987; Rock 1989a; Rock 1989b; Rock 1989c; Subudhi 2011; Wright 1983; Wright 2004). We identified no other potential sources of risk in the remaining studies.

Effects of interventions

See: [Summary of findings for the main comparison Acetazolamide compared with placebo for preventing high altitude illness](#); [Summary of findings 2 Budesonide compared with placebo for preventing high altitude illness](#); [Summary of findings 3 Dexamethasone compared with placebo for preventing high altitude illness](#)

See [Summary of findings for the main comparison](#); [Summary of findings 2](#); [Summary of findings 3](#).

GROUP 1: Carbonic anhydrase inhibitors

Comparison 1: carbonic anhydrase inhibitors: acetazolamide versus placebo

For this comparison, we included information from 28 parallel studies (2345 participants) (Anonymous 1981; Banderet 1977; Basnyat 2003; Basnyat 2008; Bradwell 1986; Burki 1992; Burtscher 2014; Carlsten 2004; Chow 2005; Ellsworth 1991; Faull 2015; Hackett 1976; HEAT 2010; Hillenbrand 2006; Hochapfel 1986; Hussain 2001; Jain 1986; Ke 2013; Larson 1982a; Mirrakhimov 1993; Moraga 2007; Parati 2013; PACE 2006; PHAIT 2004; SPACE 2011; Van Patot 2008; Wang 2013; Wright 2004).

All trials were performed in high mountain areas. Many of the studies administered acetazolamide or placebo between three and five days prior to ascent (13 out of 28; 46.4%) with doses of 500 mg/day (13 out of 28 studies, 46.4%; Anonymous 1981; Basnyat 2008; Bradwell 1986; Burki 1992; Chow 2005; Faull 2015; Hackett 1976; Hussain 2001; Moraga 2007; Parati 2013; PHAIT 2004; SPACE 2011; Wright 2004). For the assessment of AMS, the most widely-used scale was the Lake Louise Score (12 out of 28 studies, 42.8%) with scores of three or more with headache as a definition of AMS (4 out of 28 trials, 14.2%; Basnyat 2008; Carlsten 2004; Hillenbrand 2006; PHAIT 2004). Two studies involved people with a history of AMS, HAPE or HACE (Burtscher 2014; Mirrakhimov 1993).

Most of the studies reached altitudes of between 3001 to 4000 metres (Bradwell 1986; Burki 1992; Burtscher 2014; Carlsten 2004; Ellsworth 1991; Faull 2015; Jain 1986; Ke 2013; Larson 1982a; Moraga 2007; Wang 2013; Wright 2004). All but four studies included very high altitude exposure (i.e. 3500 to 5500 metres; Hochapfel 1986; Jain 1986; Mirrakhimov 1993; Wright

2004).


Seven studies did not provide any information about any of the outcomes assessed in this review (Banderet 1977; Burki 1992; Burtscher 2014; Faull 2015; Hochapfel 1986; Jain 1986; Wang 2013). Because Carlsten 2004 and PACE 2006 evaluated two different groups that had been administered doses of acetazolamide, we included this information for the following analyses. Finally, in Carlsten 2004 two different definitions of HAI were provided and we chose information according to the second definition (Lake Louise AMS score of three or more with headache).

In addition, we analysed information from five cross-over trials (Fischer 2004; Fulco 2006; Greene 1981; Larson 1982b; Subudhi 2011) with a total of 54 participants. Fischer 2004 only reported medians for scores of AMS, precluding the inclusion of this information in the following analysis.

Primary outcome 1: incidence of acute mountain sickness (AMS)


Sixteen parallel studies provided information about this outcome (Basnyat 2003; Basnyat 2008; Carlsten 2004; Chow 2005; Hackett 1976; HEAT 2010; Hillenbrand 2006; Larson 1982a; Mirrakhimov 1993; Moraga 2007; Parati 2013; PACE 2006; PHAIT 2004; SPACE 2011; Van Patot 2008; Wright 2004), registering a total of 391 events of acute mountain sickness (Incidence of AMS: 16.9%). The risk ratio (RR) for acute mountain sickness, comparing acetazolamide to placebo, was 0.47 (95% confidence interval (CI) 0.39 to 0.56; $I^2 = 0\%$; 16 trials, 2301 participants; Analysis 1.1; Figure 4).

Figure 4. Forest plot of comparison: I Carbonic anhydrase inhibitors: acetazolamide versus placebo, outcome: I.I Incidence of acute mountain sickness.



We downgraded the quality of evidence from high to moderate, due to unclear risks of selection, detection, and performance bias in most of the included studies (See [Summary of findings for the main comparison](#)). In addition, when we considered the dosage of acetazolamide, we found a non-statistically significant reduction in the risk of HAI in all groups (test for subgroup differences: Chi² = 4.55, df = 3; P = 0.21; I² = 34.0%. The RR for 250 to 255 mg is 0.60 (95% CI 0.39 to 0.94; I² = 14%; 4 trials, 855 participants). The RR for 500 mg is 0.48 (95% CI 0.38 to 0.61; I² = 0%; 8 trials, 1111 participants). The RR for 750 mg is 0.33 (95% CI 0.18 to 0.62; I² = 0%; 2 trials, 80 participants). The funnel plot did not show data asymmetry related to sample size (Figure 5).

Figure 5. Funnel plot of comparison: I Carbonic anhydrase inhibitors: acetazolamide versus placebo, outcome: I.I Incidence of acute mountain sickness.



Regarding sensitivity analyses, only one study was at low risk of bias in the three core domains selected in the Methods section (Chow 2005). For our subgroup analyses, only one study includes an extreme altitude exposure (Wright 2004), and another includes people at high risk of HAI (Mirrakhimov 1993). In addition, two cross-over studies (Fulco 2006; Larson 1982b) found four events of acute mountain sickness (total incidence of AMS = 16.6%). The odds ratios ranged from 1 to 4.3. The pooled odds ratio for AMS, comparing acetazolamide to placebo, was 2.26 (95% CI 0.54 to 9.40; $I^2 = 56\%$), showing no effect of acetazolamide in the onset of HAI, but with considerable heterogeneity.

Secondary outcome 1: incidence of high altitude pulmonary oedema (HAPE)

Seven parallel studies (1138 participants) evaluated the incidence of altitude pulmonary oedema (Basnyat 2003; Basnyat 2008; Burki 1992; Chow 2005; Ke 2013; PHAIT 2004; SPACE 2011), but they did not find any events to report (Analysis 1.2). We downgraded the quality of evidence from high to moderate due to unclear risks of selection, detection, and performance bias (See

[Summary of findings for the main comparison](#)).

Secondary outcome 2: incidence of high altitude cerebral oedema (HACE)

Six parallel studies evaluated the incidence of altitude pulmonary oedema (Basnyat 2003; Basnyat 2008; Chow 2005; Ke 2013; PHAIT 2004; SPACE 2011), but only one event was reported (incidence of HACE = 0.08%). The RR for HACE, comparing acetazolamide to placebo, was 0.32 (95% CI 0.01 to 7.48; 6 trials, 1126 participants; Analysis 1.3). We downgraded the quality of evidence from high to moderate due to unclear risks of selection, detection, and performance bias (See [Summary of findings for the main comparison](#)).

Secondary outcome 3: incidence of adverse events

Five parallel studies provide information about paraesthesias (Anonymous 1981; Basnyat 2003; Chow 2005; PACE 2006; PHAIT 2004), for 279 events (incidence of paraesthesia = 35.3%).

The RR for paraesthesia, comparing acetazolamide to placebo, was 5.53 (95% CI 2.81 to 10.88; $I^2 = 60\%$; 789 participants; [Analysis 1.4](#)). This heterogeneity is reduced to 0% when the dosage of acetazolamide is taken into account (RR from 3.09 to 12.63 by dose; [Analysis 1.4](#)). We downgraded the quality of evidence from high to low due to unclear risks of selection, performance, and detection bias, as well as inconsistency (See [Summary of findings for the main comparison](#)).

One study ([Hillenbrand 2006](#)) evaluated the incidence of side effects in general, including paraesthesia and numbness. Sixty-eight side effects were reported (incidence of side effects 17%). The risk of side effects, comparing acetazolamide to placebo, was 2.19 (95% CI 1.36 to 3.53) under intention-to-treat analysis. However, under per-protocol analysis, the risk was 2.20 (95% CI 1.55 to 3.12). When the missing subjects were considered as cases of adverse events in both arms, the estimated risk was 1.15 (95% CI 1.08 to 1.23). We downgraded the quality of evidence from high to low due to these high levels of attrition bias (See [Summary of findings for the main comparison](#)). Another study ([HEAT 2010](#)) evaluated the incidence of major events, including drug reactions and gastrointestinal bleeding. However, authors found no major events to report. Finally, in [Zell 1988](#) the authors reported the incidence of numbness in fingers, with six events in 32 participants.

One cross-over study reported the incidence of tingling ([Greene 1981](#); 24 participants). The estimated OR for this adverse event, comparing acetazolamide to placebo, was 1.44 (95% CI 0.78 to 2.68).

Secondary outcome 4: differences in HAI/AMS scores

Six parallel studies provide information about scores for AMS ([Carlsten 2004](#); [Chow 2005](#); [Hussain 2001](#); [Hillenbrand 2006](#); [Moraga 2007](#); [Wright 2004](#)). [Carlsten 2004](#) reported the scores for two doses of acetazolamide (250 mg and 500 mg) and compared them to a single common placebo group. To avoid double counting, we have presented the results as dosing subgroups only ([Analysis 1.5](#)). Pooling the data for all sets produced a heterogeneous effect estimate ($I^2 = 80.4\%$). The standardized mean difference between acetazolamide and placebo was 0.19 for doses of 250 mg/day (95% CI 0.01 to 0.37; $I^2 = 0\%$; 434 participants; [Analysis 1.5](#)). In contrast, the standardized mean difference between acetazolamide and placebo was -0.57 for doses of 500 mg/


day, but with considerable heterogeneity (95% CI -1.20 to 0.07; $I^2 = 72\%$; 92 participants; [Analysis 1.5](#)).

In addition, two cross-over studies reported differences in AMS scores, ranging from 1 to -2.7 ([Fulco 2006](#); [Subudhi 2011](#); 52 participants). The mean difference for these scores, comparing acetazolamide to placebo, was -1.25 (95% CI -4.79 to 2.29), but with considerable heterogeneity ($I^2 = 78\%$).

Trial sequential analysis for acetazolamide versus placebo

Trial sequential analysis of oral acetazolamide at any dose versus placebo for prevention of acute mountain sickness is based on the diversity-adjusted required information size (DARIS) of 2396 participants. We calculated this DARIS based upon a proportion of participants with acute mountain sickness of 23.3% in the control group; a RRR of 20% in the experimental intervention group; an alpha of 5%; a beta of 20%; and a diversity of 0%. The cumulative Z-curve (blue line) crossed the upper conventional alpha of 5% and the upper trial sequential alpha-spending monitoring boundaries, showing that we have robust data for significant efficacy ([Figure 6](#)). Likewise, trial sequential analysis of oral acetazolamide at 500 mg dose versus placebo for prevention of acute mountain sickness is based on a DARIS of 1759 participants. We calculated this DARIS based upon a proportion of participants with acute mountain sickness of 29.5% in the control group; a RRR of 20% in the experimental intervention group; an alpha of 5%; a beta of 20%; and a diversity of 0%. The cumulative Z-curve (blue line) crossed the upper conventional alpha of 5% and the upper trial sequential alpha-spending monitoring boundaries, showing that we have robust data for significant efficacy. Finally, TSA of oral acetazolamide at 250 mg dose versus placebo for prevention of acute mountain sickness is based on a DARIS of 1777 participants. We calculated this DARIS based upon a proportion of participants with acute mountain sickness of 13.1% in the control group; a RRR of 35% in the experimental intervention group; an alpha of 5%; a beta of 20%; and a diversity of 19%. The cumulative Z-curve (blue line) twice crossed twice the upper conventional alpha of 5%, but it did not cross the upper trial sequential alpha-spending monitoring boundaries, indicating that new randomized controlled trials are needed. Accordingly, after only 48.1% (855/1777) of the DARIS had been attained, we were able to reject an intervention effect of 35% or larger.

Figure 6. Trial sequential analysis on prevention of acute mountain illness in 16 oral acetazolamide at any dose vs placebo trials



Comparison 2: carbonic anhydrase inhibitors: acetazolamide 250 mg versus acetazolamide 500 mg

For this comparison, we analysed information from one study ([Carlsten 2004](#)) with 22 participants. This trial was carried out in the high mountain areas of Nepal, reaching a maximum altitude of 3630 metres.

Primary outcome 1: incidence of acute mountain sickness (AMS)

[Carlsten 2004](#) did not identify any events of acute mountain sickness.

Secondary outcome 1: incidence of high altitude pulmonary oedema (HAPE)

We found no information about this outcome in the included study.

Secondary outcome 2: incidence of high altitude cerebral oedema (HACE)

We found no information about this outcome in the included study.

Secondary outcome 3: incidence of adverse events

We found no information about this outcome in the included study.

Secondary outcome 4: differences in HAI or AMS scores

[Carlsten 2004](#) provided information about differences in AMS scores. The mean difference for these scores, comparing 250 mg/day of acetazolamide versus 500 mg/day of acetazolamide, was 0.76 (95% CI -0.16 to 1.68).

Comparison 3: carbonic anhydrase inhibitors: acetazolamide 750 mg versus acetazolamide 250mg

For this comparison, we analysed information from one study ([PACE 2006](#)) with 156 participants. This study was carried out in high mountain areas of Nepal, reaching a maximum altitude of 4928 meters.

Primary outcome 1: incidence of acute mountain sickness (AMS)

The authors of [PACE 2006](#) found 15 events of acute mountain sickness (incidence of AMS: 9.61%). The RR for acute mountain sickness, comparing 750 mg/day versus 250 mg/day of acetazolamide, was 0.60 (95% CI 0.22 to 1.61).

Secondary outcome 1: incidence of high altitude pulmonary oedema (HAPE)

We found no information about this outcome in the included study.

Secondary outcome 2: incidence of high altitude cerebral oedema (HACE)

We found no information about this outcome in the included study.

Secondary outcome 3: incidence of adverse events

Authors of [PACE 2006](#) reported information about paraesthesia, finding 117 events (incidence of paraesthesia: 75%). The RR for paraesthesias, comparing 750 mg/day versus 250 mg/day of acetazolamide, was 1.34 (95% CI 1.11 to 1.63).

Secondary outcome 4: differences in HAI/AMS scores

We found no information about this outcome in the included study.

Group 2: Steroids**Comparison 1: steroids: budesonide versus placebo**

For this comparison, we analysed the information from two studies ([Chen 2015](#); [Zheng 2014](#)) with 132 participants. Researchers administered 200 µg of inhaled budesonide twice daily in both studies. Both studies were carried out in China, reaching a maximum altitude of between 3700 to 3900 metres.

Primary outcome 1: incidence of acute mountain sickness (AMS)

Both studies provide information about the incidence of acute mountain sickness and found 45 events (incidence of AMS = 34%). The RR for AMS, comparing budesonide to placebo, was 0.37 (95% CI 0.23 to 0.61; $I^2 = 0\%$; [Analysis 2.1](#)). We downgraded the quality of evidence from high to low, due to a high risk of performance bias, as well as imprecision issues (See [Summary of findings 2](#)).

Secondary outcome 1: incidence of high altitude pulmonary oedema (HAPE)

We found no information about this outcome in the included studies.

Secondary outcome 2: incidence of high altitude cerebral oedema (HACE)

We found no information about this outcome in the included studies.

Secondary outcome 3: incidence of adverse events

[Chen 2015](#) assessed the incidence of side effects in general in all participants and did not find any events. We downgraded the quality of evidence from high to very low, due to a high risk of performance bias, as well as imprecision issues (See [Summary of findings 2](#)). Likewise, [Zheng 2014](#) evaluated the onset of persistent belching but did not find any affected participants. We downgraded the quality of evidence from high to low, due to imprecision issues (See [Summary of findings 2](#)).

Secondary outcome 4: differences in HAI/AMS scores

We found no information about this outcome in the included studies.

Comparison 2: steroids: dexamethasone versus placebo

For this comparison, we analysed the information from six studies in high mountain areas ([Bernhard 1994](#); [Hackett 1988](#); [Hussain 2001](#); [Montgomery 1989](#); [Rock 1987](#); [Zell 1988](#)), with a total of 205 participants. Two studies were carried out in the USA ([Hackett 1988](#); [Montgomery 1989](#)), two in Nepal ([Rock 1987](#); [Zell 1988](#)), and one each in Pakistan ([Hussain 2001](#)) and Bolivia ([Bernhard 1994](#)). [Hussain 2001](#) and [Montgomery 1989](#) included only men. All studies used scales other than the Lake Louise Score. [Bernhard 1994](#) included 40% of participants with previous AMS, and the altitude reached was classified as extreme (more than 5000 metres). Two studies administered 16 mg of dexamethasone ([Montgomery 1989](#); [Rock 1987](#)), and most studies administered it during one to two days ([Montgomery 1989](#); [Rock 1987](#); [Zell 1988](#)).

[Montgomery 1989](#) included the use of dexamethasone versus placebo at two different altitudes in two separate participant groups and the data for each has been presented separately ([Montgomery 1989](#) (2,700m) and [Montgomery 1989](#) (2,050m)). [Bernhard 1994](#) provided two definitions for AMS, but only one (modified Environmental Symptoms Questionnaire (ESQ) = 3 cerebral symptoms, one with intensity ≥ 2) provided information for further analyses. Data from [Bernhard 1994](#), [Hackett 1988](#) and [Hussain 2001](#) about AMS scores were provided as medians and standard errors, which needed transformation for the corresponding analyses (See [Appendix 10](#)).

We also analysed information from five cross-over studies ([Johnson 1984](#); [Rock 1989a](#); [Rock 1989b](#); [Rock 1989c](#); [Subudhi 2011](#)) with a total of 53 participants. The [Rock 1989](#) study provided information for three different doses of dexamethasone, and we extracted and analysed the data separately ([Rock 1989a](#); [Rock 1989b](#); [Rock 1989c](#)).

Primary outcome 1: incidence of acute mountain sickness (AMS)

Four parallel studies provided information about the incidence of acute mountain sickness (Bernhard 1994; Hackett 1988; Montgomery 1989; Rock 1987), and found a total of 60 events (incidence of AMS = 34.09%). The RR for AMS, comparing dexamethasone versus placebo, was 0.60 (95% CI 0.36 to 1.00; $I^2 = 39\%$; 176 participants; Analysis 3.1). We downgraded the quality of evidence from high to low, due to unclear risks of selection, performance, and detection bias, as well as imprecision issues (See Summary of findings 3). We found no numerical information about this outcome in the included cross-over studies. In Subudhi 2011 the authors reported six instances of AMS, but with no information on the number in each group.

Regarding sensitivity analyses, none of the studies included in this comparison present low risk of bias in all the three domains previously selected. Bernhard 1994 was the only study carried out at extreme altitude, and including a high-risk population. Excluding this study from these analyses modified the pooled RR from 0.60 to 0.58, but increased the heterogeneity from 39% to 56%.

Secondary outcome 1: incidence of high altitude pulmonary oedema (HAPE)

We found no information about this outcome in the included studies.

Secondary outcome 2: incidence of high altitude cerebral oedema (HACE)

We found no information about this outcome in the included studies.

Secondary outcome 3: incidence of adverse events

Bernhard 1994 assessed the incidence of adverse events in general, but found no events to report. Likewise, Zell 1988 evaluated the onset of numbness in participants, but they too found no cases to report. We downgraded the quality of evidence from high to very low, due to unclear risks of selection, performance and detection bias, as well as imprecision issues (See Summary of findings 3). From the cross-over studies, Johnson 1984 found one event of dyspepsia for this comparison (total incidence of dyspepsia = 6.25%). The RR for dyspepsia, comparing dexamethasone to placebo was 1.36 (95% CI 0.40 to 4.60).

Secondary outcome 4: differences in HAI/AMS scores

Three parallel studies provide information about AMS scores (Bernhard 1994; Hackett 1988; Hussain 2001). The standardized mean difference for these scores, comparing dexamethasone to placebo, was -0.46 (95% CI -1.21 to 0.29; $I^2 = 38\%$; 50 participants; Analysis 3.2). We downgraded the quality of evidence from

high to very low, due to unclear risks of selection, performance and detection bias, as well as imprecision issues (See Summary of findings 3). Five cross-over studies reported information about this outcome (Johnson 1984; Rock 1989a; Rock 1989b; Rock 1989c; Subudhi 2011). Mean differences ranged from -2.7 to 0.82 units. The MD for AMS scores, comparing dexamethasone to placebo, was -0.63 (95% CI -1.7 to 0.44), but with extreme heterogeneity ($I^2 = 99\%$).

Trial sequential analysis for dexamethasone versus placebo

Trial sequential analysis of dexamethasone versus placebo for prevention of acute mountain sickness is based on the diversity-adjusted required information size (DARIS) of 517 participants. We calculated this DARIS based upon a proportion of participants with acute mountain illness of 44.9% in the control group; a RRR of 35% in the experimental intervention group; an alpha of 5%; a beta of 20%; and a diversity of 43%. After the fifth trial, the cumulative Z-curve (blue line) crossed the upper conventional alpha of 5%, but it did not cross the upper trial sequential alpha-spending monitoring boundaries. Accordingly, after only 34% (176/517) of the DARIS had been attained, we were able to reject an intervention effect of 35% or larger, indicating that new randomized controlled trials are needed.

Comparison 3: steroids: prednisolone versus placebo

For this comparison, we analysed the information from one study (Basu 2002b) with 40 participants. However, this study did not provide information about any of the outcomes selected for this review.

Group 3: Bronchodilators**Comparison 1: bronchodilator drugs: salmeterol versus placebo**

For this comparison, we analysed the information from one study (Sartori 2002) with 37 participants. Researchers administered 125 mg of inhaled salmeterol twice daily. This study was carried out in Nepal, reaching a maximum altitude of 4559 metres; all participants were susceptible to HAPE.

Primary outcome 1: incidence of acute mountain sickness (AMS)

We found no information about this outcome in the included study.

Secondary outcome 1: incidence of high altitude pulmonary oedema (HAPE)

[Sartori 2002](#) provided information about the incidence of high-altitude pulmonary oedema, with 20 events (incidence of HAPE = 54.05%). The RR for HAPE, comparing salmeterol to placebo, was 0.45 (95% CI 0.22 to 0.92; 37 participants).

Secondary outcome 2: incidence of high altitude cerebral oedema (HACE)

We found no information about this outcome in the included study.

Secondary outcome 3: incidence of adverse events

We found no information about this outcome in the included study.

Secondary outcome 4: differences in HAI/AMS scores

[Sartori 2002](#) provided information about AMS scores. The mean difference for these scores, comparing salmeterol to placebo, was -5.70 (95% CI -8.50 to -2.90; 37 participants).

Comparison 2: bronchodilators drugs: theophylline versus placebo

For this comparison, we identified two parallel studies with at least 20 participants ([Fischer 2000a](#); [Küpper 2008](#)). The number of participants in [Fischer 2000a](#) was unclear, and this precludes the use of this study in further analyses. In addition, we analysed information from two cross-over studies ([Fischer 2000b](#); [Fischer 2004](#)) with a total of 24 participants. However, in [Fischer 2004](#) the authors only provided information for AMS scores as medians, precluding the inclusion of this information in further analyses.

Primary outcome 1: incidence of acute mountain sickness (AMS)

Only [Küpper 2008](#) provided information about the incidence of acute mountain sickness, with 12 events (incidence of AMS = 60%). The RR for AMS, comparing theophylline to placebo, was 0.71 (95% CI 0.34 to 1.50; 20 male participants).

Secondary outcome 1: incidence of high altitude pulmonary oedema (HAPE)

We found no information about this outcome in the included studies.

Secondary outcome 2: incidence of high altitude cerebral oedema (HACE)

We found no information about this outcome in the included studies.

Secondary outcome 3: incidence of adverse events

We found no information about this outcome in the included studies.

Secondary outcome 4: differences in HAI/AMS scores

Only [Küpper 2008](#) provided information about AMS scores for the parallel studies. The standardized mean difference for these scores, comparing theophylline to placebo, was -0.18 (95% CI -1.38 to 1.02; 20 participants). Of the cross-over studies, only [Fischer 2000b](#) reported information about scores for AMS. The mean difference between theophylline and placebo was -1.50 (95% CI -2.25 to -0.75).

Comparison 3: bronchodilator drugs: montelukast versus placebo

For this comparison, we analysed information from two cross-over studies ([Luks 2007](#); [Muza 2004](#)) with a total of 22 participants. [Muza 2004](#) provided two definitions of AMS (Lake Louise Scale ≥ 3 and ESQ AMS-C Score ≥ 0.7) and we selected the first one to include in analyses.

Primary outcome 1: incidence of acute mountain sickness (AMS)

[Muza 2004](#) found 14 events of acute mountain sickness (incidence of AMS = 58.3%). The odds ratio for AMS, comparing acetazolamide to placebo, was 1.47 (95% CI 0.61 to 3.55; 22 participants).

Secondary outcome 1: incidence of high altitude pulmonary oedema (HAPE)

We found no information about this outcome in the included studies.

Secondary outcome 2: incidence of high altitude cerebral oedema (HACE)

We found no information about this outcome in the included studies.

Secondary outcome 3: incidence of adverse events

We found no information about this outcome in the included studies.

Secondary outcome 4: differences in HAI/AMS scores

Both studies reported information about scores for AMS. Mean differences between montelukast and placebo ranged between 1.1 and -1.4. The mean difference between montelukast and placebo was -0.08 (95% CI -2.53 to 2.36; $I^2 = 81\%$) but with considerable heterogeneity.

Group 4: Selective inhibitors of phosphodiesterase-5

Comparison 1: selective inhibitors of phosphodiesterase-5: tadalafil versus placebo

For this comparison, we analysed the information from one study (Maggiorini 2006) with 19 participants. The dosage of tadalafil used was 20 mg/day. This study was carried out in Kenya, reaching a maximum altitude of 4559 metres. All participants had a history of HAPE.

Primary outcome 1: incidence of acute mountain sickness (AMS)

Maggiorini 2006 provided information about the incidence of acute mountain sickness, with 16 events (incidence of AMS = 84.2%). The RR for AMS, comparing tadalafil to placebo, was 0.90 (95% CI 0.61 to 1.32; 29 participants).

Secondary outcome 1: incidence of high altitude pulmonary oedema (HAPE)

Maggiorini 2006 provided information about the incidence of altitude pulmonary oedema, with eight events (incidence of HAPE = 42.1%). The RR for HAPE, comparing tadalafil to placebo, was 0.13 (95% CI 0.02 to 0.85; 29 participants).

Secondary outcome 2: incidence of high altitude cerebral oedema (HACE)

We found no information about this outcome in the included study.

Secondary outcome 3: incidence of adverse events

We found no information about this outcome in the included study.

Secondary outcome 4: differences in HAI/AMS scores

We found no information about this outcome in the included study.

Comparison 2: selective inhibitors of phosphodiesterase-5: sildenafil citrate versus placebo

For this comparison, we analysed the information from one study (Bates 2011) with 62 participants. The dosage of sildenafil citrate used was 150 mg/day. This study was carried out in Chile, reaching a maximum altitude of 5200 metres. Data about AMS scores were provided as medians and interquartile ranges, and we transformed them for further analyses (See Appendix 10).

Primary outcome 1: incidence of acute mountain sickness (AMS)

Bates 2011 provided information about the incidence of acute mountain sickness, with 39 events (incidence of AMS = 62.9%). The RR for AMS, comparing sildenafil citrate to placebo, was 1.31 (95% CI 0.91 to 1.89; 62 participants).

Secondary outcome 1 risk of altitude pulmonary oedema

We found no information about this outcome in the included study.

Secondary outcome 2: incidence of high altitude cerebral oedema (HACE)

We found no information about this outcome in the included study.

Secondary outcome 3: incidence of adverse events

We found no information about this outcome in the included study.

Secondary outcome 4: differences in HAI/AMS scores

Bates 2011 provided information about AMS scores. The standardized mean difference for these scores, comparing sildenafil to placebo, was -2.41 (95% CI -3.95 to -0.87; 62 participants).

Group 5: Calcium channel modulators

Comparison 1: calcium channel modulators: nifedipine versus placebo

For this comparison, we analysed the information from two studies (Bartsch 1991; Hohenhaus 1994) with a total of 48 participants. Both studies used 60 mg/day of nifedipine. Bartsch 1991 was carried out in Nepal, reaching a maximum altitude of 4559 metres, while Hohenhaus 1994 was carried out in Italy and reached the same maximum altitude. All of the participants in Bartsch 1991 had a history of HAPE, and most of the participants in Hohenhaus 1994 had susceptibility to AMS.

Primary outcome 1: incidence of acute mountain sickness (AMS)

[Hohenhaus 1994](#) provided information about the incidence of acute mountain sickness, with 17 events (incidence of AMS = 62.9%). The RR for AMS, comparing nifedipine to placebo, was 1.04 (95% CI 0.58 to 1.87; 27 participants).

Secondary outcome 1: incidence of high altitude pulmonary oedema (HAPE)

[Bartsch 1991](#) provided information about the incidence of high altitude pulmonary oedema, with eight events (incidence of HAPE = 38.09%). The RR for HAPE, comparing nifedipine to placebo, was 0.16 (95% CI 0.02 to 1.06; 21 participants).

Secondary outcome 2: incidence of high altitude cerebral oedema (HACE)

We found no information about this outcome in the included studies.

Secondary outcome 3: incidence of adverse events

We found no information about this outcome in the included studies.

Secondary outcome 4: differences in HAI/AMS scores

Both included studies provided information about AMS scores ([Bartsch 1991](#); [Hohenhaus 1994](#)). Mean differences ranged from -1.25 to 0.07. The standardized mean difference for these scores, comparing nifedipine to placebo, was -0.56, (95% CI -1.85 to 0.74; $I^2 = 78%$; 48 participants; [Analysis 4.1](#)), but with considerable heterogeneity.

Comparison 2: calcium channel modulators: flunarizine versus placebo

For this comparison, we analysed the information from one study ([Baumgartner 2003](#)) with 20 participants. [Baumgartner 2003](#) used a hypobaric chamber to assess the effectiveness of 10 mg of flunarizine at 4559 metres.

Primary outcome 1: incidence of acute mountain sickness (AMS)

[Baumgartner 2003](#) provided information about the incidence of acute mountain sickness and found 14 events (incidence of AMS = 70%). The RR for AMS, comparing flunarizine to placebo, was 1.00 (95% CI 0.56 to 1.78; 20 participants).

Secondary outcome 1: incidence of high altitude pulmonary oedema (HAPE)

We found no information about this outcome in the included study.

Secondary outcome 2: incidence of high altitude cerebral oedema (HACE)

We found no information about this outcome in the included study.

Secondary outcome 3: incidence of adverse events

We found no information about this outcome in the included study.

Secondary outcome 4: differences in HAI/AMS scores

We found no information about this outcome in the included study.

Group 6: non-steroidal anti-inflammatory drugs (NSAIDs)**Comparison 1: non-steroidal anti-inflammatory drugs (NSAIDs) and other analgesics: aspirin versus placebo**

For this comparison, we analysed the information from two studies ([Burtscher 1998](#); [Burtscher 2001](#)) with a total of 60 participants. Both studies focused on headache at altitude, using a headache score to evaluate its onset. Aspirin 320 mg was used as a prophylaxis, given from one to two hours beforehand; both studies reached a maximum altitude of 2880 metres.

Primary outcome 1: incidence of acute mountain sickness (AMS)

Both studies provided information about the incidence of acute mountain sickness ([Burtscher 1998](#); [Burtscher 2001](#)), and found a total of 31 events (incidence of AMS = 51.6%). RRs ranged from 0.13 to 0.60. The RR for AMS, comparing aspirin to placebo, was 0.35 (95% CI 0.06 to 1.95; $I^2 = 68%$; 60 participants; [Analysis 5.1](#)), but with considerable heterogeneity.

Secondary outcome 1: incidence of high altitude pulmonary oedema (HAPE)

We found no information about this outcome in the included studies.

Secondary outcome 2: incidence of high altitude cerebral oedema (HACE)

We found no information about this outcome in the included studies.

Secondary outcome 3: incidence of adverse events

[Burtscher 2001](#) assessed the incidence of major adverse events in general, but did not find any events to report.

Secondary outcome 4: differences in HAI/AMS scores

We found no information about this outcome in the included studies.

Comparison 2: non-steroidal anti-inflammatory drugs (NSAIDs) and other analgesics: ibuprofen versus placebo

For this comparison, we analysed the information from three studies ([ASCENT 2012](#); [HEAT 2010](#); [Lipman 2012](#)), with a total of 598 participants. Only [ASCENT 2012](#) and [Lipman 2012](#) provided a clear definition to determine the onset of AMS (Lake Louise AMS score ≥ 3 with headache). Ibuprofen dosage ranged from 600 to 1800 mg. [ASCENT 2012](#) and [HEAT 2010](#) were developed in Nepal, reaching a maximum altitude of 4928 metres, while [Lipman 2012](#) was developed in the USA, reaching a maximum altitude of 3810 metres. None of these studies included high-risk populations.

Primary outcome 1: incidence of acute mountain sickness (AMS)

Three studies provided information about the incidence of acute mountain sickness ([ASCENT 2012](#); [HEAT 2010](#); [Lipman 2012](#)), and found a total of 154 events (incidence of AMS = 25.7%). The RR for AMS, comparing ibuprofen to placebo, was 0.64 (95% CI 0.49 to 0.82; $I^2 = 0\%$; 598 participants; [Analysis 6.1](#)). Regarding sensitivity analyses, none of the included studies in this comparison were at low risk of bias in the three previously selected domains. Likewise, all three studies were developed at very high altitude and none of them included a population at high risk of developing HAI/AMS.

Secondary outcome 1: incidence of high altitude pulmonary oedema (HAPE)

[ASCENT 2012](#) evaluated the incidence of altitude pulmonary oedema, but did not find any events to report.

Secondary outcome 2: incidence of high altitude cerebral oedema (HACE)

[ASCENT 2012](#) evaluated the incidence of altitude cerebral oedema, but did not find any events to report.

Secondary outcome 3: incidence of adverse events

[HEAT 2010](#) assessed the incidence of major adverse events in general, but did not find any events to report. The authors of [ASCENT 2012](#) reported one event of black stools in the ibuprofen group.

Secondary outcome 4: differences in HAI/AMS scores

We found no information about this outcome in the included studies.

Trial sequential analysis for ibuprofen versus placebo

Trial sequential analysis of oral ibuprofen at any dose versus placebo for prevention of acute mountain sickness is based on a DARIS of 1532 participants. We calculated this DARIS based on a proportion of participants with acute mountain sickness of 32.6% in the control group; a RRR of 20% in the experimental intervention group; an alpha of 5%; a beta of 20%; and a diversity of 0%. After the second trial, the cumulative Z-curve (blue line) crossed the upper conventional alpha of 5%, but it did not cross the upper trial sequential alpha-spending monitoring boundaries, which were reached rather than crossed by the third trial. After only 39% (598/1532) of the DARIS had been reached, we were able to reject an intervention effect of 20% or larger, indicating that new randomized controlled trials are needed.

Comparison 3: non-steroidal anti-inflammatory drugs (NSAIDs) and other analgesics: carbasalate versus placebo

For this comparison, we analysed the information from one study ([Kayser 2008](#)) with 31 participants. [Kayser 2008](#) defined AMS in three different ways (Lake Louise AMS score ≥ 3 with headache; Lake Louise AMS score with headache and self-score + functional score ≥ 4 ; and Lake Louise AMS score with headache and self-score + functional score + clinical score ≥ 4). We chose the first definition for the following analyses.

Primary outcome 1: incidence of acute mountain sickness (AMS)

[Kayser 2008](#) provided information about the incidence of acute mountain sickness and found a total of 26 events (incidence of AMS = 83.8%). The RR for AMS, comparing carbasalate to placebo, was 0.91 (95% CI 0.67 to 1.25; 31 participants).

Secondary outcome 1: incidence of high altitude pulmonary oedema (HAPE)

We found no information about this outcome in the included study.

Secondary outcome 2: incidence of high altitude cerebral oedema (HACE)

We found no information about this outcome in the included study.

Secondary outcome 3: incidence of adverse events

We found no information about this outcome in the included study.

Secondary outcome 4: differences in HAI/AMS scores

We found no information about this outcome in the included study.

Group 7: Other comparisons**Comparison 1: other comparisons: acetazolamide versus dexamethasone**

For this comparison, we included information from three studies ([Ellsworth 1991](#); [Hussain 2001](#); [Zell 1988](#)), with a total of 46 participants. In [Ellsworth 1991](#), investigators administered 750 mg/day of acetazolamide. The study was carried out in the USA, reaching a maximum altitude of 4392 metres. [Zell 1988](#) and [Hussain 2001](#) used 500 mg/day of acetazolamide. [Zell 1988](#) was carried out in Nepal, reaching a maximum altitude of 4050 metres. We also included information from a cross-over study ([Subudhi 2011](#)), which compared acetazolamide 750 mg/day to 12 mg dexamethasone using a hypobaric chamber.

Primary outcome 1: incidence of acute mountain sickness (AMS)

We found no information about this outcome in the included studies.

Secondary outcome 1: incidence of high altitude pulmonary oedema (HAPE)

We found no information about this outcome in the included studies.

Secondary outcome 2: incidence of high altitude cerebral oedema (HACE)

We found no information about this outcome in the included studies.

Secondary outcome 3: incidence of adverse events

[Zell 1988](#) reported information about numbness in the fingers, finding six events (Incidence of numbness: 37.5%). The RR for numbness, comparing acetazolamide to dexamethasone, was 16.25 (95% CI 1.07 to 247.19; 16 participants).

Secondary outcome 4: differences in HAI/AMS scores

[Hussain 2001](#) provided information about differences in AMS scores at high altitude. The standardized mean difference for AMS scores, comparing acetazolamide to dexamethasone, was 0.292 (95% CI 0.06 to 0.52; 12 participants). We also found information about this outcome in [Subudhi 2011](#). The standardized mean difference for AMS scores, comparing acetazolamide to dexamethasone, was 0.00 (95% CI -0.23 to 0.23; 40 participants).

Comparison 2: other comparisons: acetazolamide plus dexamethasone versus acetazolamide

For this comparison, we analysed information from three studies ([Bernhard 1998](#); [Hussain 2001](#); [Zell 1988](#)), with a total of 40 participants. [Bernhard 1998](#) used 500 mg of acetazolamide/day plus 8 mg of dexamethasone/day. Forty per cent of the participants in this study had a history of previous mild or moderate AMS. This study was carried out in Italy, reaching a maximum altitude of 5334 metres. [Hussain 2001](#) and [Zell 1988](#) used 500 mg of acetazolamide/day plus 8 mg and 16 mg of dexamethasone/day respectively; there were no groups at risk of AMS, HAPE or HACE. [Zell 1988](#) was carried out in Nepal, reaching a maximum altitude of 4050 metres.

Primary outcome 1: incidence of acute mountain sickness (AMS)

[Bernhard 1998](#) found eight events of acute mountain sickness (incidence of AMS: 61.5%). The RR for acute mountain sickness, comparing acetazolamide plus dexamethasone to acetazolamide plus placebo, was 0.70 (95% CI 0.28 to 1.77; 13 participants).

Secondary outcome 1: incidence of high altitude pulmonary oedema (HAPE)

We found no information about this outcome in the included studies.

Secondary outcome 2: incidence of high altitude cerebral oedema (HACE)

We found no information about this outcome in the included studies.

Secondary outcome 3: incidence of adverse events

Zell 1988 reported information about numbness in the fingers, finding 11 events (incidence of numbness: 73.3%). The RR for numbness, comparing acetazolamide plus dexamethasone to acetazolamide, was 0.73 (95% CI 0.39 to 1.35; 15 participants).

Secondary outcome 4: differences in HAI/AMS scores

Hussain 2001 provided information about differences in AMS scores at high altitude. The mean difference for AMS scores, comparing acetazolamide to dexamethasone was -11.47 (95% CI -17.63 to -5.31; 12 participants).

Comparison 3: other comparisons: acetazolamide plus dexamethasone versus dexamethasone

For this comparison, we included information from two studies (Hussain 2001; Zell 1988), with a total of 29 participants. In Zell 1988 500 mg of acetazolamide/day plus 16 mg of dexamethasone/day were used. This study was carried out in Nepal, reaching a maximum altitude of 4050 metres.

Primary outcome 1: incidence of acute mountain sickness (AMS)

We found no information about this outcome in the included studies.

Secondary outcome 1: incidence of high altitude pulmonary oedema (HAPE)

We found no information about this outcome in the included studies.

Secondary outcome 2: incidence of high altitude cerebral oedema (HACE)

We found no information about this outcome in the included studies.

Secondary outcome 3: Incidence of adverse events

Zell 1988 reported information about numbness in fingers, finding five events (Incidence of numbness: 29.4%). The RR for numbness, comparing acetazolamide plus dexamethasone to dexamethasone was 12.22 (95% CI 0.78 to 191.46; 17 participants).

Secondary outcome 4: differences in HAI/AMS scores

Hussain 2001 provided information about differences in AMS scores at high altitude. The mean difference for AMS scores, comparing acetazolamide plus dexamethasone to dexamethasone was -9.17 (95% CI -15.62 to -2.72; 12 participants).

Comparison 4: other comparisons: acetazolamide versus ibuprofen

For this comparison, we analysed information from one study (HEAT 2010) with 254 participants. HEAT 2010 administered 225 mg of acetazolamide/day or 600 mg of ibuprofen/day.

Primary outcome 1: risk of acute mountain sickness

HEAT 2010 found 32 events of acute mountain sickness (incidence of AMS: 12.59%). The RR for AMS, comparing acetazolamide to ibuprofen, was 1.33 (95% CI 0.69 to 2.55; 163 participants).

Secondary outcome 1: risk of altitude pulmonary oedema.

We found no information about this outcome in the included study.

Secondary outcome 2: risk of high altitude cerebral oedema

We found no information about this outcome in the included study.

Secondary outcome 3: adverse events

HEAT 2010 did not identify any major adverse events.

Secondary outcome 4: differences in HAI/AMS scores

We found no information about this outcome in the included study.

Comparison 5: other comparisons: acetazolamide versus methazolamide

For this comparison, we analysed information from one study (Wright 1983) with 20 participants. Wright 1983 used 500 mg of acetazolamide/day and 100/150 mg of methazolamide/day. This study was carried out in high mountain areas of Nepal, reaching a maximum altitude of 4790 metres. Some participants in this study had a previous history of severe AMS.

Primary outcome 1: incidence of acute mountain sickness (AMS)

We found no information about this outcome in the included study.

Secondary outcome 1: incidence of high altitude pulmonary oedema (HAPE)

We found no information about this outcome in the included study.

Secondary outcome 2: incidence of high altitude cerebral oedema (HACE)

We found no information about this outcome in the included study.

Secondary outcome 3: incidence of adverse events

We found no information about this outcome in the included study.

Secondary outcome 4: differences in HAI/AMS scores

Wright 1983 reported information about AMS scores. The standardized mean difference between acetazolamide and methazolamide, was -3.00 (95% CI -21.07 to 15.07; 20 participants).

Comparison 6: other comparisons: budesonide plus formoterol versus placebo

For this comparison, we analysed the information from one study (Chen 2015) with 40 participants in the relevant arms. This study was carried out in China, reaching a maximum altitude of 3700 metres.

Primary outcome 1: incidence of acute mountain sickness (AMS)

Chen 2015 provide information about the incidence of acute mountain sickness and found 24 events (incidence of AMS = 60%). The RR for AMS, comparing budesonide plus formoterol to placebo, was 0.71 (95% CI 0.42 to 1.21; 40 participants).

Secondary outcome 1: incidence of high altitude pulmonary oedema (HAPE)

We found no information about this outcome in the included study.

Secondary outcome 2: incidence of high altitude cerebral oedema (HACE)

We found no information about this outcome in the included study.

Secondary outcome 3: incidence of adverse events

Chen 2015 assessed the incidence of side effects but found no events.

Secondary outcome 4: differences in HAI/AMS scores

We found no information about this outcome in the included study.

Comparison 7: other comparisons: budesonide versus dexamethasone

For this comparison, we analysed information from one study (Zheng 2014) with 92 participants. Zheng 2014 used 400 mg of budesonide/day and 4 mg of dexamethasone/day. This study was carried out in China, reaching a maximum altitude of 4050 metres.

Primary outcome 1: incidence of acute mountain sickness (AMS)

Zheng 2014 found 22 events of acute mountain sickness for this comparison (incidence of AMS = 23.9%). The RR for AMS, comparing budesonide to dexamethasone, was 0.83 (95% CI 0.40 to 1.73; 92 participants).

Secondary outcome 1: incidence of high altitude pulmonary oedema (HAPE)

We found no information about this outcome in the included study.

Secondary outcome 2: incidence of high altitude cerebral oedema (HACE)

We found no information about this outcome in the included study.

Secondary outcome 3: incidence of adverse events

Zheng 2014 found four events of persistent belching for this comparison (incidence of persistent belching = 4.34%). The RR for persistent belching, comparing budesonide to dexamethasone, was 0.11 (95% CI 0.01 to 2.01; 92 participants).

Secondary outcome 4: differences in HAI/AMS scores

We found no information about this outcome in the included study.

Comparison 8: other comparisons: budenoside versus budenoside plus formoterol

For this comparison, we analysed information from one study ([Chen 2015](#)) with 40 participants in the relevant arms. This study used 400 mg of budenoside/day and 9 mg of formoterol/day. It was carried out in China, reaching a maximum altitude of 3700 metres.

Primary outcome 1: incidence of acute mountain sickness (AMS)

[Chen 2015](#) found 15 events of acute mountain sickness for this comparison (total incidence of AMS = 37.5%). The RR for AMS, comparing budenoside to budenoside plus formoterol, was 0.50 (95% CI 0.21 to 1.20; 40 participants).

Secondary outcome 1: incidence of high altitude pulmonary oedema (HAPE)

We found no information about this outcome in the included study.

Secondary outcome 2: incidence of high altitude cerebral oedema (HACE)

We found no information about this outcome in the included study.

Secondary outcome 3: incidence of adverse events

[Chen 2015](#) did not find any side effects for this comparison.

Secondary outcome 4: differences in HAI/AMS scores

We found no information about this outcome in the included study.

Comparison 9: other comparisons: dexamethasone versus prednisolone

For this comparison, we analysed the information from one study ([Basu 2002a](#)) with 40 participants. However, this study did not provide information about any of outcomes selected for this review.

Comparison 10: other comparisons: tadalafil versus dexamethasone

For this comparison, we analysed information from one study ([Maggiorini 2006](#)) with 20 participants. [Maggiorini 2006](#) used 20 mg of tadalafil/day and 16 mg of dexamethasone/day. This study was carried out in Kenya, reaching a maximum altitude of 4559 metres. All participants had a history of HAPE.

Primary outcome 1: incidence of Acute mountain sickness (AMS)

[Maggiorini 2006](#) found 11 events of acute mountain sickness for this comparison (incidence of AMS = 55%). The RR for AMS, comparing tadalafil to dexamethasone, was 2.67 (95% CI 0.98 to 7.22; 20 participants).

Secondary outcome 1: incidence of high altitude pulmonary oedema (HAPE)

[Maggiorini 2006](#) found one event of altitude pulmonary oedema for this comparison (incidence of AMS = 5%). The RR for HAPE, comparing tadalafil to dexamethasone, was 3.0 (95% CI 0.14 to 65.9; 20 participants).

Secondary outcome 2: incidence of high altitude cerebral oedema (HACE)

We found no information about this outcome in the included study.

Secondary outcome 3: incidence of adverse events

We found no information about this outcome in the included study.

Secondary outcome 4: differences in HAI/AMS scores

We found no information about this outcome in the included study.

ADDITIONAL SUMMARY OF FINDINGS *[Explanation]*

Budesonide compared with placebo for preventing high altitude illness						
Patient or population: people at risk of high altitude illness Setting: High altitude; studies undertaken in India, South America and USA. Intervention: budesonide Comparison: placebo						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	placebo	Budesonide				
Incidence of acute mountain sickness (AMS) - Follow-up: From arrival to 24 hours later	606 per 1000	224 per 1000 (139 to 370)	RR 0.37 (0.23 to 0.61)	132 (2 studies)	⊕⊕○○ low ^{1,2}	
Incidence of high altitude pulmonary oedema (HAPE) - not reported	See comment	See comment	Not estimable	-	See comment	This outcome was not reported for selected trials.
Incidence of high altitude cerebral oedema (HACE) - not reported	See comment	See comment	Not estimable	-	See comment	This outcome was not reported for selected trials.
Adverse events: Side effects - Follow-up: From arrival to 24 hours later	See comment	See comment	Not estimable	40 (1 study)	⊕○○○ very low ^{3,4}	This trial reported no events

The **corresponding risk** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).
CI: Confidence interval; **RR:** Risk ratio;

GRADE Working Group grades of evidence

High quality: Further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: We are very uncertain about the estimate.

¹Risk of bias downgraded (-1) due to high risk of performance bias in one out of two studies included.

²Imprecision downgraded (-1) due to insufficient sample size to determine whether there are differences or not between these two groups.

³Risk of bias downgraded (-1) due to high risk of performance bias.

⁴Imprecision downgraded (-2) due to insufficient sample size to determine whether there are differences or not between these two groups.

Dexamethasone compared with placebo for preventing high altitude illness						
Patient or population: people at risk of high altitude illness						
Setting: High altitude; studies undertaken in India, South America and USA.						
Intervention: dexamethasone						
Comparison: placebo						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	placebo	Dexamethasone				
Incidence of acute mountain sickness (AMS) - Follow-up: From arrival to 24 hours later	449 per 1000	270 per 1000 (162 to 449)	RR 0.6 (0.36 to 1)	176 (4 studies)	⊕⊕○○ low ^{1,2}	
Incidence of high altitude pulmonary oedema (HAPE) - not reported	See comment	See comment	Not estimable	-	See comment	This outcome was not reported for selected trials.
Incidence of high altitude cerebral oedema (HACE) - not reported	See comment	See comment	Not estimable	-	See comment	This outcome was not reported for selected trials.
Adverse events: General - Follow-up: From arrival to 24 hours later	See comment	See comment	Not estimable	21 (1 study)	⊕○○○ very low ^{3,4}	This trial reported no events

The **corresponding risk** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).
CI: Confidence interval; **RR:** Risk ratio;

GRADE Working Group grades of evidence

High quality: Further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: We are very uncertain about the estimate.

¹Risk of bias downgraded (-1) due to unclear risk of selection, performance and detection bias in almost all studies included.

²Imprecision downgraded (-1) due to insufficient sample size to determine whether there are differences or not between these two groups.

³Risk of bias downgraded (-1) due to unclear risk of selection, performance and detection bias.

⁴Imprecision downgraded (-2) due to insufficient sample size to determine whether there are differences or not between these two groups.

DISCUSSION

Summary of main results

Evidence from 65 studies showed important findings for interventions included in this review (commonly-used pharmacological interventions). We report results for the three more important comparisons:

Acetazolamide versus placebo (28 parallel studies; 2345 participants)

Our systematic review included data from 28 parallel clinical studies ($n = 2345$ participants) and five cross-over studies ($n = 54$ participants) that assessed the effectiveness of acetazolamide compared with a placebo for the prevention of high altitude illness. The risk of AMS was reduced with acetazolamide (RR 0.47; 95% CI 0.39 to 0.56; $I^2 = 0\%$; 16 trials; 2301 participants; moderate quality of evidence). No events of HAPE were reported and only one event of HACE (RR 0.32; 95% CI 0.01 to 7.48; 6 parallel trials; 1126 participants; moderate quality of evidence). Few studies reported side effects for this comparison, and they showed an increase in the risk of paraesthesia with the intake of acetazolamide (5 studies, 789 participants; RR from 3.09 to 12.63 by acetazolamide dosage).

Budenoside versus placebo (2 parallel studies; 132 participants)

Data on budenoside showed a reduction in the incidence of AMS compared with placebo (2 studies, 132 participants; RR 0.37; 95% CI 0.23 to 0.61; $I^2 = 0\%$; low quality of evidence). The included studies did not report any events of HAPE or HACE, and they did not find side effects (low quality of the evidence).

Dexamethasone versus placebo (7 parallel studies; 205 participants)

For dexamethasone, data did not show benefits of dexamethasone at any dosage (four studies, 176 participants; RR 0.60; 95% CI 0.36 to 1.00; $I^2 = 39\%$; low quality of evidence). The studies did not report any events of HAPE or HACE, and we rated the evidence about adverse events as of very low quality.

We did not find any studies comparing methazolamide with a placebo. We also did not find evidence of benefits of theophylline, montelukast, selective inhibitors of phosphodiesterase-5 (such as tadalafil and sildenafil), nifedipine, flunarizine, aspirin or carbasalate in reducing the incidence of AMS. Finally, we found little information on other comparisons between different agents included in this review (i.e. ibuprofen versus placebo, acetazolamide versus dexamethasone). Combinations of these drugs did not deliver any benefits.

Overall completeness and applicability of evidence

We carried out a thorough search and identified an important number of studies addressing effectiveness and safety in the most

commonly-used pharmacological interventions for the prevention of HAI or AMS. We included 65 studies in our review, with more than 2000 participants. Those studies addressed around 15 comparisons with placebos, and 11 comparisons between different drugs. The data included participants of different age groups and both genders, as well as different high-altitude settings, different final altitudes reached, transportation, and prophylaxis times. Our systematic search for studies and our data extraction procedures should have minimized the likelihood of missing relevant studies. The funnel plot for acetazolamide versus placebo was highly symmetrical, suggesting that the chance of having missed relevant studies was minimal, with no evidence of publication bias. Despite all this, we found a lack of reports of the duration of prophylaxis, duration of ascent, criteria to diagnose AMS, HAPE or HACE, or statistical data (such as standard deviations) in several of the included studies. The sparsity of reports of adverse events was the most frequent limitation of the included studies, as well as the wide range of criteria and scales used to determine the onset of acute mountain sickness. The identification of only one study for several of the comparisons was a common factor limiting the scope and strength of this review.

The trial sequence analyses performed with on acetazolamide for the prevention of AMS suggest we have robust data for significant efficacy, which can be applied with some confidence in the field.

Quality of the evidence

We conducted GRADE assessments on outcomes of meta-analyses and single trials. We were unable to rate the evidence from either pooled or non-pooled estimates as high, due to either or both of the following reasons:

1. small sample sizes
2. the risk of bias from multiple sources, including the lack of adequate randomization methods, lack of blinding, high attrition, unclear reporting of outcomes, and bias in the presentation of data, among others.

We also downgraded the evidence because of uncertainty in clinically relevant outcomes, reflected in wide confidence intervals, i.e. imprecision. See [Summary of findings for the main comparison](#), [Summary of findings 2](#) and [Summary of findings 3](#) for detailed assessments and the rationale for ratings.

Potential biases in the review process

In all cases, we followed the methodology for systematic reviews outlined in the *Cochrane Handbook for Systematic Reviews of Interventions* (Higgins 2011). However, we had to make extensive modifications to the published protocol (Martí-Carvajal 2012), due to the need to update the methods under the current methodological guidelines for Cochrane Reviews. Readers should be aware of the potential biases related to these modifications (detailed in

Differences between protocol and review), as well as the decision to split the review into three parts, considering the numerous interventions assessed for HAI prevention.

In this review we undertook a comprehensive search to identify clinical trials addressing the issue of effectiveness and safety of commonly-used classes of drugs for preventing acute HAI. Twelve studies did not provide enough information to classify them as included or excluded, because they were published only as conference proceedings, or because we did not have access to the full texts when we were completing this review. We have also considered 12 additional studies as ongoing because they are published only as protocols and we may be able to decide whether or not to include them once they have been published. A potential source of bias in the review process is that most of the studies (more than 75%), did not report adverse events associated with the classes of drugs commonly-used for the prevention of AMS. This constitutes a lack of information about the safety profile of the drugs in question. Likewise, we did not expect to encounter any unit of analysis issues, as we did not expect to find cross-over studies. However, we identified 12 cross-over studies (20%). In order to avoid bias in the development of our review, we have analysed those studies separately.

Agreements and disagreements with other studies or reviews

There are several examples of published reviews evaluating different interventions to prevent high altitude illness. We found that our results are similar to other non-Cochrane reviews (Low 2012; Kayser 2012; Ritchie 2012; Seupaul 2012; Zafren 2014), regarding HAI/AMS prevention (CATMAT 2007; Luks 2010; Luks 2014). Most of these reviews recommend acetazolamide (at doses of 500 mg/day) as the first choice for the prevention of this condition. A systematic review developed by Dumont 2000 concludes that doses of 750 mg/day are more effective than 500 mg/day; however, our findings showed that effectiveness is similar for these two options, but there is no clear information on whether the incidence of adverse events is greater, due to the lack of information in the studies for this outcome.

In 2014, Tang 2014 published evidence in favour of the use of oral dexamethasone for the prevention of AMS. The authors of this review reported that dexamethasone could reduce the incidence of AMS, with an odds ratio of 6.03 (95% CI 2.23 to 21.00), compared with placebo. While they only identified eight studies comparing dexamethasone to placebo, we found six parallel trials and five cross-over studies. Our analysis did not produce definitive evidence about the effectiveness of dexamethasone, but we rated this evidence as being of low quality. In addition, our trial sequential analyses suggest that new randomized controlled trials are needed for this intervention. We note that current guidelines about AMS prevention include recommendations about the use of dexamethasone to prevent HAI/AMS, in 2 mg doses every six

hours or 4 mg every 12 hours (Luks 2010; Luks 2014). For the use of non-steroidal anti-inflammatory drugs (NSAIDs), our results are similar to those published by Pandit 2014, and support the use of ibuprofen as an alternative for acetazolamide, despite the fact that they provide analyses for all pooled NSAIDs (OR 0.43; 95% CI 0.27 to 0.69, $I^2 = 0\%$). We did not find any reviews about other options such as tadalafil, sildenafil, nifedipine, flunarizine or theophylline, and these are not recommended in current clinical practice guidelines for the prevention of this condition.

AUTHORS' CONCLUSIONS

Implications for practice

Our analysis suggests that acetazolamide, administered between three and five days prior to ascent, is an effective pharmacological agent to prevent acute altitude sickness in dosages of 250 to 750 mg/day. This information is based on evidence of moderate quality. Acetazolamide is associated with an increased risk of paraesthesia, which should be balanced against the suggested benefit. The clinical benefits and harms from other pharmacological interventions are unclear. There is little evidence relating to the prevention of HAPE and HACE, due to the low number of events reported.

Implications for research

There is a need for further high-quality research in this area. Future studies should be adequately powered to assess the effectiveness of these agents for the prevention of more serious forms of AMS, in combination as well as single agents. The design of future trials might be improved by the following suggestions:

1. Refining the clinical definition of AMS, HAPE and HACE.
2. Improving the reporting of statistical data related to important results, in order to avoid missing data, including information about elevation where HAI occurs.
3. Adding adverse events as an important endpoint in assessment of these preventive strategies.
4. Comparing pharmacological agents against interventions of established effectiveness (such as acetazolamide).

Finally, we suggest performing a network meta-analysis of all interventions (pharmacological and non-pharmacological) used for high altitude illness prevention, in order to determine which interventions are effective in avoiding the onset of new cases of this condition.

ACKNOWLEDGEMENTS

We would like to thank Mike Bennett (content editor), Vibeke E Horstmann (statistical editor), Janne Vendt (Information Scientist), Jeffrey H Gertsch, Martin Burtcher (peer reviewers), Matiram Pun (consumer referee) and Jane Cracknell (Managing Editor) for their help and editorial advice during the preparation of this systematic review.

REFERENCES

References to studies included in this review

Anonymous 1981 *{published data only}*

Anonymous. Acetazolamide in control of acute mountain sickness. *Lancet* 1981;**1**(8213):180–3. PUBMED: 6109857]

ASCENT 2012 *{published data only}*

Gertsch JH, Corbett B, Holck PS, Mulcahy A, Watts M, Stillwagon NT, et al. Altitude Sickness in Climbers and Efficacy of NSAIDs Trial (ASCENT): randomized, controlled trial of ibuprofen versus placebo for prevention of altitude illness. *Wilderness and Environmental Medicine* 2012;**23**(4):307–15.

Banderet 1977 *{published data only}*

Banderet LE. Self-rated moods of humans at 4300 m pretreated with placebo or acetazolamide plus staging. *Aviation, Space, and Environmental Medicine* 1977;**48**(1): 19–22. PUBMED: 831706]

Bartsch 1991 *{published data only}*

* Bartsch P, Maggiorini M, Ritter M, Noti C, Vock P, Oelz O. Prevention of high-altitude pulmonary edema by nifedipine. *New England Journal of Medicine* 1991;**325**(18): 1284–9. PUBMED: 1922223]

Oelz O, Maggiorini M, Ritter M, Noti C, Waber U, Vock P, et al. Prevention and treatment of high altitude pulmonary edema by a calcium channel blocker. *International Journal of Sports Medicine* 1992;**13**(1):S65–8. PUBMED: 1483797]

Basnyat 2003 *{published data only}*

Basnyat B, Gertsch JH, Johnson EW, Castro-Marin F, Inoue Y, Yeh C. Efficacy of low-dose acetazolamide (125 mg BID) for the prophylaxis of acute mountain sickness: a prospective, double-blind, randomized, placebo-controlled trial. *High Altitude Medicine & Biology* 2003;**4**(1):45–52. PUBMED: 12713711]

Basnyat 2008 *{published data only}*

Basnyat B, Hargrove J, Holck PS, Srivastav S, Alekh K, Ghimire LV, et al. Acetazolamide fails to decrease pulmonary artery pressure at high altitude in partially acclimatized humans. *High Altitude Medicine & Biology* 2008;**9**(3):209–16. PUBMED: 18800957]

Basu 2002a *{published data only}*

Basu M, Sawhney RC, Kumar S, Pal K, Prasad R, Selvamurthy W. Glucocorticoids as prophylaxis against acute mountain sickness. *Clinical Endocrinology* 2002;**57**(6):761–7. PUBMED: 12460326]

Basu 2002b *{published data only}*

Basu M, Sawhney RC, Kumar S, Pal K, Prasad R, Selvamurthy W. Hypothalamic-pituitary-adrenal axis following glucocorticoid prophylaxis against acute mountain sickness. *Hormone and Metabolic Research* 2002;**34**(6): 318–24. PUBMED: 12173072]

Bates 2011 *{published data only}*

Bates MG, Thompson AA, Baillie JK, Sutherland AI, Irving JB, Hirani N, et al. Sildenafil citrate for the prevention of high altitude hypoxic pulmonary hypertension: double blind, randomized, placebo-controlled trial. *High Altitude Medicine & Biology* 2011;**12**(3):207–14. PUBMED: 21962063]

Baumgartner 2003 *{published data only}*

Baumgartner RW, Keller S, Regard M, Bartsch P. Flunarizine in prevention of headache, ataxia, and memory deficits during decompression to 4559 m. *High Altitude Medicine & Biology* 2003;**4**(3):333–9. PUBMED: 14561238]

Bernhard 1994 *{published data only}*

Bernhard WN, Miller SL, Gittelsohn A. Dexamethasone for prophylaxis against acute mountain sickness during rapid ascent to 5334 m. *Journal of Wilderness Medicine* 1994;**5**(3):331–8.

Bernhard 1998 *{published data only}*

Bernhard WN, Schalick LM, Delaney PA, Bernhard TM, Barnas GM. Acetazolamide plus low-dose dexamethasone is better than acetazolamide alone to ameliorate symptoms of acute mountain sickness. *Aviation, Space, and Environmental Medicine* 1998;**69**(9):883–6. PUBMED: 9737760]

Bradwell 1986 *{published data only}*

Bradwell AR, Dykes PW, Coote JH, Forster PJ, Milles JJ, Chesner I, et al. Effect of acetazolamide on exercise performance and muscle mass at high altitude. *Lancet* 1986;**1**(8488):1001–5. PUBMED: 2871285]

Burki 1992 *{published data only}*

Burki NK, Khan SA, Hameed MA. The effects of acetazolamide on the ventilatory response to high altitude hypoxia. *Chest* 1992;**101**(3):736–41. PUBMED: 1541140]

Burtcher 1998 *{published data only}*

Burtcher M, Likar R, Nachbauer W, Philadelphia M. Aspirin for prophylaxis against headache at high altitudes: randomised, double blind, placebo controlled trial. *BMJ* 1998;**316**(7137):1057–8.

- Burtscher 2001** *{published data only}*
 Burtscher M, Likar R, Nachbauer W, Philadelphia M, Pühringer R, Lämmle T. Effects of aspirin during exercise on the incidence of high-altitude headache: a randomized, double-blind, placebo-controlled trial. *Headache* 2001;**41**(6):542–5.
- Burtscher 2014** *{published data only}*
 Burtscher M, Gatterer H, Faulhaber M, Burtscher J. Acetazolamide pre-treatment before ascending to high altitudes: When to start?. *International Journal of Clinical and Experimental Medicine* 2014;**15**(11):4378–83. PUBMED: 25550957]
- Carlsten 2004** *{published data only}*
 Carlsten C, Swenson ER, Ruoss S. A dose-response study of acetazolamide for acute mountain sickness prophylaxis in vacationing tourists at 12,000 feet (3630 m). *High Altitude Medicine & Biology* 2004;**5**(1):33–9. PUBMED: 15072715]
- Chen 2015** *{published data only}*
 * Chen GZ, Zheng CR, Qin J, Yu J, Wang H, Zhang JH, et al. Inhaled budesonide prevents acute mountain sickness in young Chinese men. *Journal of Emergency Medicine* 2015;**48**(2):197–206. PUBMED: 25294611]
 Xiangjun L, Lan H. Inhaled budesonide for the prevention of acute mountain sickness in unacclimatization young men: A double-blind randomized controlled trial. *Journal of the American College of Cardiology* 2014;**16** Suppl 1(64): C69.
- Chow 2005** *{published data only}*
 Chow T, Browne V, Heilesen HL, Wallace D, Anholm J, Green SM. Ginkgo biloba and acetazolamide prophylaxis for acute mountain sickness: a randomized, placebo-controlled trial. *Archives of Internal Medicine* 2005;**165**(3): 296–301. PUBMED: 15710792]
- Ellsworth 1991** *{published data only}*
 Ellsworth AJ, Meyer EF, Larson EB. Acetazolamide or dexamethasone use versus placebo to prevent acute mountain sickness on Mount Rainier. *Western Journal of Medicine* 1991;**154**(3):289–93. PUBMED: 2028586]
- Faull 2015** *{published data only}*
 Faull OK, Robertson J, Thomas O, Bradwell AR, Antoniadis CA, Pattinson KTS, et al. The effect of acetazolamide on saccadic latency at 3459 meters. *Wilderness and Environmental Medicine* 2015;**26**(1):72–7.
- Fischer 2000a** *{published data only}*
 Fischer R, Hautmann H, Toepfer M, Tschoep M, Pongratz H, Huber RM. Oral slow release theophylline (σT) is effective in prevention of acute mountain sickness [abstract]. *European Respiratory Journal. Supplement*. 1998;**12**(Suppl 28):407S. CN–00382822]
 * Fischer R, Lang SM, Steiner U, Toepfer M, Hautmann H, Pongratz H, et al. Theophylline improves acute mountain sickness. *European Respiratory Journal* 2000;**15**(1):123–7. PUBMED: 10678632]
- Fischer 2000b** *{published data only}*
 Fischer R, Vollmar C, Thiery M, Born C, Leitl M, Pfluger T, et al. No evidence of cerebral oedema in severe acute mountain sickness. *Cephalalgia* 2004;**24**(1):66–71. PUBMED: 14687016]
- Fischer 2004** *{published data only}*
 Fischer R, Vollmar C, Thiery M, Born C, Leitl M, Pfluger T, et al. No evidence of cerebral oedema in severe acute mountain sickness. *Cephalalgia* 2004;**24**(1):66–71. PUBMED: 14687016]
- Fulco 2006** *{published data only}*
 Fulco CS, Muza SR, Ditzler D, Lammi E, Lewis SF, Cymerman A. Effect of acetazolamide on leg endurance exercise at sea level and simulated altitude. *Clinical Science* 2006;**110**(6):683–92. PUBMED: 16499476]
- Greene 1981** *{published data only}*
 Greene MK, Kerr AM, McIntosh IB, Prescott RJ. Acetazolamide in prevention of acute mountain sickness: a double-blind controlled cross-over study. *BMJ* 1981;**283**(6295):811–3. PUBMED: 6794709]
- Hackett 1976** *{published data only}*
 Hackett PH, Rennie D, Levine HD. The incidence, importance, and prophylaxis of acute mountain sickness. *Lancet* 1976;**2**(7996):1149–55. PUBMED: 62991]
- Hackett 1988** *{published data only}*
 Hackett PH, Roach RC, Wood RA, Foutch RG, Meehan RT, Rennie D, et al. Dexamethasone for prevention and treatment of acute mountain sickness. *Aviation, Space, and Environmental Medicine* 1988;**59**(10):950–4. PUBMED: 3190622]
- HEAT 2010** *{published data only}*
 Gertsch JH, Lipman GS, Holck PS, Merritt A, Mulcahy A, Fisher RS, et al. Prospective, double-blind, randomized, placebo-controlled comparison of acetazolamide versus ibuprofen for prophylaxis against high altitude headache: the Headache Evaluation at Altitude Trial (HEAT). *Wilderness & Environmental Medicine* 2010;**21**(3):236–43. PUBMED: 20832701]
- Hillenbrand 2006** *{published data only}*
 Hillenbrand P, Pahari AK, Soon Y, Subedi D, Bajracharya R, Gurung P, et al. Prevention of acute mountain sickness by acetazolamide in Nepali porters: a double-blind controlled trial. *Wilderness & Environmental Medicine* 2006;**17**(2): 87–93. PUBMED: 16805144]
- Hochapfel 1986** *{published data only}*
 Hochapfel G, Schlemmer W. Prophylaxis of acute mountain sickness. Experiences with acetazolamide in Nepal. *Munchener Medizinische Wochenschrift* 1986;**128**(8):137–8.
- Hohenhaus 1994** *{published data only}*
 Hohenhaus E, Goerre S, Niroomand F, Oelz O, Bartsch P. Nifedipine does not prevent acute mountain sickness [abstract]. *European Respiratory Journal. Supplement*. 1993; Vol. 7 Suppl 17:187S. CN–00393692]
 * Hohenhaus E, Niroomand F, Goerre S, Vock P, Oelz O, Bartsch P. Nifedipine does not prevent acute mountain sickness. *American Journal of Respiratory and Critical Care Medicine* 1994;**150**(3):857–60. PUBMED: 8087361]
- Hussain 2001** *{published data only}*
 Hussain MM, Aslam M, Khan Z. Acute mountain sickness score and hypoxemia. *Journal of the Pakistan Medical Association* 2001;**5**(5):173–9.

Jain 1986 {published data only}

Jain SC, Singh MV, Sharma VM, Rawal SB, Tyagi AK. Amelioration of acute mountain sickness: comparative study of acetazolamide and spironolactone. *International Journal of Biometeorology* 1986;**30**(4):293–300. PUBMED: 3804482]

Johnson 1984 {published data only}

Johnson TS, Rock PB, Fulco CS, Trad LA, Spark RF, Maher JT. Prevention of acute mountain sickness by dexamethasone. *New England Journal of Medicine* 1984;**310**(11):683–6. PUBMED: 6700643]

Kayser 2008 {published data only}

Kayser B, Hulsebosch R, Bosch F. Low-dose acetylsalicylic acid analog and acetazolamide for prevention of acute mountain sickness. *High Altitude Medicine & Biology* 2008;**9**(1):15–23. PUBMED: 18331216]

Ke 2013 {published data only}

Ke T, Wang J, Swenson ER, Zhang X, Hu Y, Chen Y, et al. Effect of acetazolamide and ginkgo biloba on the human pulmonary vascular response to an acute altitude ascent. *High Altitude Medicine & Biology* 2013;**14**(2):162–7. PUBMED: 23795737]

Küpper 2008 {published data only}

Küpper TE, Strohl KP, Hofer M, Gieseler U, Netzer CM, Netzer NC. Low-dose theophylline reduces symptoms of acute mountain sickness. *Journal of Travel Medicine* 2008;**15**(5):307–14. PUBMED: 19006503]

Larson 1982a {published data only}

Larson EB, Roach RC, Schoene RB, Hornbein TF. Acute mountain sickness and acetazolamide. Clinical efficacy and effect on ventilation. *JAMA* 1982;**248**(3):328–32. PUBMED: 7045433]

Larson 1982b {published data only}

Lipman 2012 {published data only}

Lipman GS, Kanaan NC, Holck P, Gertsch JH, Constance BB, Corbett B, et al. Ibuprofen prevents altitude illness: A prospective, double-blind, randomized controlled trial. *Academic Emergency Medicine* 2011;**18**(5 Suppl 1):S15.
* Lipman GS, Kanaan NC, Holck PS, Constance BB, Gertsch JH, PAINS Group. Ibuprofen prevents altitude illness: a randomized controlled trial for prevention of altitude illness with nonsteroidal anti-inflammatories. *Annals of Emergency Medicine* 2012;**58**(6):484–90. PUBMED: 22440488]

Luks 2007 {published data only}

Luks AM, Henderson WR, Swenson ER. Leukotriene receptor blockade does not prevent acute mountain sickness induced by normobaric hypoxia. *High Altitude Medicine & Biology* 2007;**8**(2):131–8. PUBMED: 17584007]

Maggiorini 2006 {published data only}

Bernheim AM, Kiencke S, Fischler M, Dorschner L, Debrunner J, Mairbaurl H, et al. Acute changes in pulmonary artery pressures due to exercise and exposure

to high altitude do not cause left ventricular diastolic dysfunction. *Chest* 2007;**132**:380–7.

Fischler M, Maggiorini M, Dorschner L, Debrunner J, Bernheim A, Kiencke S, et al. Dexamethasone but not tadalafil improves exercise capacity in adults prone to high-altitude pulmonary edema. *American Journal of Respiratory and Critical Care Medicine* 2009;**180**(4):346–52. PUBMED: 19520908]

* Maggiorini M, Brunner-La Rocca HP, Peth S, Fischler M, Bohm T, Bernheim A, et al. Both tadalafil and dexamethasone may reduce the incidence of high-altitude pulmonary edema: a randomized trial. *Annals of Internal Medicine* 2006;**145**(7):497–506. PUBMED: 17015867]

Mairbaurl H, Brunner-La Rocca HP, Peth S, Dehnert C, Fischler M, Bohm T, et al. Phosphodiesterase-5 inhibitors and glucocorticoids prevent a high-altitude pulmonary oedema. *Deutsche Zeitschrift für Sportmedizin* 2005; Vol. 56, issue 7/8:218. CN-00727391]

Mirrahlmov 1993 {published data only}

Mirrahlmov M, Brimkulov N, Cielicki J, Tobiasz M, Kudaiberdiev Z, Moldotashev I, et al. Asthma treatment at high altitude effects of acetazolamide on overnight oximetry and prevention of acute mountain sickness [Abstract]. *European Respiratory Journal* 1992; Vol. 5, issue Suppl 15: 366s. CN-00494881]

* Mirrahlmov M, Brimkulov N, Cieslicki J, Tobiasz M, Kudaiberdiev Z, Moldotashev I, et al. Effects of acetazolamide on overnight oxygenation and acute mountain sickness in patients with asthma. *European Respiratory Journal* 1993;**6**(4):536–40. PUBMED: 8491304]

Montgomery 1989 {published data only}

Montgomery AB, Luce JM, Michael P, Mills J. Effects of dexamethasone on the incidence of acute mountain sickness at two intermediate altitudes. *JAMA* 1989;**261**(5):734–6. PUBMED: 2911170]

Moraga 2007 {published data only}

Moraga FA, Flores A, Serra J, Esnaola C, Barriento C. Ginkgo biloba decreases acute mountain sickness in people ascending to high altitude at Ollague (3696 m) in northern Chile. *Wilderness & Environmental Medicine* 2007;**18**(4): 251–7. PUBMED: 18076292]

Muza 2004 {published data only}

Muza SR, Kaminsky D, Fulco CS, Banderet LE, Cymerman A. Cysteinyl leukotriene blockade does not prevent acute mountain sickness. *Aviation, Space, and Environmental Medicine* 2004;**75**(5):413–9. PUBMED: 15152893]

PACE 2006 {published data only}

Basnyat B, Gertsch JH, Holck PS, Johnson EW, Luks AM, Donham BP, et al. Acetazolamide 125 mg BD is not significantly different from 375 mg BD in the prevention of acute mountain sickness: the prophylactic acetazolamide dosage comparison for efficacy (PACE) trial. *High Altitude Medicine & Biology* 2006;**7**(1):17–27. PUBMED: 16544963]

Parati 2013 {published data only}

Caravita S, Faini A, Lombardi C, Valentini M, Gregorini F, Rossi J, et al. Sex and acetazolamide effects on chemoreflex and periodic breathing during sleep at altitude. *Chest* 2015; **147**(1):120–31.

* Parati G, Revera M, Giuliano A, Faini A, Bilo G, Gregorini F, et al. Effects of acetazolamide on central blood pressure, peripheral blood pressure, and arterial distensibility at acute high altitude exposure. *European Heart Journal* 2013; **10**(10):759–66.

Salvi P, Revera M, Faini A, Giuliano A, Gregorini F, Agostoni, et al. Changes in subendocardial viability ratio with acute high-altitude exposure and protective role of acetazolamide. *Hypertension* 2013; **61**(4):793–9.

PHAIT 2004 {published data only}

Gertsch JH, Basnyat B, Johnson EW, Onopa J, Holck PS. Randomised, double blind, placebo controlled comparison of ginkgo biloba and acetazolamide for prevention of acute mountain sickness among Himalayan trekkers: the prevention of high altitude illness trial (PHAIT). *BMJ* 2004; **328**(7443):797. PUBMED: 15070635]

Rock 1987 {published data only}

Rock PB, Johnson TS, Cymerman A, Burse RL, Falk LJ, Fulco CS. Effect of dexamethasone on symptoms of acute mountain sickness at Pikes Peak, Colorado (4,300 m). *Aviation, Space, and Environmental Medicine* 1987; **58**(7):668–72. PUBMED: 3619842]

Rock 1989a {published data only}

Rock PB, Johnson TS, Larsen RF, Fulco CS, Trad LA, Cymerman A. Dexamethasone as prophylaxis for acute mountain sickness. Effect of dose level. *Chest* 1989; **95**(3):568–73. PUBMED: 2920585]

Rock 1989b {published data only}

Rock 1989c {published data only}

Sartori 2002 {published data only}

Sartori C, Allemann Y, Duplain H, Lepori M, Egli M, Lipp E, et al. Salmeterol for the prevention of high-altitude pulmonary edema. *New England Journal of Medicine* 2002; **346**(21):1631–6. PUBMED: 12023995]

SPACE 2011 {published data only}

Basnyat B, Holck PS, Pun M, Halverson S, Szawarski P, Gertsch J, et al. Spironolactone does not prevent acute mountain sickness: a prospective, double-blind, randomized, placebo-controlled trial by SPACE Trial Group (spironolactone and acetazolamide trial in the prevention of acute mountain sickness group). *Wilderness & Environmental Medicine* 2011; **22**(1):15–22. PUBMED: 21377114]

Subudhi 2011 {published data only}

Julian CG, Subudhi AW, Wilson MJ, Dimmen AC, Pecha T, Roach RC. Acute mountain sickness, inflammation, and permeability: New insights from a blood biomarker study. *Journal of Applied Physiology* 2011; **111**(2):392–9.

* Subudhi AW, Dimmen AC, Julian CG, Wilson MJ, Panerai RB, Roach RC. Effects of acetazolamide and dexamethasone on cerebral hemodynamics in hypoxia.

Journal of Applied Physiology 2011; **110**(5):1219–25. PUBMED: 21393464]

Van Patot 2008 {published data only}

* Van Patot MC, Leadbetter G 3rd, Keyes LE, Maakestad KM, Olson S, Hackett PH. Prophylactic low-dose acetazolamide reduces the incidence and severity of acute mountain sickness. *High Altitude Medicine & Biology* 2008; **9**(4):289–93. PUBMED: 19115912]

Wang 2013 {published data only}

Wang J, Ke T, Zhang X, Chen Y, Liu M, Chen J, et al. Effects of acetazolamide on cognitive performance during high-altitude exposure. *Neurotoxicology and Teratology* 2013; **35**:28–33. PUBMED: 23280141]

Wright 1983 {published data only}

Wright AD, Bradwell AR, Fletcher RF. Methazolamide and acetazolamide in acute mountain sickness. *Aviation, Space, and Environmental Medicine* 1983; **54**(7):619–21. PUBMED: 6349608]

Wright 2004 {published data only}

Wright AD, Beazley MF, Bradwell AR, Chesner IM, Clayton RN, Forster PJ, et al. Medroxyprogesterone at high altitude. The effects on blood gases, cerebral regional oxygenation, and acute mountain sickness. *Wilderness & Environmental Medicine* 2004; **15**(1):25–31. PUBMED: 15040503]

Zell 1988 {published data only}

Zell SC, Goodman PH. Acetazolamide and dexamethasone in the prevention of acute mountain sickness. *Western Journal of Medicine* 1988; **148**(5):541–5. PUBMED: 3051673]

Zheng 2014 {published data only}

Zheng CR, Chen GZ, Yu J, Qin J, Song P, Bian SZ, et al. Inhaled budesonide and oral dexamethasone prevent acute mountain sickness. *American Journal of Medicine* 2014; **127**(10):1001–9.e2. PUBMED: 24784698]

References to studies excluded from this review

ACME-1 2006 {published data only}

Modesti PA, Vanni S, Morabito M, Modesti A, Marchetta M, Gamberi T, et al. Role of endothelin-1 in exposure to high altitude: Acute Mountain Sickness and Endothelin-1 (ACME-1) study. *Circulation* 2006; **114**(13):1410–6. PUBMED: 16982943]

Agostoni 2013 {published data only}

Agostoni P, Swenson ER, Fumagalli R, Salvionia E, Cattadoria G, Farina S, et al. Acute high-altitude exposure reduces lung diffusion: Data from the HIGHCARE Alps project. *Respiratory Physiology & Neurobiology* 2013; **188**(2):223–8.

Bartsch 1993 {published data only}

* Bärtsch P, Merki B, Hofstetter D, Maggiorini M, Kayser B, Oelz O. Treatment of acute mountain-sickness by simulated descent - a randomized controlled trial. *BMJ* 1993; **306**(6885):1098–101.

Bärtsch 1994 {published data only}

* Bärtsch P, Maggi S, Kleger GR, Ballmer PE, Baumgartner RW. Sumatriptan for high-altitude headache. *Lancet* 1994; **344**(8934):1445. PUBMED: 7968111]

Bilo 2015 {published data only}

Bilo G, Lombardi C, Revera M, Diazzi E, Giuliano A, Faini A, et al. Acetazolamide counteracts ambulatory blood pressure increase under acute exposure to high altitude. *High Blood Pressure and Cardiovascular Prevention* 2011; **18**(3):118.

* Bilo G, Villafuerte FC, Faini A, Anza-Ramírez C, Revera M, Giuliano A, et al. Ambulatory blood pressure in untreated and treated hypertensive patients at high altitude: The High Altitude Cardiovascular Research-Andes study. *Hypertension* 2015; **65**(6):1266–72.

Bilo, G, Villafuerte F, Anza C, Revera, Giuliano A, Faini A, et al. Combined antihypertensive treatment and blood pressure responses to acute high altitude exposure in patients with hypertension. HIGHCARE-ANDES Lowlanders Study. *European Heart Journal* 2011; **34**(Suppl 1):269–70.

Bloch 2009 {published data only}

Bloch KE, Turk AJ, Maggiorini M, Hess T, Merz T, Bosch MM, et al. Effect of ascent protocol on acute mountain sickness and success at Muztagh Ata, 7546 m. *High Altitude Medicine & Biology* 2009; **10**(1):25–32. PUBMED: 19326598]

Broome 1994 {published data only}

Broome JR, Stoneham MD, Beeley JM, Milledge JS, Hughes AS. High altitude headache: treatment with ibuprofen. *Aviation, Space, and Environmental Medicine* 1994; **65**(1):19–20. PUBMED: 8117220]

Cain 1966 {published data only}

Cain SM, Dunn JE 2nd. Low doses of acetazolamide to aid accommodation of men to altitude. *Journal of Applied Physiology* 1966; **21**(4):1195–200. PUBMED: 5916650]

Debevec 2015 {published data only}

Debevec T, Pialoux V, Saugy J, Schmitt L, Cejuela R, Mury P, et al. Prooxidant/Antioxidant balance in hypoxia: A cross-over study on normobaric vs. hypobaric “live high-train low”. *PLoS ONE* 2015; **10**(9):e0137957.

Dumont 1999 {published data only}

Dumont L, Mardirosoff C, Soto-Debeuf G, Tassonyi E. Magnesium and acute mountain sickness. *Aviation, Space, and Environmental Medicine* 1999; **70**(6):625. PUBMED: 10373058]

Forster 1982 {published data only}

Forster P. Methazolamide in acute mountain sickness. *Lancet* 1982; **1**(8283):1254. PUBMED: 6123014]

Forwand 1968 {published data only}

Forwand SA, Landowne M, Follansbee JN, Hansen JE. Effect of acetazolamide on acute mountain sickness. *New England Journal of Medicine* 1968; **279**(16):839–45. PUBMED: 4877992]

Fulco 2011 {published data only}

Fulco CS, Muza SR, Beidleman BA, Demes R, Staab JE, Jones JE, et al. Effect of repeated normobaric hypoxia

exposures during sleep on acute mountain sickness, exercise performance, and sleep during exposure to terrestrial altitude. *American Journal of Physiology. Regulatory, Integrative and Comparative Physiology* 2011; **300**(2):R428–36. PUBMED: 21123763]

Gertsch 2002 {published data only}

Gertsch JH, Seto TB, Mor J, Onopa J. Ginkgo biloba for the prevention of severe acute mountain sickness (AMS) starting one day before rapid ascent. *High Altitude Medicine & Biology* 2002; **3**(1):29–37. PUBMED: 12006162]

Gray 1971 {published data only}

Gray GW, Bryan AC, Frayser R, Houston CS, Rennie ID. Control of acute mountain sickness. *Aerospace Medicine* 1971; **42**(1):81–4. PUBMED: 4925130]

Harris 2003 {published data only}

Harris NS, Wenzel RP, Thomas SH. High altitude headache: efficacy of acetaminophen vs. ibuprofen in a randomized, controlled trial. *Journal of Emergency Medicine* 2003; **24**(4):383–7. PUBMED: 12745039]

Johnson 1988 {published data only}

Johnson TS, Rock PB, Young JB, Fulco CS, Trad LA. Hemodynamic and sympathoadrenal responses to altitude in humans: effect of dexamethasone. *Aviation, Space, and Environmental Medicine* 1988; **59**(3):208–12. PUBMED: 3355474]

Jonk 2007 {published data only}

Jonk AM, Van den Berg IP, Olfert IM, Wray DW, Arai T, Hopkins SR, et al. Effect of acetazolamide on pulmonary and muscle gas exchange during normoxic and hypoxic exercise. *Journal of Physiology* 2007; **579**(Pt 3):909–21. PUBMED: 17218362]

Kotwal 2015 {published data only}

Kotwal J, Kotwal A, Bhalla S, Singh PK, Nair V. Effectiveness of homocysteine lowering vitamins in prevention of thrombotic tendency at high altitude area: A randomized field trial. *Thrombosis Research* 2015; **136**(4):758–62.

Lalande 2009 {published data only}

Lalande S, Snyder EM, Olson TP, Hulsebus ML, Orban M, Somers VK, et al. The effects of sildenafil and acetazolamide on breathing efficiency and ventilatory control during hypoxic exercise. *European Journal of Applied Physiology* 2009; **106**(4):509–15. PUBMED: 19337745]

Lawley 2012 {published data only}

Lawley JS, Oliver SJ, Mullins P, Morris D, Junglee NA, Jelleyman C, et al. Optic nerve sheath diameter is not related to high altitude headache: A randomized controlled trial. *High Altitude Medicine and Biology* 2012; **3**:193–9.

Levine 1989 {published data only}

Levine BD, Yoshimura K, Kobayashi T, Fukushima M, Shibamoto T, Ueda G. Dexamethasone in the treatment of acute mountain sickness. *New England Journal of Medicine* 1989; **321**(25):1707–13. PUBMED: 2687688]

Liu 2013 {published data only}

Liu C, Croft QP, Kalidhar S, Brooks JT, Herigstad M, Smith TG, et al. Dexamethasone mimics aspects of physiological

- acclimatization to 8 hours of hypoxia but suppresses plasma erythropoietin. *Journal of Applied Physiology* 2013;**114**(7): 948–56. PUBMED: 23393065]
- Mairer 2012** *{published data only}*
Mairer K, Gobel M, Defrancesco M, Wille M, Messner H, Loizides A, et al. MRI evidence: acute mountain sickness is not associated with cerebral edema formation during simulated high altitude. *PLoS One* 2012;**7**(11):e50334. PUBMED: 23226263]
- McIntosh 1986** *{published data only}*
McIntosh IB, Prescott RJ. Acetazolamide in prevention of acute mountain sickness. *Journal of International Medical Research* 1986;**14**(5):285–7. PUBMED: 3533677]
- Purkayastha 1995** *{published data only}*
Purkayastha SS, Ray US, Arora BS, Chhabra PC, Thakur L, Bandopadhyay P, et al. Acclimatization at high altitude in gradual and acute induction. *Journal of Applied Physiology* 1995;**79**(2):487–92. PUBMED: 7592207]
- Reinhart 1994** *{published data only}*
Reinhart WH, Goerre S, Bartsch P. Acetazolamide reduces the erythropoietin response to hypoxia at high altitude in humans. *Journal of Wilderness Medicine* 1994;**5**(3):312–7.
- Sandoval 2000** *{published data only}*
Sandoval M, Silva J, Parra M, Cifuentes F, Maluenda R, Escobar J. Effects of consumption of antioxidants in altitude exposure [Efectos del consumo de antioxidantes en exposición a altitud]. *Boletín Científico de la Asociación Chilena de Seguridad* 2000;**2**(3):67–73.
- Scalzo 2015** *{published data only}*
Scalzo RL, Binns SE, Klochak AL, Giordano GR, Paris HLR, Sevits KJ, et al. Methazolamide plus aminophylline abrogates hypoxia-mediated endurance exercise impairment. *High Altitude Medicine & Biology* 2015;**16**:331–42.
- Serra 2001** *{published data only}*
Serra J. Chachacoma: An alternative treatment for the prevention of acute mountain sickness? [Chachacoma ¿un tratamiento alternativo para la prevención del mal agudo de montaña?]. *Revista de Ciencias de la Salud* 2001;**5**(1):36–42.
- Siebenmann 2011** *{published data only}*
Siebenmann C, Bloch KE, Lundby C, Nussbamer-Ochsner Y, Schoeb M, Maggiorini M. Dexamethasone improves maximal exercise capacity of individuals susceptible to high altitude pulmonary edema at 4559 m. *High Altitude Medicine & Biology* 2011;**12**(2):169–77. PUBMED: 21718165]
- Singh 1969** *{published data only}*
Singh I, Khanna PK, Srivastava MC, Lal M, Roy SB, Subramanyam CS. Acute mountain sickness. *New England Journal of Medicine* 1969;**280**(4):175–84.
- Solís 1984** *{published data only}*
Solís JV. Acetazolamide: Effects on the “acute mountain sickness”, aggravated by physical activity. *Boletín de la Fundación Jiménez Díaz* 1984;**11**(3):117–20.
- Suh 2015** *{published data only}*
Suh KS, Kim T, Yi NJ, Hong G. Preparation for high altitude expedition and changes in cardiopulmonary and biochemical laboratory parameters with ascent to high altitude in transplant patients and live donors. *Clinical Transplantation* 2015;**29**(11):1013–20.
- Teppema 2007** *{published data only}*
Teppema LJ, Balanos GM, Steinback CD, Brown AD, Foster GE, Duff HJ, et al. Effects of acetazolamide on ventilatory, cerebrovascular, and pulmonary vascular responses to hypoxia. *American Journal of Respiratory and Critical Care Medicine* 2007;**175**(3):277–81. PUBMED: 17095745]
- Vuyk 2006** *{published data only}*
Vuyk J, Van den Bos J, Terhell K, De Bos R, Vletter A, Valk P, et al. Acetazolamide improves cerebral oxygenation during exercise at high altitude. *High Altitude Medicine & Biology* 2006;**7**(4):290–301. PUBMED: 17173514]
- White 1984** *{published data only}*
White AJ. Cognitive impairment of acute mountain sickness and acetazolamide. *Aviation, Space, and Environmental Medicine* 1984;**55**(7):598–603. PUBMED: 6466255]
- Wright 1988** *{published data only}*
Wright AD, Bradwell AR, Jensen J, Lassen N. Cerebral blood flow in acute mountain sickness and treatment with acetazolamide. *Clinical Science* 1988;**74** Suppl 18:1P.

References to studies awaiting assessment

- Dugas 1995** *{published data only}*
Dugas L, Dubray C, Herry JP, Olsen NV, Court-Payen M, Hansen JM, et al. Cardiovascular effects of a calcium channel blocker in hypoxia caused by altitude [Effets cardiovasculaires d'un bloqueur calcique en hypoxie d'altitude]. *Presse Medicale* 1995;**24**(16):763–8. PUBMED: 7784415]
- Ellsworth 1987** *{published data only}*
Ellsworth AJ, Larson EB, Strickland D. A randomized trial of dexamethasone and acetazolamide for acute mountain sickness prophylaxis. *American Journal of Medicine* 1987;**83**(6):1024–30. PUBMED: 3332564]
- Furian 2016** *{published data only}*
Furian M, Lichtblau M, Aeschbacher SS, Bisang M, Hartmann SE, Poulin M, et al. Altitude related adverse health effects in lowlanders with COPD travelling to 3, 200 m. randomized trial of preventive dexamethasone treatment. *Respiration* 2016;**91**(5):415.
- Hefli 2014** *{published data only}*
Hefli JP, Stutz M, Geiser T, Hefli U, Merz T, Huber A. Effect of antioxidant supplements on AMS and endothelial function during a high altitude expedition: A prospective randomized double-blind trial. *High Altitude Medicine & Biology* 2014;**15**(2):A262.
- Kasic 1991** *{published data only}*
Kasic JF, Yaron M, Nicholas RA, Lickteig JA, Roach R. Treatment of acute mountain-sickness - hyperbaric versus

oxygen-therapy. *Annals of Emergency Medicine* 1991;**20**(10): 1109–12.

Lee 2011 *{published data only}*

Lee JH, Choi PC. Comparison of methazolamide and acetazolamide for prevention of acute mountain sickness in adolescents. *Journal of the Korean Society of Emergency Medicine* 2011;**22**(5):523–30.

Pun 2014 *{published data only}*

Pun M, Neupane M, Lohani A, Thapa GB, Yadav S, Holck PS, et al. Role of low-dose acetazolamide (125 mg BID) in prevention of acute mountain sickness in pilgrims ascending rapidly: A prospective double-blind placebocontrolled randomized trial. *High Altitude Medicine & Biology* 2014;**15**(2):A233.

Roncin 1996 *{published data only}*

Roncin JP, Schwartz F, Darbigny P. EGb 761 in control of acute mountain sickness and vascular reactivity to cold exposure. *Aviation, Space, and Environmental Medicine* 1996;**67**(5):445–52.

Swenson 1997 *{published data only}*

Swenson ER, Macdonald A, Vatheuer M, Maks C, Treadwell A, Allen R, et al. Acute mountain sickness is not altered by a high carbohydrate diet nor associated with elevated circulating cytokines. *Aviation, Space, and Environmental Medicine* 1997;**68**:499–503.

Utz 1970 *{published data only}*

Utz G, Schlierf G, Barth P, Linhart P, Wollenweber J. Prevention of acute mountain sickness using acetazolamide [Prophylaxe der akuten hohlenkrankheit mit acetazolamid]. *Munchener Medizinische Wochenschrift* 1970;**112**(23): 1122–4. PUBMED: 5467747]

Wang 1998 *{published data only}*

Wang W, Zhang X, Ma Y. Low-concentration nitrous oxide inhalation in the treatment of high-altitude pulmonary edema. *Chinese Journal of Tuberculosis and Respiratory Diseases* 1998;**21**(4):212–4.

Xiangjun 2014 *{published data only}*

Xiangjun L, Lan H. Inhaled budesonide for the prevention of acute mountain sickness in unacclimatization young men: a double-blind randomized controlled trial. *Journal of the American College of Cardiology* 2014;**64**(16 Suppl 1): C69.

References to ongoing studies

ChiCTR-TRC-13003319 *{published data only}*

ChiCTR-TRC-13003319. Oral zolpidem for improving sleep and then prevention of acute mountain sickness: a single centre, randomized, double-blind, controlled, prospective trial. www.chictr.org.cn/hvshowproject.aspx?id=6053 (first received 28th June 2013).

ChiCTR-TRC-13003590 *{published data only}*

ChiCTR-TRC-13003590. The Meaning of Intravenous Iron Supplementation In Acute Mountain Sickness: A Randomized, Double-Blinded, Placebo-Controlled Trial. <http://www.chictr.org.cn/showproj.aspx?proj=5970> (first published 23 June 2013).

NCT00886912 *{published data only}*

NCT00886912. Prevention of Acute Mountain Sickness (AMS) by Intermittent Hypoxic Training. <http://clinicaltrials.gov/show/NCT00886912> Date of registration: April 21th 2009.

NCT01606527 *{published data only}*

NCT01606527. Prospective, Double-blind, Randomized, Placebo-controlled Trial of Ibuprofen Versus Placebo for Prevention of Neurologic Forms of Altitude Sickness. <http://clinicaltrials.gov/show/NCT01606527> April 13th 2012.

NCT01682551 *{published data only}*

NCT01682551. Evaluation of the Prevention and Treatment Effects of Chinese Medicine on High Altitude Illness. <http://clinicaltrials.gov/show/NCT01682551> September 5th 2012.

NCT01794078 *{published data only}*

NCT01794078. A Randomized, 4-Sequence, Double-Blind Study to Test the Safety of Combined Dosing With Aminophylline and Ambrisentan in Exercising Healthy Human Volunteers at Simulated High Altitude. <https://clinicaltrials.gov/show/NCT01794078> February 10th 2013.

NCT01993667 *{published data only}*

NCT01993667. Acetazolamide for the Prevention of High Altitude Illness: a Comparison of Dosing. <https://clinicaltrials.gov/show/NCT01993667> November 12th 2013.

NCT02244437 *{published data only}*

NCT02244437. Ibuprofen vs Acetaminophen in the Prevention of Acute Mountain Sickness: A Double Blind, Randomized Controlled Trial. <https://clinicaltrials.gov/show/NCT02244437> September 14th 2014.

NCT02450968 *{published data only}*

NCT02450968. Dexamethasone for Prophylaxis of Acute Mountain Sickness in Patients With Chronic Obstructive Pulmonary Disease Travelling to Altitude. <https://clinicaltrials.gov/show/NCT02450968> May 18th 2015.

NCT02604173 *{published data only}*

NCT02604173. A Randomized Controlled Trial of Altitude Sickness Prevention and Efficacy of Comparative Treatments. <https://clinicaltrials.gov/show/NCT02604173> November 6th 2015.

NCT02811016 *{published data only}*

NCT02811016. Effect of Inhaled Budesonide on the Incidence and Severity of Acute Mountain Sickness at 4559 m. <https://clinicaltrials.gov/show/NCT02811016> June 14th 2016.

NCT02941510 *{published data only}*

NCT02941510. Inhaled Budesonide for Altitude Illness Prevention. <https://clinicaltrials.gov/show/NCT02941510> October 19th 2016.

Additional references

Adams 2004

Adams J. Ginkgo biloba and acetazolamide for acute mountain sickness: exclusion of high risk, low status groups

- perpetuates discrimination and inequalities. *BMJ* 2004;**329**(7458):171. MEDLINE: 15258082
- Anonymous 1892**
Anonymous. Mountain sickness. *British Medical Journal* 1892;**1**(1633):829.
- Bailey 2009a**
Bailey DM, Bärtsch P, Knauth M, Baumgartner RW. Emerging concepts in acute mountain sickness and high-altitude cerebral edema: from the molecular to the morphological. *Cellular and Molecular Life Sciences* 2009;**66**(22):3583–94. MEDLINE: 19763397
- Balslem 2011**
Balslem H, Helfand M, Schunemann HJ, Oxman AD, Kunz R, Brozek J, et al. GRADE guidelines: 3. Rating the quality of evidence. *Journal of Clinical Epidemiology* 2011;**64**(4):401–6. [PUBMED: 21208779]
- Bates 2007**
Bates MG, Thompson AA, Baillie JK. Phosphodiesterase type 5 inhibitors in the treatment and prevention of high altitude pulmonary edema. *Current Opinion in Investigational Drugs* 2007;**8**(3):226–31. MEDLINE: 17408118
- Brok 2009**
Brok J, Thorlund K, Wetterslev J, Gluud C. Apparently conclusive meta-analyses may be inconclusive—Trial sequential analysis adjustment of random error risk due to repetitive testing of accumulating data in apparently conclusive neonatal meta-analyses. *International Journal of Epidemiology* 2009;**38**(1):287–98. [PUBMED: 18824466]
- Bärtsch 1992**
Bärtsch P. Treatment of high altitude diseases without drugs. *International Journal of Sports Medicine* 1992;**13 Suppl 1**: 71–4. MEDLINE: 1483799
- Bärtsch 2004**
Bärtsch P, Bailey DM, Berger MM, Knauth M, Baumgartner RW. Acute mountain sickness: controversies and advances. *High Altitude Medicine & Biology* 2004;**5**(2):110–2. MEDLINE: 15265333
- Bärtsch 2007**
Bärtsch P, Gibbs JS. Effect of altitude on the heart and the lungs. *Circulation* 2007;**116**(19):2191–202. MEDLINE: 17984389
- CATMAT 2007**
Committee to Advise on Tropical Medicine and Travel (CATMAT). Statement on high-altitude illnesses. An Advisory Committee Statement (ACS). *Canada Communicable Disease Report (Relevé des maladies transmissibles au Canada)* 2007;**33**(ACS-5):1–20. MEDLINE: 17520777
- Chandramoorthi 2008**
Chandramoorthi GD, Piramanayagam S, Marimuthu P. An insilico approach to high altitude pulmonary edema - Molecular modeling of human beta2 adrenergic receptor and its interaction with salmeterol & nifedipine. *Journal of Molecular Modeling* 2008;**14**(9):849–56. MEDLINE: 18512086
- CTU 2011 [Computer program]**
Copenhagen Trial Unit. Trial Sequential Analysis Version. Copenhagen: Copenhagen Trial Unit, 2011.
- DailyMed**
DailyMed. dailymed.nlm.nih.gov/dailymed/about.cfm 2011; Vol. (Accessed 25 May 2011).
- Dehnert 2010**
Dehnert C, Bärtsch P. Can patients with coronary heart disease go to high altitude?. *High Altitude Medicine & Biology* 2010;**11**(3):183–8. [PUBMED: 20919884]
- DerSimonian 1986**
DerSimonian R, Laird N. Meta-analysis in clinical trials. *Controlled Clinical Trials* 1986;**7**(3):177–88.
- Dumont 2000**
Dumont L, Mardirosoff C, Tramèr MR. Efficacy and harm of pharmacological prevention of acute mountain sickness: quantitative systematic review. *BMJ* 2000;**321**(7256): 267–72. MEDLINE: 10915127
- Elbourne 2002**
Elbourne DR, Altman DG, Higgins JB, Curtin F, Worthington HV, Vail A. Meta-analyses involving cross-over trials: methodological issues. *International Journal of Epidemiology* 2002;**31**(1):140–9. [PUBMED: 11914310]
- Elphick 2004**
Elphick HL, Elphick DA. Ginkgo biloba and acetazolamide for acute mountain sickness: bias in participants may underestimate effectiveness of agents. *BMJ* 2004;**329**(7458):172. MEDLINE: 15258084
- Guyatt 2008**
Guyatt GH, Oxman AD, Kunz R, Vist GE, Falck-Ytter Y, Schunemann HJ. What is “quality of evidence” and why is it important to clinicians. *BMJ* 2008;**336**(7651):995–8. MEDLINE: 18456631
- Guyatt 2011a**
Guyatt G, Oxman AD, Akl EA, Kunz R, Vist G, Brozek J, et al. GRADE guidelines: 1. Introduction-GRADE evidence profiles and summary of findings tables. *Journal of Clinical Epidemiology* 2011;**64**(4):383–94. [PUBMED: 21195583]
- Guyatt 2011b**
Guyatt GH, Oxman AD, Kunz R, Atkins D, Brozek J, Vist G, et al. GRADE guidelines: 2. Framing the question and deciding on important outcomes. *Journal of Clinical Epidemiology* 2011;**64**(4):395–400. [PUBMED: 21194891]
- Guyatt 2011c**
Guyatt GH, Oxman AD, Kunz R, Brozek J, Alonso-Coello P, Rind D, et al. GRADE guidelines 6. Rating the quality of evidence-imprecision. *Journal of Clinical Epidemiology* 2011;**64**(12):1283–93. [PUBMED: 21839614]
- Guyatt 2011d**
Guyatt GH, Oxman AD, Kunz R, Woodcock J, Brozek J, Helfand M, et al. GRADE guidelines: 8. Rating the quality of evidence-indirectness. *Journal of Clinical Epidemiology* 2011;**64**(12):1303–10. [PUBMED: 21802903]

Guyatt 2011e

Guyatt GH, Oxman AD, Kunz R, Woodcock J, Brozek J, Helfand M, et al. GRADE guidelines: 7. Rating the quality of evidence-inconsistency. *Journal of Clinical Epidemiology* 2011;**64**(12):1294–302. [PUBMED: 21803546]

Guyatt 2011f

Guyatt GH, Oxman AD, Montori V, Vist G, Kunz R, Brozek J, et al. GRADE guidelines: 5. Rating the quality of evidence-publication bias. *Journal of Clinical Epidemiology* 2011;**64**(12):1277–82. [PUBMED: 21802904]

Guyatt 2011g

Guyatt GH, Oxman AD, Sultan S, Glasziou P, Akl EA, Alonso-Coello P, et al. GRADE guidelines: 9. Rating up the quality of evidence. *Journal of Clinical Epidemiology* 2011;**64**(12):1311–6. [PUBMED: 21802902]

Guyatt 2011h

Guyatt GH, Oxman AD, Vist G, Kunz R, Brozek J, Alonso-Coello P, et al. GRADE guidelines: 4. Rating the quality of evidence-study limitations (risk of bias). *Journal of Clinical Epidemiology* 2011;**64**(4):407–15. [PUBMED: 21247734]

Hackett 1992

Hackett PH, Roach RC, Hartig GS, Greene ER, Levine BD. The effect of vasodilators on pulmonary hemodynamics in high altitude pulmonary edema: a comparison. *International Journal of Sports Medicine* 1992;**13 Suppl 1**: 68–71. MEDLINE: 1483798

Hackett 1999

Hackett PH. High altitude cerebral edema and acute mountain sickness. A pathophysiology update. *Advances in Experimental Medicine and Biology* 1999;**474**:23–45. [PUBMED: 10634991]

Hackett 2004

Hackett PH, Roach RC. High altitude cerebral edema. *High Altitude Medicine & Biology* 2004;**5**(2):136–46. MEDLINE: 15265335

Higgins 2003

Higgins JPT, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ* 2003;**327** (7414):557–60. MEDLINE: 12958120

Higgins 2011

Higgins JP, Green S, editor(s). *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 (updated March 2011). The Cochrane Collaboration, 2011. Available from handbook.cochrane.org.

Hozo 2005

Hozo SP, Djulbegovic B, Hozo I. Estimating the mean and variance from the median, range, and the size of a sample. *BMC Medical Research Methodology* 2005;**5**:13. [PUBMED: 15840177]

Hussain 2004

Hussain MM, Aslam M. Hypoxia and pulmonary acclimatisation at 4578 m altitude: the role of acetazolamide and dexamethasone. *Journal of the Pakistan Medical Association* 2003;**53**(10):451–8. MEDLINE: 14696884

Höhne 2007

Höhne C, Pickerodt PA, Francis RC, Boemke W, Swenson ER. Pulmonary vasodilation by acetazolamide during hypoxia is unrelated to carbonic anhydrase inhibition. *American Journal of Physiology. Lung Cellular and Molecular Physiology* 2007;**292**(1):L178–84. [PUBMED: 16936246]

Imray 2010

Imray C, Wright A, Subudhi A, Roach R. Acute mountain sickness: pathophysiology, prevention, and treatment. *Progress in Cardiovascular Diseases* 2010;**52**(6):467–84. MEDLINE: 20417340

Kallenberg 2007

Kallenberg K, Bailey DM, Christ S, Mohr A, Roukens R, Menold E, et al. Magnetic resonance imaging evidence of cytotoxic cerebral edema in acute mountain sickness. *Journal of Cerebral Blood Flow and Metabolism* 2007;**27**(5): 1064–71. [PUBMED: 17024110]

Kayser 2012

Kayser B, Dumont L, Lysakowski C, Combescure C, Haller G, Tramer MR. Reappraisal of acetazolamide for the prevention of acute mountain sickness: a systematic review and meta-analysis. *High Altitude Medicine & Biology* 2012;**13**(2):82–92. [PUBMED: 22724610]

Kleinsasser 2002

Kleinsasser A, Loeckinger A. Are sildenafil and theophylline effective in the prevention of high-altitude pulmonary edema?. *Medical Hypotheses* 2002;**59**(2):223–5. MEDLINE: 12208214

Lan 1983

Lan KKG, Demets D. Discrete sequential boundaries for clinical trials. *Biometrika* 1983;**70**(3):659–63.

Leaf 2007

Leaf DE, Goldfarb DS. Mechanisms of action of acetazolamide in the prophylaxis and treatment of acute mountain sickness. *Journal of Applied Physiology* 2007;**102** (4):1313–22. MEDLINE: 17023566

Leissner 2009

Leissner KB, Mahmood FU. Physiology and pathophysiology at high altitude: considerations for the anesthesiologist. *International Journal of Anesthesia* 2009;**23** (4):543–53. MEDLINE: 19921365

Low 2012

Low EV, Avery AJ, Gupta V, Schedlbauer A, Grocott MP. Identifying the lowest effective dose of acetazolamide for the prophylaxis of acute mountain sickness: systematic review and meta-analysis. *BMJ (Clinical research ed.)* 2012;**345**: e6779. [PUBMED: 23081689]

Luks 2008a

Luks AM. Do we have a “best practice” for treating high altitude pulmonary edema?. *High Altitude Medicine & Biology* 2008;**9**(2):111–4. MEDLINE: 18578641

Luks 2008b

Luks AM, Swenson ER. Medication and dosage considerations in the prophylaxis and treatment of high-altitude illness. *Chest* 2008;**133**(3):744–55. MEDLINE: 18321903

- Luks 2010**
Luks AM, McIntosh SE, Grissom CK, Auerbach PS, Rodway GW, Schoene RB, Wilderness Medical Society. Wilderness Medical Society consensus guidelines for the prevention and treatment of acute altitude illness. *Wilderness & Environmental Medicine* 2010;**21**(2):146–55. MEDLINE: 20591379
- Luks 2014**
Luks AM, McIntosh SE, Grissom CK, Auerbach PS, Rodway GW, Schoene RB, et al. Wilderness Medical Society practice guidelines for the prevention and treatment of acute altitude illness: 2014 update. *Wilderness & Environmental Medicine* 2014;**25**(4 Suppl):S4–14. [PUBMED: 25498261]
- Maggiorini 1998**
Maggiorini M, Müller A, Hofstetter D, Bärtsch P, Oelz O. Assessment of acute mountain sickness by different score protocols in the Swiss Alps. *Aviation, Space, and Environmental Medicine* 1998;**69**(12):1186–92. [PUBMED: 9856545]
- Maggiorini 2010**
Maggiorini M. Prevention and treatment of high-altitude pulmonary edema. *Progress in Cardiovascular Diseases* 2010;**52**(6):500–6. MEDLINE: 20417343
- Monge 1942**
Monge C. Life in the Andes and chronic mountain sickness. *Science* 1942;**95**(2456):79–84. MEDLINE: 17757318
- Murata 2004**
Murata T, Hori M, Sakamoto K, Karaki H, Ozaki H. Dexamethasone blocks hypoxia-induced endothelial dysfunction in organ-cultured pulmonary arteries. *American Journal of Respiratory and Critical Care Medicine* 2004;**170**(6):647–55. MEDLINE: 15184203
- Murata 2005**
Murata T, Suzuki N, Yamawaki H, Sato K, Hori M, Karaki H, et al. Dexamethasone prevents impairment of endothelium-dependent relaxation in arteries cultured with fetal bovine serum. *European Journal of Pharmacology* 2005;**515**(1-3):134–41. MEDLINE: 15907323
- Naeije 2010**
Naeije R. Physiological adaptation of the cardiovascular system to high altitude. *Progress in Cardiovascular Diseases* 2010;**52**(6):456–66. MEDLINE: 20417339
- Ozaki 2001**
Ozaki M, Kawashima S, Yamashita T, Ohashi Y, Rikitake Y, Inoue N, et al. Reduced hypoxic pulmonary vascular remodeling by nitric oxide from the endothelium. *Hypertension* 2001;**37**(2):322–7. MEDLINE: 11230292
- Palmer 2010**
Palmer BF. Physiology and pathophysiology with ascent to altitude. *American Journal of the Medical Sciences* 2010;**340**(1):69–77. MEDLINE: 20442648
- Pandit 2014**
Pandit A, Karmacharya P, Pathak R, Giri S, Aryal MR. Efficacy of NSAIDs for the prevention of acute mountain sickness: a systematic review and meta-analysis. *Journal of Community Hospital Internal Medicine Perspectives* 2014;**4**(4):1–7. [PUBMED: 25317267]
- Paralíkar 2010**
Paralíkar SJ, Paralíkar JH. High-altitude medicine. *Indian Journal of Occupational and Environmental Medicine* 2010;**14**(1):6–12. MEDLINE: 20808661
- Pasha 2010**
Pasha MA, Newman JH. High-altitude disorders: pulmonary hypertension: pulmonary vascular disease: the global perspective. *Chest* 2010;**137** Suppl(6):13–9. MEDLINE: 20522576
- RevMan 5.3 [Computer program]**
The Nordic Cochrane Centre, The Cochrane Collaboration. Review Manager (RevMan). Version 5.3.. Copenhagen: The Nordic Cochrane Centre, The Cochrane Collaboration, 2014.
- Ri-Li 2003**
Ri-Li G, Chase PJ, Witkowski S, Wyrick BL, Stone JA, Levine BD, et al. Obesity: associations with acute mountain sickness. *Annals of Internal Medicine* 2003;**139**(4):253–7. [PUBMED: 12965980]
- Richalet 2005**
Richalet JP, Grataudour P, Robach P, Pham I, Déchaux M, Joncquiert-Latarjet A, et al. Sildenafil inhibits altitude-induced hypoxemia and pulmonary hypertension. *American Journal of Respiratory and Critical Care Medicine* 2005;**171**(3):275–81. MEDLINE: 15516532
- Ritchie 2012**
Ritchie ND, Baggott AV, Andrew Todd WT. Acetazolamide for the prevention of acute mountain sickness—a systematic review and meta-analysis. *Journal of Travel Medicine* 2012;**19**(5):298–307. [PUBMED: 22943270]
- Rodríguez de Romo 2002**
Rodríguez de Romo AC. Daniel Vergara Lope and Carlos Monge Medrano: two pioneers of high altitude medicine. *High Altitude Medicine & Biology* 2002;**3**(3):299–309. MEDLINE: 12396886
- Scherrer 2010**
Scherrer U, Rexhaj E, Jayet PY, Allemann Y, Sartori C. New insights in the pathogenesis of high-altitude pulmonary edema. *Progress in Cardiovascular Diseases* 2010;**52**(6):485–92. MEDLINE: 20417341
- Schneider 2002**
Schneider M, Bernasch D, Weymann J, Holle R, Bärtsch P. Acute mountain sickness: influence of susceptibility, preexposure, and ascent rate. *Medicine and Science in Sports and Exercise* 2002;**34**(12):1886–91. [PUBMED: 12471292]
- Schoene 2004**
Schoene RB. Unraveling the mechanism of high altitude pulmonary edema. *High Altitude Medicine & Biology* 2004;**5**(2):125–35. MEDLINE: 15265334
- Schoene 2008**
Schoene RB. Illnesses at high altitude. *Chest* 2008;**134**(2):402–16. MEDLINE: 18682459

Schoonman 2008

Schoonman GG, Sandor PS, Nirkko AC, Lange T, Jaermann T, Dydak U, et al. Hypoxia-induced acute mountain sickness is associated with intracellular cerebral edema: a 3 T magnetic resonance imaging study. *Journal of Cerebral Blood Flow and Metabolism* 2008;**28**(1):198–206. [PUBMED: 17519973]

Seupaul 2012

Seupaul RA, Welch JL, Malka ST, Emmett TW. Pharmacologic prophylaxis for acute mountain sickness: a systematic shortcut review. *Annals of Emergency Medicine* 2012;**59**(4):307–317.e1. [PUBMED: 22153998]

Stedman 2011

Stedman MR, Curtin F, Elbourne DR, Kesselheim AS, Brookhart MA. Meta-analyses involving cross-over trials: methodological issues. *International Journal of Epidemiology* 2011;**40**(6):1732–4. [PUBMED: 20026595]

Stream 2008

Stream JO, Grissom CK. Update on high-altitude pulmonary edema: pathogenesis, prevention, and treatment. *Wilderness & Environmental Medicine* 2008;**19**(4):293–303. MEDLINE: 19099331

Swenson 2006

Swenson ER. Carbonic anhydrase inhibitors and hypoxic pulmonary vasoconstriction. *Respiratory Physiology & Neurobiology* 2006;**151**(2-3):209–16. [PUBMED: 16376158]

Swenson 2007

Swenson E, Teppema L. Prevention of acute mountain sickness by acetazolamide: as yet an unfinished story. *Journal of Applied Physiology* 2007;**102**(4):1305–7. MEDLINE: 17194729

Tang 2014

Tang E, Chen Y, Luo Y. Dexamethasone for the prevention of acute mountain sickness: systematic review and meta-analysis. *International Journal of Cardiology* 2014;**173**(2):133–8. [PUBMED: 24679688]

Thorlund 2009

Thorlund K, Devereaux PJ, Wetterslev J, Guyatt G, Ioannidis JP, Thabane L, et al. Can trial sequential monitoring boundaries reduce spurious inferences from meta-analyses?. *International Journal of Epidemiology* 2009;**38**(1):276–86. [PUBMED: 18824467]

Tintinger 2010

Tintinger GR, Feldman C, Theron AJ, Anderson R. Montelukast: more than a cysteinyl leukotriene receptor antagonist?. *Scientific World Journal* 2010;**10**:2403–13. [PUBMED: 21170491]

Wan 2014

Wan X, Wang W, Liu J, Tong T. Estimating the sample mean and standard deviation from the sample size, median, range and/or interquartile range. *BMC Medical Research Methodology* 2014;**14**:135. [PUBMED: 25524443]

Wetterslev 2008

Wetterslev J, Thorlund K, Brok J, Gluud C. Trial sequential analysis may establish when firm evidence is reached in cumulative meta-analysis. *Journal of Clinical Epidemiology* 2008;**61**(1):64–75. [PUBMED: 18083463]

Wilson 2009

Wilson MH, Newman S, Imray CH. The cerebral effects of ascent to high altitudes. *Lancet Neurology* 2009;**8**(2):175–91. MEDLINE: 19161909

Windsor 2009

Windsor JS, Firth PG, Grocott MP, Rodway GW, Montgomery HE. Mountain mortality: a review of deaths that occur during recreational activities in the mountains. *Postgraduate Medical Journal* 2009;**85**(1004):316–21. [PUBMED: 19528307]

Wright 2008

Wright A, Brearey S, Imray C. High hopes at high altitudes: pharmacotherapy for acute mountain sickness and high-altitude cerebral and pulmonary oedema. *Expert Opinion on Pharmacotherapy* 2008;**9**(1):119–27. MEDLINE: 18076343

Yang 2007

Yang B, Wang GY, Chen B, Qin RB, Xi SL, Chen L. Anti-hypoxia and anti-oxidation effects of aminophylline on human with acute high-altitude exposure. *Chinese Medical Sciences Journal* 2007;**22**(1):62–5. [PUBMED: 17441321]

Zafren 2014

Zafren K. Prevention of high altitude illness. *Travel Medicine and Infectious Disease* 2014;**12**(1):29–39. [PUBMED: 24393671]

Zhao 2001

Zhao L, Mason NA, Morrell N, Kojonazarov B, Sadykov A, Maripov A, et al. Sildenafil inhibits hypoxia-induced pulmonary hypertension. *Circulation* 2001;**104**(4):424–8. MEDLINE: 11468204

References to other published versions of this review**Martí-Carvajal 2012**

Martí-Carvajal Arturo J, Hidalgo R, Simancas-Racines D. Interventions for preventing high altitude illness. *Cochrane Database of Systematic Reviews* 2012, Issue 4. [DOI: 10.1002/14651858.CD009761]

* Indicates the major publication for the study

CHARACTERISTICS OF STUDIES

Characteristics of included studies *[ordered by study ID]*

Anonymous 1981

Methods	<ol style="list-style-type: none"> 1. Design: Parallel, 2 arms 2. Country: Ecuador 3. Multisite: No 4. International: No 5. Treatment duration: 3 days 6. Follow-up: 8 days 7. Rate of ascent: unclear 8. Final altitude reached: 5000 metres 9. AMS scale: clinical arbitrary score (0 - 100) 10. Randomization unit: participants 11. Analysis unit: Groups
Participants	<ol style="list-style-type: none"> 1. 20 participants enrolled (age 20 - 52 , all normally resided at less than 200 metres, all medically qualified) <p>Randomized to:</p> <p>Acetazolamide group (n = 10, 50%)</p> <p>Placebo group (n = 10, 50%)</p> <ol style="list-style-type: none"> 2. No participant randomized was excluded 3. No participant was lost to follow-up 4. Main characteristics of participants: <p>Age: 20 - 52 years</p> <p>100% men</p> <p>History of AMS: not stated</p> <p>Percentage/number type of HAI reported: not reported</p>
Interventions	<ol style="list-style-type: none"> 1. Acetazolamide group: acetazolamide 500 mg/day for 3 days, oral 2. Placebo group (control): unclear
Outcomes	<p>This RCT did not specify its primary or secondary outcomes</p> <ol style="list-style-type: none"> 1. Assessment of Acute Mountain Sickness by clinical interview: arbitrary scores (0 - 100) 2. Peer review: rank order according to subjective impression 3. Blood gas measurements included: hydrogen ion concentration, oxygen tension and carbon dioxide tension
Notes	<ol style="list-style-type: none"> 1. Trial Registration: Not stated 2. Funder: Boehringer ingelheim ltda. Financial Mathematics Ltd. Geigy pharmaceuticals, laboratoire d'étude de recherches scientifiques lederle pharmaceuticals, the Arthur Thompson Trust fund and West Midlands Regional Health Authority and many other companies that gave financial aid 3. Role of Funder: Not stated 4. A priori sample size estimation: No 5. Conducted: Not stated 6. Declared conflicts of interest: Not reported

Anonymous 1981 (Continued)

<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote "...were randomly allocated..." (Page 181) The method of sequence generation was not specified
Allocation concealment (selection bias)	Unclear risk	The method of sequence generation was not specified
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Quote "Details of medication were concealed until after descent" (Page 181) There was insufficient information to assess whether blinding was likely to introduce bias in the results
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Incomplete outcome data (attrition bias) All outcomes	Low risk	No participants were reported as lost to follow-up
Selective reporting (reporting bias)	Low risk	Reporting bias was not detected
Other bias	Unclear risk	Possible industry bias. The trial is sponsored by the industry or has received other kind of for-profit support

ASCENT 2012

Methods	<ol style="list-style-type: none"> 1. Design: A randomized, doubled-blind, placebo-controlled trial. 2 arms: placebo group, ibuprofen group 2. Country: Nepal 3. Multisite: No 4. International: No 5. Treatment duration: 1 day 6. Intention-to-treat: Yes 7. Follow-up: 1 day after arrival 8. Rate of ascent: unclear 9. Final altitude reached: 4928 metres 10. AMS scale: Lake Louise AMS questionnaire (LLQ)
Participants	<ol style="list-style-type: none"> 1. 294 participants enrolled, 183 completed the entire protocol. 49 broke protocol, but allowed data collection; at the end 62 participants were lost to follow-up 2. 232 participants completed the study. (Healthy men and women, 37 ± 12 years),

	<p>recruited at 4280 or 4358 metres on the Everest approach: Placebo (109, 47%) Ibuprofen (123, 53%) 3. Main characteristics of participants: Age 36 ± 11 (placebo) 38 ± 12 (ibuprofen) Number/Percentage of women: 35 (32.4%) placebo, 46 (37.7%) ibuprofen Percentage/number history of AMS: 5/109 (4.7% placebo) 7/123 (5.8% ibuprofen) Percentage/number type of HAI reported: This study reported: Severe high altitude headache (HAH), evaluated by LLQ > 2: 16/109 (14.7% placebo) 6/123 (4.9% ibuprofen) AMS incidence evaluated by LLQ > 3: 44/109 (40.4% placebo), 30/123 (24.4% ibuprofen)</p>	
Interventions	<p>1. Placebo group: placebo 3 times daily orally for at least 3 doses before ascent 2. Ibuprofen group: 600 mg of ibuprofen 3 times a day orally for at least 3 doses before ascent 3. In both groups there was a period of acclimatization, approximately 3.4 ± 0.8 nights</p>	
Outcomes	<p>Primary outcome 1. Incidence of headache, severe headache, AMS, severe AMS. Measured by a value of 2, 3 or 5 respectively on the LLQ Secondary endpoint 1. SpO₂ decreased from baseline (end point SpO₂%)</p>	
Notes	<p>1. Trial Registration “Not stated” 2. Funder: Wellcome Trust UK 3. Role of Funder: Financial support 4. A priori sample size estimation: Yes, 164 participants (84 per arm) 5. Conducted: Enrolment took place between October and November 2009; start date not specified or when the study ended 6. Declared conflicts of interest: No</p>	
Risk of bias		
Bias	Authors’ judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: “Study medications were randomized via computer-generated code” (Page 308)
Allocation concealment (selection bias)	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias

ASCENT 2012 (Continued)

Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	21% of participants randomized were not analysed (62 participants). A modified ITT analysis was performed
Selective reporting (reporting bias)	Low risk	Selective reporting of information was not detected
Other bias	Low risk	No additional biases were identified

Banderet 1977

Methods	<ol style="list-style-type: none"> 1. Design: Paralell longitudinal study, 2 arms 2. Country: USA 3. Multisite: No 4. International: No 5. Treatment duration: 4 days 6. Follow-up: 3 weeks 7. Rate of ascent: unclear 8. Final altitude reached: 4300 metres 9. AMS scale: The Clyde Mood Scale and the High Altitude Symptom Questionnaire 10. Randomization unit: participant 11. Analysis unit: group
Participants	<p>35 participants enrolled (volunteers)</p> <p>Randomized to:</p> <p>Treatment group (n = 18, 51%)</p> <p>Placebo group (n = 17, 49%)</p> <p>Main characteristics of participants:</p> <p>Age: 19 - 28 years</p> <p>women/men: n = 16 / 19</p> <p>History of AMS: none</p>
Interventions	<ol style="list-style-type: none"> 1. Treatment group (intervention): acetazolamide 500 twice a day during last 2 days of staging at 1600 metres and during the first 2 days at 4300 metres 2. Placebo group (control): placebo 2 tablets twice a day each day throughout the study
Outcomes	This trial did not specify by primary or secondary outcomes Scores of Clyde Mood Scale (by symptom)
Notes	<ol style="list-style-type: none"> 1. Trial Registration: Not stated 2. Funder: Not stated 3. Role of funder: Not stated 4. A priori sample size estimation: No 5. Conducted: Not stated

Banderet 1977 (Continued)

6. Declared conflicts of interest: No		
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "Subjects were assigned randomly..." (Page 20) Insufficient information to score this item as low or high risk of bias
Allocation concealment (selection bias)	Unclear risk	Quote: "Subjects were assigned randomly..." (Page 20) Insufficient information to score this item as low or high risk of bias
Blinding of participants and personnel (performance bias) All outcomes	High risk	Quote: "All were informed initially that some of them would receive placebo tablets" (Page 23)
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Incomplete outcome data (attrition bias) All outcomes	Low risk	No participant were lost to follow-up
Selective reporting (reporting bias)	High risk	Important participant-important outcomes, such as adverse events, were not reported
Other bias	Low risk	No additional biases were identified

Bartsch 1991

Methods	<ol style="list-style-type: none"> 1. Design: Parallel (2 arms) 2. Country: Italy 3. Multisite: No 4. International: No 5. Treatment duration: 4 days 6. Follow-up: unclear 7. Rate of ascent: 155 metres/hour 8. Final altitude reached: 4559 metres 9. AMS scale: AMS score 10. Randomization unit: participants 11. Analysis unit: 2rroup
---------	---

Participants	<p>21 participants enrolled (mountaineers who had radiographically-documented episodes of high-altitude pulmonary oedema and who had continued alpine-style climbing to peaks above 4000 metres after there episodes of HAPE)</p> <p>Randomized to:</p> <p>Nifedipine (n = 10, 47.6%)</p> <p>Placebo (n = 11, 52.3%)</p> <p>6 participants left the study early</p> <p>1 person in placebo group left because of high-altitude pulmonary oedema on day 2</p> <p>3 people in placebo group left because of high-altitude pulmonary oedema on day 3</p> <p>1 person left the trial on the day of arrival at 4559 metres because of prodromal symptoms of pulmonary oedema</p> <p>4. Main characteristics of participants:</p> <p>Age (mean, range): placebo group 41 years, 20 - 58; nifedipine group 44 years, 23 - 62</p> <p>Number of women/men: 1 / 20</p> <p>Number of participants with 1 episode of HAPE: 6 placebo and 6 nifedipine</p>	
Interventions	<p>1. Nifedipine group: administration of slow-release preparation of nifedipine (Adalat, 20 mg) given at 10 p.m. on the third and second days before the ascent and at 8 a.m. and 10 p.m. on the day before. Starting on the day of the ascent the medication was taken 3 times daily (at 6 a.m., 2 p.m. and 10 p.m.)</p> <p>2. Placebo group (control): capsules taken orally 3 times daily for 4 days</p>	
Outcomes	<p>This trial did not specify by primary or secondary outcomes</p> <p>1. Presence of HAPE (documented by doppler. Susceptible mountaineers with documented histories of high-altitude pulmonary oedema)</p> <p>2. AMS score by clinical examination</p> <p>3. Blood and end expiratory gas analysis: SaO₂, PaO₂, PaCO₂, end exploratory PO₂</p>	
Notes	<p>1. Trial Registration: Not stated</p> <p>2. Funder: supported from the Swiss National Science Foundation</p> <p>3. Role of funder: Not stated</p> <p>4. A priori sample size estimation: No</p> <p>5. Conducted: Not stated</p> <p>6. Declared conflicts of interest: Not reported</p>	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote "...was assigned randomly.." (Page 1285) Insufficient information to score this item as low or high risk of bias
Allocation concealment (selection bias)	Unclear risk	Insufficient information to score this item as low or high risk of bias

Bartsch 1991 (Continued)

Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: "The clinical examination were directed toward the signs and symptoms of AMS, and were always performed by the same investigator, who was not aware of the subjects medication" (Page 1285)
Incomplete outcome data (attrition bias) All outcomes	High risk	5/11 placebo participants were not included in analyses of AMS scores at 4559 metres
Selective reporting (reporting bias)	High risk	Participant-important outcomes, such as adverse events, were not reported
Other bias	Low risk	No additional biases were identified

Basnyat 2003

Methods	<ol style="list-style-type: none"> 1. Design: Parallel, 2 arms 2. Country: Nepal 3. Multisite: No 4. International: No 5. Treatment : 2 or 3 days 6. Follow-up: unclear 7. Rate of ascent: 4.3 ± 1.1 days (range 3 - 6) 8. Final altitude reached: 4937 metres 9. AMS scale: The Lake Louise Acute Mountain Sickness Scoring System 10. Randomization unit: participant 11. Analysis unit: group
Participants	<p>197 participants enrolled (healthy non-Nepali male and female trekkers of > 18 years of age travelling between the villages from 4243 metres to 4937 metres)</p> <p>Exclusion criteria: Already had a diagnosis of AMS, HACE or HAPE; Had been on a high-altitude trek 2 weeks prior to this trek; Were not trekking directly to 4937 metres; Had taken acetazolamide or ginkgo biloba in the week prior to presentation; Has diabetes, serious heart or pulmonary disease or a sulfa allergy</p> <p>Randomized to:</p> <p>Acetazolamide (n = 96, 48.7%) Placebo (n = 101, 51.2%)</p> <p>2 . 42 participants lost at follow-up (they did not retrieve the questionnaire at Lobujr): Acetazolamide group (n = 22, 22.9%) Placebo group (n = 20, 19.8%)</p> <p>3. Main characteristics of participants :</p> <p>Age (mean; SD): acetazolamide group 35.8 ± 12.1; placebo group 33.9 ± 11.4 Percentage women/men: acetazolamide group 64.9% men/35.1% women; placebo</p>

	group 69.1% men/30.9% women O ₂ saturation at Periche: Acetazolamide group: 86.9 ± 3.9; placebo group: 86.9 ± 4	
Interventions	<p>1. Acetazolamide group (intervention): acetazolamide 125 mg twice daily for 2 to 3 days before the final evaluation at 4937 metres</p> <p>2. Placebo group (control): visually-matched placebo twice daily for 2 to 3 days before the final evaluation at 4937 metres</p> <p>Cointerventions: None stated</p>	
Outcomes	<p>Primary outcome:</p> <p>1. Incidence and severity of AMS by the LLQ Score at Lobuje</p> <p>Secondary outcomes:</p> <p>1. The presence or absence of high-altitude headache</p> <p>2. Diagnosis of HAPE or HACE</p> <p>3. Pulse oximetry differential between 4243 metres and 4937 metres</p> <p>4. Acute symptoms suggestive of infection at 4937 metres (sore throat, cough, sinusitis, diarrhoea)</p> <p>5. Incidence of paraesthesias</p> <p>6. Missed capsules</p>	
Notes	<p>1. Trial Registration: Not stated</p> <p>2. Funder: The Himalayan Rescue Association and Nepal International Clinic, Kathmandu, Nepal; and Deurali Pharmaceutical Company</p> <p>3. Role of funder: Donated the placebo capsules. Study administrators paid their own expenses</p> <p>4. A priori sample size estimation: No</p> <p>5. Conducted: November 1 to 22 of 2001</p> <p>6. Declared conflicts of interest: Yes, Page 52</p>	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote " Random allocation occurred on site, ..." (Page 47). The method of sequence generation was not specified
Allocation concealment (selection bias)	Low risk	Quote: "Randomization code was drawn up by a neutral party and was securely kept in Katmandu, completely unavailable to the study administrators" Page 47
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Quote " ...were visually indistinguishable, and neither study administrators nor participants knew the identity of the study capsules" (Page 47)

Basnyat 2003 (Continued)

Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Incomplete outcome data (attrition bias) All outcomes	High risk	The results were likely to be biased due to missing data 1. Losses at follow up in experimental group: 19.8%. (22/96) 2. Losses at follow up in control group: 22.9%. (20/101)
Selective reporting (reporting bias)	Low risk	Reporting bias was not detected
Other bias	Low risk	No additional biases were identified

Basnyat 2008

Methods	<ol style="list-style-type: none"> 1. Design: Parallel - 2 arms 2. Country: Nepal 3. Multisite: No 4. International: No 5. Treatment duration: 4 days 6. Follow-up: unclear 7. Rate of ascent: 36 hours to a maximum of 96 hours 8. Final altitude reached: 5000 metres 9. AMS scale: LLQ 10. Randomization unit: participant 11. Analysis unit: group
Participants	<p>364 healthy people were enrolled between the ages of 18 and 65, non-Nepali, without AMS or any concurrent illness, and not already taking acetazolamide or any other drug for the prevention of altitude illness</p> <p>Exclusion criteria: Mild AMS; significantly depressed oxygen saturation; women know to be pregnant or unable to exclude the possibility of being pregnant. or having missed menses by over 7 days; individuals with a known drug allergy to acetazolamide or other sulfa drugs; individuals who had spent 24 hours at altitude of 4500 metres or higher within the last 9 days; anyone know to have taken any of the following in the last 2 days: acetazolamide, steroids, theophylline or diuretics; individuals who had known intracranial space-occupying lesions or a history of elevated intracranial pressure</p> <p>Randomized to:</p> <p>Acetazolamide group (n = 187; 51.3%) Placebo group (n = 177; 48.6%)</p> <p>25 patients randomized were excluded due to:</p> <p>Dropped out or disqualified for stopping study drugs or taking non-study acetazolamide Acetazolamide group (n = 13; 6.9%) Placebo group (n = 12; 6.7%)</p> <p>3. Main characteristics of participants :</p> <p>Age (mean, SD): acetazolamide 37.9 ± 12.5; placebo 39.4 ± 12.1</p>

	<p>Percentage/number of women/men: acetazolamide: women 42.2% (79), men 57.8% (108); placebo: women 32.2% (57), men 67.8% (120)</p> <p>Percentage/number history of AMS: acetazolamide: 36.4% (68), placebo; 39.5% (70)</p> <p>Percentage/number Type of HAI reported: placebo 21.9%, acetazolamide group 10.2%</p> <p>Pulse oximetry (mean, SD): acetazolamide = 86.45 ± 3.39; placebo = 85.91 ± 4.08</p> <p>Heart rate (mean, SD): acetazolamide = 82.6 ± 12; placebo = 82.5 ± 12</p>
Interventions	<ol style="list-style-type: none"> 1. Acetazolamide group (intervention) = acetazolamide tablets 250 mg twice day for 4 days 2. Placebo group (control) = visually identical-appearing placebo tablets twice day for 4 days
Outcomes	<p>Primary outcomes:</p> <ol style="list-style-type: none"> 1. HAPE diagnosis (signs and symptoms): AMS (LLS ≥ 3, at least 1 symptom) + 2 signs and 2 symptoms of pulmonary involvement <p>Determination of pulmonary artery systolic pressure</p> <p>Secondary outcomes</p> <ol style="list-style-type: none"> 1. Pulse oxygen saturation of < 70% in participants meeting HAPE diagnosis 2. Incidence of AMS, HAPE and HACE
Notes	<ol style="list-style-type: none"> 1. Trial Registration: Not stated 2. Funder: Sonosite Micromaxx, Wellcome Trust of Great Britain 3. Role of funder: Provision of ultrasonographer 4. A priori sample size estimation: Yes.(Page 211) 5. Conducted: October and November, 2006 6. Declared conflicts of interest: Yes (Page 215)

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: " Computer generated randomization of commercial pharmaceutical grade acetazolamide and placebo were carried out by Deuralu Janata pharmaceuticals" (Page 210)
Allocation concealment (selection bias)	Low risk	3 sealed master lists of the randomization code were held by the manufacturer and independent clinicians. Only opened by an independent clinician when there was a concern (Page 211)
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Quote: "The code was only to be opened during the trial by an independent clinician who was not a study author when there was concern of allergic reaction or any other adverse event (...)" (Page 211) Study drug and placebo had a visually iden-

Basnyat 2008 (Continued)

		tical appearance (Page 210)
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Incomplete outcome data (attrition bias) All outcomes	Low risk	Missing data were unlikely to make treatment effects depart from plausible values Loss to follow-up in experimental group: 6.95% (13/187) Loss to follow-up in control group: 6.77% (12/177)
Selective reporting (reporting bias)	High risk	Participant-important outcomes, such as adverse events, were not reported
Other bias	Low risk	No additional biases were identified

Basu 2002a

Methods	<ol style="list-style-type: none"> 1. Design: Parallel, 5 arms 2. Country: India 3. Multisite: No 4. International: No 5. Treatment duration: 5 days 6. Follow-up: 7 days 7. Rate of ascent: unclear 8. Final altitude reached: 3450 metres 9. AMS scale: Lake Louise AMS scoring system 10. Randomization unit: participant 11. Analysis unit: group
Participants	<p>50 healthy men enrolled (none of them taking any medication and had not taken steroid preparations; excluded if any disorders or contraindication to steroid therapy)</p> <p>Patients randomized to:</p> <p>Group I (n = 10, 20%) Group II (n = 10, 20%) Group III (n = 10, 20%) Group IV (n = 10, 20%) Group V (n = 10, 20%)</p> <p>Unclear if any people were excluded</p> <p>Unclear if participants were lost to follow-up (See Table 2, only 9 participants in dexamethasone group)</p> <p>Main characteristics of participants:</p> <p>Age: 19 - 24 years for all participants Percentage of men: 100% Body weight: 55 - 70 kg History of AMS: Not stated</p>

Basu 2002a (Continued)

Interventions	<ol style="list-style-type: none"> 1. Group I (intervention): prednisolone 10 mg, oral single dose a day for 5 days 2. Group II (intervention): prednisolone 20 mg oral single dose a day for 5 days 3. Group III (intervention): prednisolone 40 mg oral single dose a day for 5 days 4. Group IV (intervention): dexamethasone IV 0.5 mg dose a day for 5 days 5. Group V (control): placebo once a day in the morning at 08:00 hours before breakfast Coninterventions: None declared	
Outcomes	This RCT did not specify by primary or secondary outcomes <ol style="list-style-type: none"> 1. Symptoms of AMS. Score of AMS 2. Physiological variables: BP, SaO₂, heart rate 3. Hormonal estimations: cortisol and ACTH 	
Notes	<ol style="list-style-type: none"> 1. Trial Registration: Not stated 2. Funder: Not stated 3. Role of funder: Not stated 4. A priori sample size estimation: No 5. Declared conflicts of interest: Not stated 	
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote "...randomized trial..." (Page 762) The method of sequence generation was not specified
Allocation concealment (selection bias)	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias Quote "The placebo and drugs looked alike..." (Page 762)
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Selective reporting (reporting bias)	High risk	Participant-important outcomes, such as adverse events, were not reported
Other bias	Low risk	No additional biases were identified

Methods	<ol style="list-style-type: none"> 1. Design: Randomized placebo controlled trial 2. Country: India 3. Multisite: No 4. International: No 5. Treatment duration: unclear 6. Follow-up: followed over 1 week at high altitude and 2 weeks on return to sea level 7. Rate of ascent: 3450 metres by air 8. Final altitude reached: 3450 metres 9. AMS scale: Lake Louise AMS scoring system 10. Randomization unit: Participants 11. Analysis unit: Group 	
Participants	<p>40 participants enrolled. Those selected had no contraindication to steroid therapy and had not taken any steroid preparations within the preceding year</p> <p>Randomized to:</p> <p>20 prednisolone (50%) Group I</p> <p>20 placebo (50%) Group II</p> <p>Number randomized who were excluded: not reported</p> <p>Participants lost to follow-up: not reported</p> <p>Main characteristics of participants:</p> <p>Age: 19 - 26 years</p> <p>Percentage/number of women/men: 40 men (100%)</p> <p>Percentage/number history of AMS: In placebo group 11 participants showed AMS</p> <p>prednisolone group unclear</p>	
Interventions	<p>Group I: received prednisolone 20 mg</p> <p>Group II: received placebo</p> <p>Once a day at 8:00 a.m. before breakfast for 2 days prior to induction, and for 3 days on arrival at high altitude</p>	
Outcomes	<p>This trial did not specify by primary or secondary outcomes</p> <ol style="list-style-type: none"> 1. AMS scores 2. Circulatory levels of ACTH, cortisol, epinephrine and norepinephrine 	
Notes	<ol style="list-style-type: none"> 1. Trial Registration: Not stated 2. Funder: Not stated 3. Role of funder: Not stated 4. A priori sample size estimation: No 5. Conducted: Unclear 6. Declared conflicts of interest: No reported 	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "The subjects were randomly divided into two groups of twenty each." (Page 319) Insufficient information to score this item

Basu 2002b (Continued)

		as low or high risk of bias
Allocation concealment (selection bias)	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	No participants were reported as lost to follow-up
Selective reporting (reporting bias)	High risk	Participant-important outcomes, such as adverse events, were not reported
Other bias	Unclear risk	No additional biases were identified

Bates 2011

Methods	<ol style="list-style-type: none"> 1. Design: Parallel design, 2 arms 2. Country: Bolivia 3. Multisite: No 4. International: No 5. Treatment duration: 12 days 6. Follow-up: 12 days 7. Rate of ascent: unclear 8. Final altitude reached: 5200 metres 9. AMS scale: Lake Louise Consensus Symptom Score 10. Randomization unit: participant 11. Analysis unit: participant
Participants	<p>This trial was conducted concurrently with a similar trial of an oral antioxidant vitamin supplement, addressing a different aspect of altitude illness (information not included in this review)</p> <p>62 healthy native lowlanders</p> <p>Randomized to:</p> <p>Sildenafil 20 (32.3%)</p> <p>Placebo 42 (67.7 %)</p> <p>1 participant in the placebo group developed HAPE while at 3650 metres, did not ascend to the high altitude laboratory, and was excluded from the trial. Throughout the trial, it proved technically impossible to obtain satisfactory PASP measurements from 7 participants (all from the placebo group) and they were excluded from all the PASP analyses. 12 more participants were evacuated from the high altitude laboratory because of severe symptoms of AMS and thereby withdrew from the trial (5 in the sildenafil group and 7 in the placebo group). PASP and AMS data for these participants were included</p>

	<p>until their evacuation. 8 further individual PASP measurements (all at 5200 metres, 2 in the sildenafil group, 6 in the placebo group) were rejected as technically unsatisfactory following independent review after the expedition. All AMS data and PASP data from these participants at different time points are included in the analysis</p> <p>Main characteristics of participants: Male: placebo: 62% (26/42), sildenafil 55% (11/20) Age mean: placebo 21.5 ± 2.7 years, sildenafil 21.2 ± 3 years History of AMS: Not stated</p>
Interventions	<p>Sildenafil Citrate group (intervention): 50 mg/day before ascending, then 50 mg/3 times a day orally, during trek Placebo group: Unclear</p>
Outcomes	<p>Primary outcomes: PASP assessed by transthoracic echocardiography at 6 hours, 3 days, and 1 week following high-altitude exposure at 5200 metres Secondary outcomes were oxygen saturations by pulse oximetry, severity of AMS using the LLS for the first 7 days at high altitude, and proportion of participants with LLS > 3 on day 2 at 5200 metres</p>
Notes	<ol style="list-style-type: none"> 1. Trial Registration: NCT00627965 2. Funder: None 3. A priori sample size estimation: Stated on page 209 4. Conducted: Not stated 5. Declared conflicts of interest: No, stated on page 213

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "All 103 Apex 2 expedition participants were randomly assigned to three groups using a computer programme operated by an independent statistician, as determined..." (Page 208)
Allocation concealment (selection bias)	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Quote: "All researchers and participants were unaware of the group assignments and independent of the individual responsible for the randomization process" "Supplies of sildenafil and masked placebo were obtained directly from the manufacturers. Packs of these tablets were identically packaged in the UK under the supervision of the head of clinical trial facility, and distributed to the trial participants for per-

Bates 2011 (Continued)

		sonal administration.” (Page 208)
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Incomplete outcome data (attrition bias) All outcomes	Low risk	8 participants lost to follow-up at day 2 (13%)
Selective reporting (reporting bias)	High risk	Patient-important outcomes, such as adverse events, were not reported
Other bias	Low risk	No additional biases were identified

Baumgartner 2003

Methods	<ol style="list-style-type: none"> 1. Design: Parallel, 2 arms 2. Country: Switzerland 3. Multisite: No 4. International: No 5. Treatment duration: 7 days 6. Follow-up: 6 hours 7. Rate of ascent: unclear 8. Final altitude reached: 4559 metres (hypobaric chamber) 9. AMS scale: Environmental Symptom Questionnaire of Sampson 10. Randomization unit: participant 11. Analysis unit: group
Participants	<p>20 participants enrolled (Healthy white men living at altitudes below 500 metres. Men with a history of migraine or other headaches were not included)</p> <p>Randomized to:</p> <p>Flunarizine (10, 50%)</p> <p>Placebo (10, 50%)</p> <p>No participants lost to follow-up:</p> <p>Main characteristics of participants:</p> <p>Mean age (SD; range) = 24 (4, 20 to 35)</p>
Interventions	<ol style="list-style-type: none"> 1. Flunarizine group (intervention): 2 tablets of flunarizine 5 mg daily for 7 days 2. Placebo group (control): 2 tablets of placebo 5 mg daily for 7 days, identical form, colour and weight as intervention <p>Cointerventions: Not stated</p>
Outcomes	<p>This trial did not specify by primary or secondary outcomes</p> <ol style="list-style-type: none"> 1. Assessment of HAH and symptoms of AMS 2. Static posturography 3. Memory test. 4. BP and SaO₂ measurements

Notes	1. Trial Registration: Not stated 2. Funder: Not stated 3. Role of funder: Not stated 4. A priori sample size estimation: No 5. Conducted: Not stated 6. Declared conflicts of interest: Not reported	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Insufficient information to score this item as low or high risk of bias
Allocation concealment (selection bias)	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Quote: "The trial medication consisted of two tablets containing 5 mg of flunarizine or two tablets of identical form, colour and weight containing placebo" (Page 334) Insufficient information to score this item as low or high risk of bias
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Incomplete outcome data (attrition bias) All outcomes	Low risk	No participants were lost to follow-up
Selective reporting (reporting bias)	High risk	Participant-important outcomes, such as adverse events, were not reported
Other bias	Low risk	No additional biases were identified

Bernhard 1994

Methods	<ol style="list-style-type: none"> 1. Design: Paralell, 2 arms 2. Country: Bolivia 3. Multisite: No 4. International: No 5. Treatment duration: 4 days 6. Follow-up: Unclear 7. Rate of ascent: unclear 8. Final altitude reached: 5334 metres 9. AMS scale: Modified Enviromental Symptom Questionnaire 10. Randomization unit: participant 11. Analysis unit: group 	
Participants	<p>23 participants enrolled (healthy lowland-living volunteers interested in high-altitude research)</p> <p>Exclusion criteria: People who had been to high altitude 4 weeks prior to study; prior history of any chronic medical conditions including peptic ulcer disease, psychiatric illness or sensitivity to dexamethasone</p> <p>Randomized to:</p> <p>Dexamethasone (n = 11, 48%)</p> <p>Placebo (n = 12, 52%)</p> <p>No participants lost to follow-up</p> <p>Main characteristics of participants:</p> <p>Age (mean, SEM): dexamethasone 43 ± 3.9 years. placebo 32 ± 1.6 years</p> <p>Number of women/men: 15 women / 8 men</p> <p>History of AMS: 40% experienced mild to moderate AMS at altitudes less than 4000 metres</p>	
Interventions	<ol style="list-style-type: none"> 1. Dexamethasone group (intervention): dexamethasone capsules 4 mg every 12 hours orally for 4 days 2. Placebo group (control): placebo capsules 4 mg every 12 hours orally for 4 days (identical capsules to intervention) 3. Cointervention: None declared 	
Outcomes	<p>This trial did not specify by primary or secondary outcomes</p> <ol style="list-style-type: none"> 1. AMS definition 1: Presence of at least 3 cerebral symptoms with a minimum of 1 symptom at intensity score > 2 2. AMS: definition 2: Scores > 0.7 for AMS-C and 0.6 for AMS-R 3. SaO₂ and heart rate 4. Side effects 	
Notes	<ol style="list-style-type: none"> 1. Trial Registration: Not stated 2. Funder: Dr Clark Watts, Organon Inc and Nellcor Inc 3. Role of funder: Not stated; pharmaceutical supplies 4. A priori sample size estimation: No 5. Conducted: Not stated 6. Declared conflicts of interest: No 	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement

Bernhard 1994 (Continued)

Random sequence generation (selection bias)	Unclear risk	Insufficient information to score this item as low or high risk of bias
Allocation concealment (selection bias)	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Quote: "Subjects were randomly assigned to receive identical capsules of either dexamethasone 4 mg or placebo..." (Page 333) Insufficient information to score this item as low or high risk of bias
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Incomplete outcome data (attrition bias) All outcomes	Low risk	No participants were lost to follow-up
Selective reporting (reporting bias)	High risk	Participant-important outcomes, such as adverse events, were not reported
Other bias	Unclear risk	Possible industry bias: "The authors gratefully acknowledge...Organon Inc, for pharmaceutical supplies" (Page 338)

Bernhard 1998

Methods	<ol style="list-style-type: none"> 1. Design: Paralell, 2 arms 2. Country: Bolivia 3. Multisite: No 4. International: No 5. Treatment duration: 4 days 6. Follow-up: Unclear 7. Rate of ascent: unclear 8. Final altitude reached: 5334 metres 9. AMS scale: Modified Environmental Symptom Questionnaire. 10. Randomization unit: participant 11. Analysis unit: group
Participants	<p>13 participants enrolled (healthy volunteers, none of them normally resident at altitudes above 2000 metres, none of them had been to high altitude during the 4 weeks prior to the ascent)</p> <p>Randomized to:</p> <p>Dexamethasone + acetazolamide group (n = 6, 47%)</p> <p>Placebo + acetazolamide group (n = 7, 53%)</p> <p>No participant was lost to follow-up</p>

Bernhard 1998 (Continued)

	<p>Main characteristics of participants: Age (mean, SE): dexamethasone + acetazolamide 42 ± 4.7 years. placebo + acetazolamide 44 ± 3.1 years Number of women/men: 9 men + 4 women: dexamethasone + acetazolamide 4 men + 2 women; placebo + acetazolamide 5 men + 2 women 50% of participants had experienced mild to moderate AMS</p>	
Interventions	<p>1. Dexamethasone group (intervention): dexamethasone capsules 4 mg twice a day and sustained 500 mg acetazolamide given once daily for 4 days 2. Placebo group (control): placebo (identical capsules to dexamethasone) and sustained 500 mg acetazolamide capsules given once daily for 4 days 3. Cointerventions: Not stated</p>	
Outcomes	<p>This trial did not specify by primary or secondary outcomes 1. AMS definition 1: Presence of at least 3 cerebral symptoms with a minimum of 1 symptom having an intensity score of > 2 2. AMS definition 2: Scores > 0.7 for AMS-C and 0.6 for AMS-R 3. SaO₂ and heart rate</p>	
Notes	<p>1. Trial Registration: Not stated 2. Funder: Not stated 3. Role of funder: Not stated 4. A priori sample size estimation: No 5. Conducted: Not stated 6. Declared conflicts of interest: No</p>	
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "Using a randomized number system, subjects were assigned to two groups..." (Page 884)
Allocation concealment (selection bias)	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Quote: "Both researchers and subjects were blinded as to type of medication given" (Page 884) "... and a placebo in capsules identical to those used for the dexamethasone" (Page 884)
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias

Bernhard 1998 (Continued)

Incomplete outcome data (attrition bias) All outcomes	Low risk	No participants were lost to follow-up
Selective reporting (reporting bias)	High risk	Participant-important outcomes, such as adverse events, were not reported
Other bias	Low risk	No additional biases were identified

Bradwell 1986

Methods	<ol style="list-style-type: none"> 1. Design: Paralell, 2 arms 2. Country: Nepal 3. Multisite: No 4. International: No 5. Treatment duration: Unclear 6. Follow-up: Unclear 7. Rate of ascent: 5 - 10 miles per day 8. Final altitude reached: 4846 metres 9. AMS scale: Unclear (usual clinical criteria) 10. Randomization unit: participant 11. Analysis unit: group
Participants	<p>21 participants enrolled (all normally resident at < 200 metres, none was acclimatized to high altitude, and all were in good general health)</p> <p>Randomized, after being stratified by age and sex, to:</p> <p>Acetazolamide group (n = 11, 52.3%)</p> <p>Placebogroup (n = 10, 47.6%)</p> <p>Unclear number of participants were excluded: "These people and others who missed a test were excluded from the relevant analyses" (Page 1002)</p> <p>Unclear if participants were lost to follow-up</p> <p>Main characteristics of participants:</p> <p>Age (Range): 22 - 56 years</p> <p>Number of women/men: 19 men and 2 women</p> <p>12 had been on previous expedition</p>
Interventions	<ol style="list-style-type: none"> 1. Acetazolamide group (intervention): Release capsules 500 mg. No further details were provided 2. Placebo group (control): No details were provided 3. Cointerventions: Not stated
Outcomes	<p>This trial did not specify by primary or secondary outcomes</p> <ol style="list-style-type: none"> 1. Exercise performance tests 2. Tissue measurements 3. Blood gas measurements 4. AMS scores (unclear information)

Bradwell 1986 (Continued)

Notes	<p>1. Trial Registration: Not stated 2. Funder: Wellcome Trust, Wyeth Laboratories, the Arthur Thompson Trust, Birmingham Regional Health Authority, the Samuel Scott Trust, The Royal Society, the Physiological society, Squibb Medical supplies, Lederle Laboratories, among others 3. Role of funder: Not stated 5. A priori sample size estimation: No 6. Conducted: Not stated 7. Declared conflicts of interest: No</p>	
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: " At Kathmandu, subjects were randomized to acetazolamide or placebo by an independent observer after he had stratified them by age and sex" (Page 1002) Insufficient information to score this item as low or high risk of bias
Allocation concealment (selection bias)	Unclear risk	Quote: " At Kathmandu, subjects were randomized to acetazolamide or placebo by an independent observer after he had stratified them by age and sex" (Page 1002) Insufficient information to score this item as low or high risk of bias
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Quote: " At Kathmandu, subjects were randomized to acetazolamide or placebo by an independent observer after he had stratified them by age and sex" (Page 1002) "Details of medication were concealed from all subjects until treatment was withdrawn" (Page 1002)
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Unclear if participants were lost to follow-up
Selective reporting (reporting bias)	Low risk	Reporting bias was not detected
Other bias	Unclear risk	Possible industry bias: "We thank ... Wyeth laboratories... Squibb medical supplies, Lederle Laboratories, and many other societies and companies for grants " (Page

Burki 1992

Methods	<ol style="list-style-type: none"> 1. Design: Parallel design, 2 arms 2. Country: Pakistan (Karakorum Mountain) 3. Multisite: No 4. International: No 5. Treatment duration: 1 day 6. Follow-up: 4 days 7. Rate of ascent: 491.5 metres/hour 8. Final altitude reached: 4450 metres 9. AMS scale: Clinical observation: Evaluation of dizziness, nausea/vomiting and headache on a scale of 0 to 2 10. Randomization unit: participant 11. Analysis unit: participant
Participants	<p>12 healthy men signed informed consent</p> <p>Randomized to:</p> <p>Acetazolamide: 6 (50%)</p> <p>Ascorbic acid: 6 (50%)</p> <p>1 person in placebo group was excluded due to severe mountain sickness</p> <p>No losses to follow-up reported</p> <p>Main characteristics of participants</p> <p>Age: Acetazolamide: 20.2 ± 1.5, placebo: 20.7 ± 1.4</p> <p>History of AMS: Not stated</p>
Interventions	<p>Acetazolamide 250 mg twice daily at sea level (518 metres)</p> <p>Visually identical ascorbic acid 500 mg twice daily at sea level</p>
Outcomes	<p>Main outcomes were ventilatory response measured at sea level before and after taking the allocated drug, then another 2 measures were taken at 32 and 56 hours later at 4450 metres</p>
Notes	<ol style="list-style-type: none"> 1. Trial Registration: Not reported 2. Funder: Not stated 3. A priori sample size estimation: Not stated 4. Conducted: Unclear 5. Declared conflicts of interest: Not stated

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "The subjects were randomly divided into two groups..." (Page 736) Insufficient information to score this item as low or high risk of bias

Burki 1992 (Continued)

Allocation concealment (selection bias)	Unclear risk	Quote: “The subjects were randomly divided into two groups...” (Page 736) Insufficient information to score this item as low or high risk of bias
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Quote: “The subjects were given either the placebo tablets... or acetazolamide tablets... in a double-blind fashion...” (Page 736) Insufficient information to score this item as low or high risk of bias
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Incomplete outcome data (attrition bias) All outcomes	Low risk	No participants were lost to follow-up
Selective reporting (reporting bias)	High risk	Participant-important outcomes, such as adverse events, were not reported
Other bias	Low risk	No additional biases were identified

Burtscher 1998

Methods	<ol style="list-style-type: none"> 1. Design: Parallel, 2 arms 2. Country: Austria 3. Multisite: No 4. International: No 5. Treatment duration: 1 hour 6. Follow-up: 24 hours 7. Rate of ascent: Unclear 8. Final altitude reached: 3480 metres 9. AMS scale: Headache scoring
Participants	<p>29 participants enrolled (with a history of headache)</p> <p>Randomized to:</p> <p>Aspirin group (n = 15)</p> <p>Placebo group (n = 14)</p> <p>Main characteristics of participants:</p> <p>Age (mean): aspirin group 38 ± 14 years, placebo group 38 ± 14 years</p> <p>Men: aspirin group n = 9/15, placebo group n = 8/14</p> <p>History of Headache: All</p>
Interventions	<ol style="list-style-type: none"> 1. Aspirin group (intervention): aspirin 320 mg, 3 tablets at 4-hour intervals, beginning 1 hour before arrival at high altitude 2. Placebo group (control): 3 tablets at 4-hour intervals, beginning 1 hour before arrival

Burtscher 1998 (Continued)

	at high altitude	
Outcomes	1. Primary outcome Incidence and severity of headache 2. Secondary outcome Heart rate Blood pressure Arterial oxygen saturation	
Notes	1. Trial Registration: Not stated 2. Funder: Austrian Society for Mountain Medicine, the Health Section of the Austrian Alpine Club, and HoffmannLa Roche 3. Role of funder: Not stated 4. A priori sample size estimation: No 5. Conducted: Not stated 6. Declared conflicts of interest: Yes	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "Twenty nine volunteers with a history of headache at high altitude were randomly assigned in a double blind fashion to receive placebo (...)" (Page 1057) Insufficient information to score this item as low or high risk of bias
Allocation concealment (selection bias)	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Incomplete outcome data (attrition bias) All outcomes	Low risk	No participants were lost to follow-up
Selective reporting (reporting bias)	High risk	Participant-important outcomes, such as adverse events, were not reported
Other bias	Unclear risk	Unclear role of funder in this trial

Burtscher 2001

Methods	<ol style="list-style-type: none"> 1. Design: Parallel, 2 arms 2. Country: Austria 3. Multisite: No 4. International: No 5. Treatment duration: 12 hours 6. Follow-up: 2 days 7. Rate of ascent: Not clear 8. Final altitude reached: 3480 metres 9. AMS scale: Headache scoring
Participants	<p>31 participants enrolled (healthy men and women whose medical history contained reports of at least one episode of headache after ascent to altitudes above 2000 metres)</p> <p>Randomized to:</p> <p>Aspirin group (n = 16)</p> <p>Placebo group (n = 15)</p> <p>Main characteristics of participants:</p> <p>Age (median): aspirin group 39 ± 22 to 58 ; placebo group 40 ± 23 - 59</p> <p>Men: aspirin group n = 12/16, placebo group n = 8/15</p> <p>History of AMS: None</p> <p>Type of HAI reported: None</p>
Interventions	<ol style="list-style-type: none"> 1. Aspirin group (intervention): aspirin 320 mg with 150 ml water, 3 times at 4-hour intervals, beginning 2 hours before arrival at high altitude 2. Placebo group (control): tablets with 150 ml water, 3 times at 4-hour intervals, beginning 2 hours before arrival at high altitude
Outcomes	<ol style="list-style-type: none"> 1. Primary outcome: Incidence of headache 2. Secondary outcome: Arterial oxygen saturation
Notes	<ol style="list-style-type: none"> 1. Trial Registration: Not stated 2. Funder: This study was supported by the Austrian Science Fund grant P13009-MED, Grunenthal GMBH and Hoffmann-La Roche 3. Role of funder: Not stated 4. A priori sample size estimation: No 5. Conducted: Not stated 6. Declared conflicts of interest: No

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "Thirty-one subjects were randomly assigned in a double-blind fashion to receive placebo (...)" (Page 543) Insufficient information to score this item as low or high risk of bias
Allocation concealment (selection bias)	Unclear risk	Insufficient information to score this item as low or high risk of bias

Burtscher 2001 (Continued)

Blinding of participants and personnel (performance bias) All outcomes	Low risk	Quote: “Tablets (placebo or 320 mg aspirin) were administered three times at 4-hour intervals, beginning 2 hours before arrival at high altitude. Placebos were nearly identical to aspirin in appearance and taste. Tablets were administered by a person who was not involved in scoring or testing procedures” (Page 5430)
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk
Incomplete outcome data (attrition bias) All outcomes	Low risk	No participants were lost to follow-up
Selective reporting (reporting bias)	Low risk	Selective reporting of informations was not detected
Other bias	Unclear risk	Unclear role of funder in this trial

Burtscher 2014

Methods	<ol style="list-style-type: none"> 1. Design: Parallel (two arms) 2. Country: Italy 3. Multisite: No 4. International: No 5. Treatment duration: Pills were taken 10 and 1 hour before high altitude exposure 6. Follow-up: unclear 7. Rate of ascent: car travel from 600 to 3480 metres. Second and third day climbed 106 metres/hour 8. Final altitude reached: 3800 metres 9. AMS scale: Lake Louise Consensus scoring system 10. Randomization unit: patient 11. Analysis unit: group
Participants	<p>15 volunteers enrolled, all of them had history of AMS</p> <p>Exclusion criteria: Any type of acute or chronic illness; regular smoking (> 5 cigarettes per day); regular medications; stops at an altitude > 2500 metres during the previous 4 weeks; Age < 20 or > 60 years; pregnancy or lactation; haemoglobin concentration < 12.0 g/dL</p> <p>Randomized to:</p> <p>Acetazolamide group (n = 7, 46.6%)</p> <p>Placebo group (n = 8, 53.4%)</p> <p>No participants randomized were excluded</p> <p>No participants lost to follow-up</p> <p>Main characteristics of participants:</p> <p>Age (median/mean, SD): 43.6 ± 13.4, placebo 44.7 ± 8.6</p>

	Number of men/woman: Acetazolamide: 4 men: 3 women, placebo: 4 men: 4 women	
Interventions	Acetazolamide group (intervention): received 2 tablets (2 × 125 mg acetazolamide) to be taken 10 hours and 1 hour before high altitude exposure Placebo group received placebo the same way	
Outcomes	This trial did not specify by primary or secondary outcomes 1. AMS symptoms according to the Lake Louise Score. Participants were considered to be suffering from AMS when the score was ≥ 3 2. Physiological variables: heart rate, minute ventilation, arterial blood gases analysis	
Notes	1. Trial Registration: Not stated 2. Funder: Not specified 3. Role of funder: Not specified 4. A priori sample size estimation: No 5. Conducted: Not stated 6. Declared conflicts of interest: yes	
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote " Study participants were randomly assigned in a double blind fashion to receive placebo or acetazolamide before exposure to high altitude" (Page 4379)
Allocation concealment (selection bias)	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Quote: "Subjects received two tablets (2 × 125 mg acetazolamide or placebo) to be taken 10 hours and 1 hour before arrival at high altitude. Tablets were administered by a person who was not involved in evaluations and the timely intake of tablets has been checked" (Page 4379)
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Incomplete outcome data (attrition bias) All outcomes	Low risk	No participants were lost to follow-up
Selective reporting (reporting bias)	High risk	Participant-important outcomes, such as adverse events, were not reported

Burtscher 2014 (Continued)

Other bias	Low risk	No additional biases were identified
------------	----------	--------------------------------------

Carlsten 2004

Methods	<ol style="list-style-type: none"> 1. Design: Parallel design, 3 arms 2. Country: Bolivia 3. Multisite: No 4. International: No 5. Treatment duration: 24 hours 6. Follow-up: 24 hours 7. Rate of ascent: Unknown 8. Final altitude reached: 3630 metres 9. AMS scale: LLS 10. Randomization unit: participant 11. Analysis unit: participant
Participants	<p>32 healthy vacationers who had flown from Miami to Bolivia</p> <p>Randomized to:</p> <p>Acetazolamide 250 mg: 11 (34.3%)</p> <p>Acetazolamide 500 mg: 11 (34.3%)</p> <p>Placebo (ascorbic acid): 10 (31.2%)</p> <p>No losses to follow-up reported</p> <p>Main characteristics of participants: Not stated</p>
Interventions	Acetazolamide 250 mg, acetazolamide 125 mg, ascorbic acid every 8 hours, 2 doses
Outcomes	<p>AMS score</p> <p>Absolute change from evaluation at 0 hours to evaluation at 24 hours</p>
Notes	<ol style="list-style-type: none"> 1. Trial Registration: Not reported 2. Funder: Houston Award Fund, Emge Travelling Scholars programme at Stanford University School of Medicine and the Center for Latin American Studies at Stanford University 3. Role of funder: Financial support 4. A priori sample size estimation: Not stated 5. Conducted: Unclear 6. Declared conflicts of interest: Not stated

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "33 subjects were randomly given one of three identical packets, each packet containing two tablets..." (Page 35) Insufficient information to score this item as low or high risk of bias

Carlsten 2004 (Continued)

Allocation concealment (selection bias)	Unclear risk	Quote: “33 subjects were randomly given one of three identical packets, each packet containing two tablets...” (Page 35) Insufficient information to score this item as low or high risk of bias
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Incomplete outcome data (attrition bias) All outcomes	Low risk	3% of participants were lost to follow-up
Selective reporting (reporting bias)	High risk	Participant-important outcomes, such as adverse events, were not reported
Other bias	Low risk	No additional biases were identified

Chen 2015

Methods	<ol style="list-style-type: none"> 1. Design: Parallel, 4 arms 2. Country: China 3. Multisite: No 4. International: No 5. Treatment duration: 3 days 6. Follow-up: unclear 7. Rate of ascent: flight from 500 metres to 3700 metres in 2½ hours 8. Final altitude reached: 3700 metres 9. AMS scale: Lake Louise Score 10. Randomization unit: participant 11. Analysis unit: group
Participants	<p>80 healthy young men, lowland residents of Chengdu, China</p> <p>Inclusion criteria: Resident at or below 500 metres; Healthy; 18 - 35 years of age</p> <p>Exclusion criteria: HAI (> 2500 metres) exposure history in the past year; organic diseases such as congenital heart disease, dysrhythmia, liver or kidney dysfunction, or psychological or neurological disorders</p> <p>Randomized to:</p> <p>Budesonida inhaled group (n = 20, 25%)</p> <p>Procaterol tablet group (n = 20, 25%)</p> <p>Budesonida/formoterol inhaled group (n = 20, 25%)</p> <p>Placebo Group (n = 20, 25%)</p> <p>No participants randomized were excluded</p> <p>No participants lost to follow-up</p>

	Main characteristics of participants: Age (median/mean \pm SD): budesonide 21.85 \pm 3.23 procaterol 20.3 \pm 2.03 budesonide/ formoterol 20.6 \pm 2.76 placebo 21.65 \pm 3.31 Number of men/women: Not specified	
Interventions	1. Group A received Budesonide 200 mg twice daily 2. Group B received procaterol 25 mg twice daily 3. Group C received Formoterol/budesonide 160 mg 4.5 mg twice daily 4. Group D received placebo tablets, one tablet twice daily	
Outcomes	Primary outcomes: 1. Symptoms of AMS at 20, 72, and 120 hours after arrival at 3700 metres altitude Secondary outcomes 1. HAPE or HACE Other outcomes: 1. Adverse reactions 2. Heart rate and SpO ₂ 3. Pulmonary function test	
Notes	1. Trial Registration: ChiCTRPRC-12002748 2. Funder: Special Health Research Project, Ministry of Health of P.R. China (grant No. 201002012) 3. Role of Funder: None 4. A priori sample size estimation: No 5. Conducted: between June 4 and June 16, 2012 6. Declared conflicts of interest: no	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote " Subjects were randomly assigned to four groups (n = 20), by a physician who did not participate in later parts of the study, using a computer-generated random number List" (Page 198)
Allocation concealment (selection bias)	Low risk	Quote: "The physician who made group assignments prepared one medicine box for each subject. The physician then gave these boxes to other researchers and kept the blinding code. The subjects were fully informed and knew that they could be assigned to any of four groups and that one group would take a placebo." (Page 198)
Blinding of participants and personnel (performance bias) All outcomes	High risk	Quote: "We initially intended to design a double-blind trial. However, the procaterol tablet and placebo groups used oral tablets,

Chen 2015 (Continued)

		and the budesonide and budesonide/formoterol groups used inhalants. So subjects might assume that they were given a different drug than those in another group, although they could not know specifically what drug they were taking” (Page 204)
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Incomplete outcome data (attrition bias) All outcomes	Low risk	There were no losses to follow-up
Selective reporting (reporting bias)	High risk	There was no information on HAPE and HACE
Other bias	Low risk	No additional biases were identified

Chow 2005

Methods	<ol style="list-style-type: none"> 1. Design: Parallel design, 3 arms 2. Country: USA 3. Multisite: No 4. International: No 5. Treatment duration: 5 days 6. Follow-up: 1 day 7. Rate of ascent: 1285 metres/hour 8. Final altitude reached: 3800 metres 9. AMS scale: The Lake Louise acute mountain sickness scoring system
Participants	<p>68 enrolled and randomized</p> <p>Exclusion criteria: travelled to an elevation above 2400 metres within 30 days of the study; contraindications to high altitude exposure; pregnant; pre-existing use of acetazolamide or ginkgo biloba; known hypersensitivity of acetazolamide or ginkgo biloba; known bleeding disorders or receiving anticoagulant therapy; scheduled a surgical or dental procedure within 14 days of study participations</p> <p>Randomized to:</p> <p>Acetazolamide: 24/68 (35.3%) 3 withdrew before ascent</p> <p>Ginkgo biloba: 21/68 (30.9%) 4 withdrew before ascent</p> <p>Placebo: 23/68 (33.8%) 3 withdrew before ascent. 1 person in the acetazolamide group withdrew after ascent for personal reasons</p> <p>Main characteristics of participants</p> <p>Age: Acetazolamide: 32 (25 - 42); Ginkgo biloba: 40 (25 - 62); Placebo: 33.5 (24 - 65)</p> <p>No. of men: Acetazolamide: 13 (65%); Ginkgo biloba: 10 (58.8%); Placebo: 10 (50%)</p> <p>History of AMS: Not stated</p>

Chow 2005 (Continued)

Interventions	1. Acetazolamide 250 mg twice a day 2. Gingko Biloba 120 mg twice a day 3. Control: placebo twice a day
Outcomes	1. Primary: LLS self-report questionnaire score and the incidence of AMS 2. Secondary: Number of participants requesting analgesics Number of participants requesting anti-emetics Number of participants experiencing high-altitude pulmonary oedema or high-altitude cerebral oedema Incidence of other symptoms
Notes	1. Trial Registration: Not reported 2. Funder: Not stated 3. Role of funder: Not stated 4. A priori sample size estimation: Yes, page 298 5. Conducted: Not stated 6. Declared conflicts of interest: Not stated

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "We developed a randomization sequence by drawing cards out of a hat, using 25 labeled cards for each group" (Page 297)
Allocation concealment (selection bias)	Low risk	Quote: "Study medications were prepared (...) with enclosed administration instructions and affixed with serial numbers" (Page 297)
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Quote: "To maintain blinding, subjects in acetazolamide group started taking placebo 5 days before ascent and switched to a typical dose for AMS prophylaxis" (Page 297)
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: "in the event of an emergency, an investigator had access to the study key, which was stored within a sealed envelope" (Page 297)
Incomplete outcome data (attrition bias) All outcomes	Low risk	Percentage of participants lost at follow-up: 16.1%
Selective reporting (reporting bias)	Low risk	Reporting bias was not detected

Chow 2005 (Continued)

Other bias	Low risk	No other biases were identified
------------	----------	---------------------------------

Ellsworth 1991

Methods	<ol style="list-style-type: none"> 1. Design: randomized, double-blind, concurrent, placebo-controlled (declared as cross-over by authors) 2. Country: Mount Rainer, Seattle Washington (USA) 3. Multisite: No 4. International: No 5. Treatment duration: 2 days 6. Follow-up: unclear 7. Rate of ascent: 1800 metres/7 hours, day 2 1392 metres/7 hours 8. Final altitude reached: 4392 metres 9. AMS scale: Environmental Symptoms Questionnaire, second revision (ESQ-III)
Participants	<p>18 participants were enrolled. They normally resided at sea level and had not been exposed to high altitude within 3 weeks before the study. All were free of cardiorespiratory disease, and none had a history of diabetes mellitus, sulfa drug allergy, acid peptic disease, or psychiatric illness</p> <p>Randomized to:</p> <p>Acetazolamide 8 (44%) Dexamethasone 10 (56%)</p> <p>Authors did not report exclusions and losses during trial</p> <p>Main characteristics of participants:</p> <p>Age: Acetazolamide 32.6 ± 3.9; dexamethasone 36.2 ± 2.4 Percentage/number of women/men: Acetazolamide: 6 men (75%), 2 women (25%); dexamethasone: 5 men (50%) 5 women (50%) Percentage/number History of AMS: Acetazolamide: 5 (62%); dexamethasone: 3 (30%)</p>
Interventions	<ol style="list-style-type: none"> 1. Acetazolamide, 250 mg (750 in 24 hours) 2. Dexamethasone, 4 mg (12 mg in 24 hours) 3. Lactose placebo
Outcomes	<p>Outcomes were not classified as primary or secondary</p> <p>Incidence of AMs (unclear data)</p> <p>AMS-C scores</p> <p>AMS-R scores</p>
Notes	<ol style="list-style-type: none"> 1. Trial Registration: Not stated 2. Funder: Not stated 3. Role of funder: Not stated 4. A priori sample size estimation: No 5. Conducted: Unclear 6. Declared conflicts of interest: Not reported

Risk of bias

Bias	Authors' judgement	Support for judgement
------	--------------------	-----------------------

Ellsworth 1991 (Continued)

Random sequence generation (selection bias)	Low risk	Quote: “Using a random numbers table (. . .)” (Page 289)
Allocation concealment (selection bias)	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Quote: “The drugs were packaged in identical appearing pink capsules by Pharmaceutical services” (Page 289)
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Quote: “(...) a clinical interview and examination were conducted by one of the investigators (AJE) without knowledge of the subjects response to the questionnaire” (Page 290)
Incomplete outcome data (attrition bias) All outcomes	Low risk	No participants were lost to follow-up
Selective reporting (reporting bias)	High risk	Participant-important outcomes, such as adverse events, were not reported. Report of incidence of AMS is unclear
Other bias	Low risk	No other biases were identified

Faull 2015

Methods	<ol style="list-style-type: none"> 1. Design: Parallel, 2 arms 2. Country: Italy 3. Multisite: No 4. International: No 5. Treatment duration: 4 days 6. Follow-up: 2 days 7. Rate of ascent: Unclear 8. Final altitude reached: 3459 metres 9. AMS scale: Lake Louise Score
Participants	<p>20 participants enrolled (healthy men and women residing at elevations between 50 metres and 150 metres, without recent (within 2 months) exposure to high altitudes)</p> <p>Randomized to:</p> <p>Acetazolamide group (n = 10)</p> <p>Placebo Group (n = 10)</p> <p>Main characteristics of participants:</p> <p>Age (median): total group: 43 ± 16</p> <p>Men: acetazolamide group: n = 7; placebo group: n = 7</p> <p>History of AMS: None</p> <p>Type of HAI reported: None</p>

Interventions	<ol style="list-style-type: none"> 1. Acetazolamide group (intervention): acetazolamide 250 mg taken every 12 hours starting 3 days before ascent 2. Placebo group (control): tablets 250 mg taken every 12 hours starting 3 days before ascent
Outcomes	<ol style="list-style-type: none"> 1. Primary outcome: Prosaccadic and antisaccadic eye movements 2. Secondary outcome: Presence of AMS
Notes	<ol style="list-style-type: none"> 1. Trial Registration: Not stated 2. Funder: Jabbs Foundation 3. Role of funder: Not stated 4. A priori sample size estimation: No 5. Conducted: Not stated 6. Declared conflicts of interest: No

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "Minimization was used to reduce group differences in AMS susceptibility, age, and sex. Subjects were randomly allocated to receive either 250 mg acetazolamide or identically matching placebo(...)" (Page 73)
Allocation concealment (selection bias)	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	1 participant (5%) was removed from final analysis
Selective reporting (reporting bias)	High risk	Participant-important outcomes, such as adverse events, were not reported
Other bias	Low risk	No other biases were identified

Fischer 2000a

Methods	<ol style="list-style-type: none"> 1. Design: Parallel randomized design 2. Country: Switzerland 3. Multisite: No 4. International: No 5. Treatment duration: 3 hours 6. Follow-up: unclear 7. Rate of ascent: unclear 8. Final altitude reached: 3454 metres 9. AMS scale: Acute Mountain Sickness Score 	
Participants	<p>21 participants enrolled (healthy mountaineers with normal weight and constant good health)</p> <p>Exclusion criteria: women, smoking, non-compliance with study protocol, previous pulmonary disease</p> <p>Randomized to:</p> <p>Theophylline group (unclear)</p> <p>Placebo group (unclear)</p> <p>No participants were lost to follow-up</p> <p>Main characteristics of participants:</p> <p>Age (median/mean): 29 ± 8</p> <p>Percentage of men: 100%</p>	
Interventions	<ol style="list-style-type: none"> 1. Theophylline (intervention): 375 mg slow-release tablets taken twice daily for 3 days, or 250 mg twice daily for participants < 70 kg. This was stopped 12 hours after arrived at altitude 2. Placebo group (control): placebo tablets twice daily for 3 days 	
Outcomes	<p>Outcomes were not classified as primary or secondary</p> <ol style="list-style-type: none"> 1. AMS scores by LLS 2. Measurements of respiratory frequency 3. Pulse rate 4. Oxygen saturation 5. Serum theophylline level 	
Notes	<ol style="list-style-type: none"> 1. Trial Registration: Not stated 2. Funder: Byk Gulden, Constance, Germany 3. Role of funder: Not stated 4. A priori sample size estimation: No 5. Conducted: Not stated 6. Declared conflicts of interest: Not reported 	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "21 subjects were randomly allocated to placebo (...)" (Page 124)

Fischer 2000a (Continued)

Allocation concealment (selection bias)	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	No participants were lost to follow-up
Selective reporting (reporting bias)	Low risk	Reporting bias was not detected
Other bias	Unclear risk	Unclear role of funder

Fischer 2000b

Methods	<ol style="list-style-type: none"> 1. Design: Cross-over design 2. Country: Germany 3. Multisite: No 4. International: No 5. Treatment duration: first phase 3 days 6. Follow-up: unclear 7. Rate of ascent: unclear 8. Final altitude reached: first phase 4500 metres 9. AMS scale: Acute Mountain Sickness Score
Participants	<p>14 participants enrolled (healthy mountaineers with normal weight and constant good health)</p> <p>Exclusion criteria: women, smoking, non-compliance with study protocol, previous pulmonary disease</p> <p>Randomized to:</p> <p>Theophylline group (unclear)</p> <p>Placebo group (unclear)</p> <p>No participants were lost to follow-up</p> <p>Main characteristics of participants:</p> <p>Age (median/mean): 29 ± 8 first study</p> <p>Percentage of men: 100%</p>
Interventions	<ol style="list-style-type: none"> 1. Theophylline (intervention): 375 mg slow-release tablets taken twice daily for 3 days, or 250 mg twice daily for participants < 70 kg 2. Placebo group (control): placebo tablets twice daily for 3 days
Outcomes	<p>Outcomes were not classified as primary or secondary</p> <ol style="list-style-type: none"> 1. AMS scores by LLS

Fischer 2000b (Continued)

	2. Measurements of respiratory frequency 3. Pulse rate 4. Oxygen saturation 5. Serum theophylline level	
Notes	1. Trial Registration: Not stated 2. Funder: Byk Gulden, Constance , Germany 3. Role of funder: Not stated 4. A priori sample size estimation: No 5. Conducted: Not stated 6. Declared conflicts of interest: Not reported	
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "14 subjects were randomly allocated to placebo or study medication for the first session(...)" (Page 124)
Allocation concealment (selection bias)	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Quote: "or matched placebo tablets twice daily" (Page 124) Insufficient information to score this item as low or high risk of bias
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Incomplete outcome data (attrition bias) All outcomes	Low risk	No participants were lost to follow-up
Selective reporting (reporting bias)	High risk	Participant-important outcomes, such as adverse events, were not reported
Other bias	Unclear risk	Unclear role of funder. It is unclear if previous events of HAI (specifically in phase 1) affected the probability of new events in second phase of cross-over trials

Fischer 2004

Methods	<ol style="list-style-type: none"> 1. Design: cross-over trial 2. Country: Germany 3. Multicentre study: No 4. Altitude setting: 4500 metres 5. AMS scale: The Lake Louise self-assessment questionnaire (LLS) and the ESQ were used to assess symptoms of AMS at 0, 3, 6 and 9 hours
Participants	<p>10 participants enrolled (male volunteers) Exclusion criteria: Not provided Randomized to each group with an interval of 2 weeks between each of the three chamber sessions Theophylline group Acetazolamide group Placebo group No participants were lost to follow-up Main characteristics of participants: Age (median/mean): 24.8 years Percentage of men: 100%</p>
Interventions	<ol style="list-style-type: none"> 1. Intervention: acetazolamide (250mg twice a day) 2. Theophylline (250 mg twice a day) 3. Placebo (twice a day)
Outcomes	<p>Outcomes were not classified as primary or secondary</p> <ol style="list-style-type: none"> 1. LLS scores 2. ESQ scores 3. PaO₂, PaCO₂ and PH measurements 4. Magnetic resonance imaging
Notes	<ol style="list-style-type: none"> 1. Trial Registration: Not stated 2. Funder: Deutsche Akademik für Flug + Radiometer Inc 3. Role of funder: Not stated 4. A priori sample size estimation: No 5. Conducted: Not stated 6. Declared conflicts of interest: Not reported

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Insufficient information to score this item as low or high risk of bias
Allocation concealment (selection bias)	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias

Fischer 2004 (Continued)

Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Incomplete outcome data (attrition bias) All outcomes	Low risk	No participants were lost to follow-up
Selective reporting (reporting bias)	High risk	Participant-important outcomes, such as adverse events, were not reported. Numbers of participants by arm were not provided
Other bias	Unclear risk	Unclear role of funder. It is unclear if previous events of HAI (specifically in phase 1) affected the probability of new events in second phase of cross-over trials

Fulco 2006

Methods	<ol style="list-style-type: none"> 1. Design: Cross-over trial (4 arms) 2. Country: USA 3. Multisite: No 4. International: No 5. Treatment duration: 2 days 6. Follow-up: unclear 7. Rate of ascent: unclear 8. Final altitude reached: 4300 metres 9. AMS scale: ESQ
Participants	<p>6 participants enrolled. All were born at altitude < 1500 metres and resided near sea level for at least 6 months</p> <p>4-week long definitive testing phase</p> <p>Randomized each week to:</p> <ul style="list-style-type: none"> Sea level + placebo Sea level + acetazolamide Simulated altitude + placebo Simulated altitude + acetazolamide <p>No participants were lost to follow-up</p> <p>Main characteristics of participants:</p> <p>Age (median/mean): 20 ± 1 years</p> <p>Number of men: 5/6</p>
Interventions	<ol style="list-style-type: none"> 1. Acetazolamide 250 mg, 3 times a day for 2 days 2. Placebo group (control), 3 times a day for 2 days
Outcomes	<p>Outcomes were not classified as primary or secondary</p> <ol style="list-style-type: none"> 1. Physiological measurements 2. ESQ scores

Fulco 2006 (Continued)

	3. AMS-C 4. AMS-R	
Notes	1. Trial Registration: Not stated 2. Funder: Not stated 3. Role of funder: Not stated 4. A priori sample size estimation: No 5. Conducted: Not stated 6. Declared conflicts of interest: No	
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "The presentation of the definitive exercise testing bouts...was assigned randomly for each subject" (Page 684) Insufficient information to score this item as low or high risk of bias
Allocation concealment (selection bias)	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Quote: "Both the subjects and the investigators directly involved were blinded to drug treatment status..." (Page 684) "Acetazolamide and an identically appearing placebo capsule were prepared by a local pharmacy that had no other relationship with the study" (Page 685)
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: "Both the subjects and the investigators directly involved were blinded to drug treatment status..." (Page 684)
Incomplete outcome data (attrition bias) All outcomes	Low risk	No participants were lost to follow-up
Selective reporting (reporting bias)	High risk	Participant-important outcomes, such as adverse events, were not reported
Other bias	Unclear risk	It is unclear if previous events of HAI (specifically in phase 1) affected the probability of new events in second phase of cross-over trials

Greene 1981

Methods	<ol style="list-style-type: none"> 1. Design: Cross-over design (2 arms) 2. Country: Kenya 3. Multisite: No 4. International: No 5. Treatment duration: 4 weeks 6. Follow-up: 4 weeks 7. Final altitude reached: varied 8. AMS scale: Self-administered subjective questionnaire of AMS symptoms 	
Participants	<p>24 British climbers; none were professional sportsmen; 5 were medically trained</p> <p>2. Participants were paired for age, sex and likely activities, and each member of each pair was allocated at random to 1 of 2 treatment groups: Acetazolamide 500 mg sustained release nightly Placebo: identically presented</p> <p>No participants were lost to follow-up</p> <p>Main characteristics of participants: 2 women, 22 men</p> <p>History of AMS: Not reported Type of HAI reported: Not reported</p>	
Interventions	<ol style="list-style-type: none"> 1. Acetazolamide 500 mg nightly during 5 nights before and after exposure 2. Placebo in the same way 	
Outcomes	<p>Outcomes were not classified as primary or secondary</p> <ol style="list-style-type: none"> 1. Scores for AMS from symptom cards 2. Adverse events 	
Notes	<ol style="list-style-type: none"> 1. Trial Registration: Not stated 2. Funder: Young Explorers Trust, Lederle laboratories 3. Role of funder: Not stated 4. A priori sample size estimation: No 5. Conducted: Not stated 6. Declared conflicts of interest: No 	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "They were paired for age, sex and likely activities, and each member of each pair was allocated at random to one of two treatment groups" (Page 811) Insufficient information to score this item as low or high risk of bias
Allocation concealment (selection bias)	Unclear risk	Insufficient information to score this item as low or high risk of bias

Greene 1981 (Continued)

Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Incomplete outcome data (attrition bias) All outcomes	Low risk	No participants were lost to follow-up
Selective reporting (reporting bias)	Unclear risk	Most of information was presented as graphs
Other bias	Unclear risk	Unclear role of funder. It is unclear if previous events of HAI (specifically in phase 1) affected the probability of new events in second phase of cross-over trials

Hackett 1976

Methods	<ol style="list-style-type: none"> 1. Design: Parallel design (2 arms) 2. Country: Nepal 3. Multisite: No 4. International: No 5. Treatment duration: 4 days 6. Follow-up: Unclear 7. Final altitude reached: 4243 metres 8. AMS scale: Subjective symptoms evaluation
Participants	<p>278 hikers recruited in Namche Bazar (3440 metres) were included (volunteers). Number of participants assigned to each group is unclear</p> <p>Assigned to</p> <p>Acetazolamide 71 (24%)</p> <p>Placebo 49 (39%)</p> <p>No treatment controls 158 (69%): participants not taking tablets</p> <p>3. Number of participants lost to follow-up unclear. 52 questionnaires were excluded on their return</p> <p>Main characteristics of participants:</p> <p>Age: 33: 18 - 71 years</p> <p>Men: 71%</p>
Interventions	<ol style="list-style-type: none"> 1. Acetazolamide 250 mg starting at 3440 metres twice daily for 4 days 2. Placebo tablets (lactose, provided by the Royal Drug company, Kathmandu, Nepal) twice daily for 4 days

Hackett 1976 (Continued)

Outcomes	Outcomes were not classified as primary or secondary 1. Acute Mountain Sickness 2. Severity: HAPE or cerebral oedema	
Notes	1. Trial Registration: Not stated 2. Funder: Unclear 3. Role of funder: Not stated 4. A priori sample size estimation: No 5. Conducted: Oct 10 to Nov 10, 1975 6. Declared conflicts of interest: No	
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "Those volunteering were assigned to placebo or acetazolamide groups and subjects taking no tablets were classified as controls" (Page 1150) Insufficient information to score this item as low or high risk of bias
Allocation concealment (selection bias)	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Quote: "Tablets were packaged into small plastic bags (coded for later identification) each containing a course of medication and selected at random so that neither the subject nor the investigator knew which was being given" (Page 1150)
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: "Tablets were packaged into small plastic bags (coded for later identification) each containing a course of medication and selected at random so that neither the subject nor the investigator knew which was being given" (Page 1150)
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Number of participants lost from each arm to follow-up unclear
Selective reporting (reporting bias)	High risk	Participant-important outcomes, such as adverse events, were not reported. HAPE and HACE results are not clearly reported in "Results" section
Other bias	Low risk	No other biases were identified

Hackett 1988

Methods	<ol style="list-style-type: none"> 1. Design: Randomized double-blind study 2. Country: USA 3. Multisite: No 4. International: No 5. Treatment duration: 1 day 6. Follow-up: unclear 7. Rate of ascent: 4400 metres in 1 hour, by helicopter 8. Final altitude reached: 4400 metres 9. AMS scale: AMS Symptoms Questionnaire 	
Participants	<p>15 healthy military men on no medication were enrolled; None had been to high altitude within 3 weeks before the study</p> <p>Randomized to:</p> <p>Placebo (n = 7)</p> <p>Dexamethasone 2 mg (n = 8)</p> <p>No participants were lost to follow-up</p> <p>Main characteristics of participants:</p> <p>Men, age 28 ± 1.0 year, height 181 ± 2 cm, and weight 83±4 kg</p>	
Interventions	<p>Dexamethasone: 2 mg dexamethasone every 6 hours starting 1 hour before flying</p> <p>Placebo: no details were provided</p>	
Outcomes	<p>Outcomes were not classified as primary or secondary</p> <ol style="list-style-type: none"> 1. AMS scores and severity 2. AMS-C 3. AMS-R 4. Physiological measurements 	
Notes	<ol style="list-style-type: none"> 1. Trial Registration: Not stated 2. Funder: Unclear “Many people and organizations” 3. Role of funder: Unclear 4. A priori sample size estimation: No 5. Conducted: Unclear 6. Declared conflicts of interest: Not reported 	
<i>Risk of bias</i>		
Bias	Authors’ judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: “the 15 subjects were randomized to receive (...)” (Page 951) Insufficient information to score this item as low or high risk of bias
Allocation concealment (selection bias)	Unclear risk	Insufficient information to score this item as low or high risk of bias

Hackett 1988 (Continued)

Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Incomplete outcome data (attrition bias) All outcomes	Low risk	No participants were lost to follow-up
Selective reporting (reporting bias)	High risk	Participant-important outcomes, such as adverse events, were not reported
Other bias	Low risk	No other biases were identified

HEAT 2010

Methods	<ol style="list-style-type: none"> 1. Design: Randomized trial, parallel, 3 arms 2. Country: Nepal 3. Multisite: No 4. International: No 5. Treatment duration: 1 day 6. Intention-to-treat: Yes 7. Follow-up: 1 day 8. Rate of ascent: unclear 9. Final altitude reached: 4928 metres 10. AMS scale: Lake Louise AMS questionnaire (LLS)
Participants	<p>343 participants enrolled (healthy men and women, 18 - 65 years), to ascend from 2 villages at 4280 metres and 4358 metres respectively, to 4928 metres</p> <p>Exclusion criteria: headache at recruitment, diagnosis of AMS, signs or symptoms of a substantial acute infection, had slept above 4500 metres, or had taken any NSAIDs or acetazolamide within 1 day or 3 days prior to enrolment, respectively</p> <p>Randomized into 3 groups:</p> <p>Placebo (89)</p> <p>Ibuprofen (129)</p> <p>Acetazolamide (125)</p> <p>48 participants randomized were excluded due to protocol violations:</p> <p>Placebo (12, 13%)</p> <p>Ibuprofen (18, 14%)</p> <p>Acetazolamide (18, 14%)</p> <p>Participants lost to follow-up: 78 (22.7%) lost to follow-up for unclear reasons</p> <p>Main characteristics of participants:</p> <p>Age: 39.2 ± 12.1 (placebo), 39.1 ± 12 (acetazolamide), 37 ± 11.4 (ibuprofen)</p> <p>Number/Percentage of men: 47/65 (72.3% placebo), 65/97 (67.7% acetazolamide), 75/103 (73.5% ibuprofen)</p> <p>Percentage/number History of AMS: 3/65 (4.6% placebo), 2/97 (2.1% acetazolamide)</p>

	, 2/103 (1.9% ibuprofen)
Interventions	<ol style="list-style-type: none"> 1. Placebo group: placebo 3 times a day orally for 1 day prior to the ascent 2. Ibuprofen group: 600 mg of ibuprofen 3 times a day orally <p>Cointervention: In all 3 groups there was a period of acclimatization, approximately three nights in each group</p>
Outcomes	<p>Primary outcome</p> <ol style="list-style-type: none"> 1. Incidence of headache at the study endpoint as calculated on the Lake Louise AMS Questionnaire (LLS) <p>Secondary endpoints</p> <ol style="list-style-type: none"> 1. Evaluation of headache severity by visual analog scale (VAS) 2. Pulse oximetry 3. AMS incidence and severity as measured by the LLS 4. Side effects
Notes	<ol style="list-style-type: none"> 1. Trial Registration: Not stated 2. Funder: Himalayan Rescue Association, Deurali-Janta Pharmaceuticals of Kathmandu, Nepal and Hari Bhakta Sharma. Drs Derek and Lydia Lipman 3. Role of funder: randomization of the drugs and packaging. Drs Derek and Lydia Lipman (financial support) 4. A priori sample size estimation: Yes 5. Conducted: October 2005 to November 2005 6. Declared conflicts of interest: Yes. (Page 241)

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "Study medications were randomized via computer-generated code" (Page 237)
Allocation concealment (selection bias)	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Incomplete outcome data (attrition bias) All outcomes	High risk	22.7% of participants were lost to follow-up and not include in final analysis
Selective reporting (reporting bias)	Low risk	Reporting bias was not detected

Other bias	Unclear risk	Possible industry bias
------------	--------------	------------------------

Hillenbrand 2006

Methods	<ol style="list-style-type: none"> 1. Design: Randomized, double-blind controlled trial 2. Country: Mount Everest region of Nepal 3. Multisite: No 4. International: No 5. Follow-up: 7 days 6. Rate of ascent : 300 metres/day 7. Final altitude reached: 4930 metres 8. AMS scale: Lake Louise AMS symptom score
Participants	<p>403 male Nepali porters (adults) were enrolled for 8 Nepali doctors</p> <p>Exclusion criteria: AMS, various medical conditions, sulphonamide allergy or any other previous drug reactions, or taking a different route that did not pass through the assessment stations</p> <p>3 porters were excluded for 1 of these reasons</p> <p>Randomized to:</p> <p>Acetazolamide group = 202 (50.5%)</p> <p>Placebo group = 198 (49.5%)</p> <p>275 porters were lost to follow-up</p> <p>Most porters (275 porters; 68.75%) dropped out of the trial; 92 porters missed 1 station, 61 porters missed 2 stations, and 122 porters missed all 3 stations Treatment allocation and demographic data were similar in porters who completed the trial and in those who dropped out. 16 porters (4%) were excluded from the analysis, 8 porters for deviating from the standard trek route and 8 porters for noncompliance with medication. Three noncompliers accepted medication from a friend, 3 porters took acetazolamide, 1 porter received medicine from a trekker, and one porter simply failed to take his medication</p> <p>Main characteristics of participants (all groups):</p> <p>Age (median, range): 25, 18 - 54</p> <p>Percentage of men: 100%</p> <p>Weight: 51 kgs, 38 - 66</p>
Interventions	<ol style="list-style-type: none"> 1. Acetazolamide group: 250 mg acetazolamide, orally for 7 days 2. Placebo group: 250 mg orally for 7 days
Outcomes	<p>Outcomes were not classified as primary or secondary</p> <ol style="list-style-type: none"> 1. AMS incidence 2. Related factors 3. Side effects
Notes	<ol style="list-style-type: none"> 1. Trial Registration: Not stated 2. Funder: "Jerwood Foundation and the Sir Samuel Scott of Yews Trust for grants; and the Good Hope Hospital NHS Trust Charitable Fund, the Holy Trinity Parish Church, the Royal Sutton Fun Run, and many individuals for generous donations towards the funding of this study" (Page 93) 3. Role of funder: Wyeth donated acetazolamide

Hillenbrand 2006 (Continued)

	4. A priori sample size estimation: Yes 5. Conducted: October to November 2001 6. Declared conflicts of interest: "The authors have no conflicting interests in this work" (Page 87)	
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "The randomization code was sent directly by DPH to one of the authors of this study, who was not directly involved in performing the clinical trial. He prepared the sealed envelopes containing the trial codes" (Page 88)
Allocation concealment (selection bias)	Low risk	Quote: "The randomization code was sent directly by DPH to one of the authors of this study, who was not directly involved in performing the clinical trial. He prepared the sealed envelopes containing the trial codes" (Page 88)
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Quote: "(...) and a sealed envelope that was only to be opened in the event of illness (...)" (Page 88)
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: "Porters were asked to report to them and were assessed for AMS, using the LLS AMS symptoms score." (Page 88)
Incomplete outcome data (attrition bias) All outcomes	High risk	68.75% of porters dropped out of the trial
Selective reporting (reporting bias)	Low risk	Reporting bias was not detected
Other bias	Low risk	No other biases were identified

Hochapfel 1986

Methods	<ol style="list-style-type: none"> 1. Design: Parallel design (2 arms) 2. Country: Nepal (Annapurna) 3. Multisite: No 4. International: No 5. Treatment duration: 9 days 6. Follow-up: 9 days 7. Final altitude reached: 5500 metres 8. AMS scale: Self-administered subjective questionnaire 9. Randomization unit: patient 10. Analysis unit: patient
Participants	<p>18 trekkers (7 women, 11 men), ages ranged 27 - 53 years, were included. None of them had been at an altitude over 3000 metres over the last 12 months</p> <p>Randomized to:</p> <p>Acetazolamide group: number assigned unclear</p> <p>Placebo group: number assigned unclear</p> <p>Unclear if participants were lost to follow-up</p> <p>Characteristics of participants not reported</p>
Interventions	<ol style="list-style-type: none"> 1. Acetazolamide 250 mg 2. Placebo tablets: no different in form or taste from the acetazolamide tablets
Outcomes	<p>Outcomes were not classified as primary or secondary</p> <ol style="list-style-type: none"> 1. Subjective complaints 2. Onset of headache 3. Side effects
Notes	<ol style="list-style-type: none"> 1. Trial Registration: Not stated 2. Funder: Not stated 3. Role of funder: Not stated 4. A priori sample size estimation: No 5. Conducted: Not stated 6. Declared conflicts of interest: Not stated

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "the batches were distributed in a random process" Insufficient information to score this item as low or high risk of bias
Allocation concealment (selection bias)	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Quote: "None of participants knew the encryption(..)" "The placebo did not differ in the form nor in the taste of the Diamox

Hochapfel 1986 (Continued)

		tablet”
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Selective reporting (reporting bias)	High risk	Unknown number of participants in each arm
Other bias	Low risk	No other biases were identified

Hohenhaus 1994

Methods	<ol style="list-style-type: none"> 1. Design: Randomized trial, parallel, 2 arms 2. Country: Italy 3. Multisite: No 4. International: No 5. Treatment duration: 3 days 6. Follow-up: 1 day 7. Rate of ascent: unclear 8. Final altitude reached: 4559 metres 9. AMS scale: Score proposed at the International Hypoxia Symposium+ “Do you feel ill?” = Yes
Participants	<p>27 mountaineers were recruited. 12 had increased susceptibility to AMS, 8 normal susceptibility and 7 unknown susceptibility</p> <p>Randomized to:</p> <p>Nifedipine group: 14 (51.8%)</p> <p>Placebo group: 13 (48.1%)</p> <p>No participants were lost to follow-up</p> <p>Main characteristics of participants:</p> <p>Age 33 (24 - 60) (placebo); 37 (21 - 54) (nifedipine)</p> <p>Number of men: 9/13 (placebo); 7/14 (nifedipine)</p> <p>Number History of AMS/susceptible: 6/13 (placebo); 7/14 (nifedipine)</p>
Interventions	<ol style="list-style-type: none"> 1. Nifedipine group: Adalat retard, 20 mg. 2. Placebo group: No details provided <p>Medication was given at 10 P.M. on the third and second days before the ascent and at 8 A. M. and 10 P.M. on the day before. Starting on the day of ascent, medication was taken three times daily (at 6 A.M., 2 P.M. and 10 P.M.)</p>
Outcomes	<p>Outcomes were not classified as primary or secondary</p> <ol style="list-style-type: none"> 1. Presence of AMS 2. Blood and end-expiratory gas analysis 3. Pulmonary artery pressure 4. HAPE

Hohenhaus 1994 (Continued)

Notes	1. Trial Registration: Not stated 2. Funder: Italian Alpine Club and Swiss Army 3. Role of funder: providing locations and transportation of the radiographic equipment 4. A priori sample size estimation: No 5. Conducted: Not stated 6. Declared conflicts of interest: No	
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "was assigned randomly in a double-blind design with stratification (...)" (Page 858) Insufficient information to score this item as low or high risk of bias
Allocation concealment (selection bias)	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Incomplete outcome data (attrition bias) All outcomes	Low risk	No participants were lost to follow-up
Selective reporting (reporting bias)	High risk	Participant-important outcomes, such as adverse events, were not reported
Other bias	Low risk	No other biases were identified

Hussain 2001

Methods	<ol style="list-style-type: none"> 1. Design: Parallel (4 arms) 2. Country: Pakistan 3. Multisite: No 4. International: No 5. Treatment duration: 6 days 6. Follow-up: 3 days 7. Rate of ascent: 4578/24 hours 8. Final altitude reached: 4578 metres 9. AMS scale: Modified ESQ 	
Participants	<p>24 participants enrolled (healthy men, low altitude residents at < 500 metres with good health and not suffering from any acute or chronic systemic illness or psychiatric disease)</p> <p>Randomized to:</p> <ul style="list-style-type: none"> Acetazolamide group (n = 6) Placebo Group (n = 6) Dexamethasone group (6) Acetazolamide and dexamethasone group (6) <p>Main characteristics of participants:</p> <ul style="list-style-type: none"> Age (median): global range 25 - 35 years Men: 6 participants in each group History of AMS: None Type of HAI reported: None 	
Interventions	<ol style="list-style-type: none"> 1. Acetazolamide group : 250 mg every 12 hours, started 24 hours before ascent to 4578 metres and continued for 5 days 2. Placebo group: multivitamin tablet every 12 hours, started 24 hours before ascent to 4578 metres and continued for 5 days 3. Dexamethasone group (control): 4 mg tablet every 12 hours, started 24 hours before ascent to 4578 metres and continued for 5 days 4. Acetazolamide and dexamethasone: 250 mg and 4 mg every 12 hours, started 24 hours before ascent to 4578 metres and continued for 5 days 	
Outcomes	<ol style="list-style-type: none"> 1. Primary outcome: Presence of AMS 2. Secondary outcome: Oxygen saturation, severity of AMS 	
Notes	<ol style="list-style-type: none"> 1. Trial Registration: Not stated 2. Funder: Not stated 3. Role of funder: Not stated 4. A priori sample size estimation: No 5. Conducted: Not stated 6. Declared conflicts of interest: No 	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "The study was placebo controlled and the subjects were randomized in double blind fashion into four study groups; that

Hussain 2001 (Continued)

		is, six subjects in each group” Insufficient information to score this item as low or high risk of bias
Allocation concealment (selection bias)	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Incomplete outcome data (attrition bias) All outcomes	Low risk	No participants were reported as lost to follow-up
Selective reporting (reporting bias)	High risk	Participant-important outcomes, such as adverse events, were not reported
Other bias	Low risk	No other biases were identified

Jain 1986

Methods	<ol style="list-style-type: none"> 1. Design: Parallel trial 2. Country: Delhi, India 3. Multisite: No 4. International: No 5. Treatment duration: 4 days 6. Follow-up: 4 days 7. Rate of ascent: Simulate 4570 metres in 1 day 8. Final altitude reached: 4570 metres 9. AMS scale: General High Altitude Questionnaire (GHAQ) 10. Randomization unit: participants 11. Analysis unit: group
Participants	<p>29 participants enrolled (Indian soldiers aged between 22 and 26 years having no previous experience of being at high altitude)</p> <p>Randomized to:</p> <p>Acetazolamide tablets (n = 10)</p> <p>Spironolactone (n = 9)</p> <p>Placebo (n = 10)</p> <p>No participant randomized was excluded or lost to follow-up</p> <p>Main characteristics of participants:</p> <p>Age (median): global ranged 22 - 26 years</p> <p>Men: 100%</p> <p>History of AMS: None</p>

	Type of HAI reported: None	
Interventions	<ol style="list-style-type: none"> 1. Acetazolamide tablets 250 mg every 6 hours beginning a day before the ascent to high altitude 2. Spironolactone tablets 25 mg every 6 hours beginning a day before the ascent to high altitude 3. Placebo tablet every 6 hours beginning a day before the ascent to high altitude 	
Outcomes	<p>Outcomes were not classified as primary or secondary</p> <ol style="list-style-type: none"> 1. Presence of AMS 2. Blood and end-expiratory gas analysis 	
Notes	<ol style="list-style-type: none"> 1. Trial Registration: Not stated 2. Funder: Not stated 3. Role of funder: Not stated 4. A priori sample size estimation: No 5. Conducted: No stated 6. Declared conflicts of interest: Not reported 	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "The subjects were initially tested at an altitude of 200 m and then divided into three groups by using a random number table" (Page 294)
Allocation concealment (selection bias)	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	No participants were reported as lost to follow-up
Selective reporting (reporting bias)	High risk	Participant-important outcomes, such as adverse events, were not reported
Other bias	Low risk	No other biases were identified

Johnson 1984

Methods	<ol style="list-style-type: none"> 1. Design: Double-blind cross-over 2. Country: Boston, USA 3. Multisite: No 4. International: No 5. Treatment duration: 1 day 6. Follow-up: Unclear 7. Rate of ascent: Simulate 4570 metres in 1 day 8. Final altitude reached: 4570 metres 9. AMS scale: ESQ III, AMS-C, and AMS-R questionnaires 	
Participants	<p>12 participants enrolled (healthy men, 20 - 26 years of age, residing at sea level). They were exposed to simulated altitude on 2 separate occasions 4 participants did not participate in the cross-over phase Main characteristics of participants: Age (median): global ranged 22 - 26 years Men: 100% History of AMS: None Type of HAI reported: None</p>	
Interventions	<p>Dexamethasone 4 mg every 6 hours by mouth Placebo</p>	
Outcomes	<p>Outcomes were not classified as primary or secondary <ol style="list-style-type: none"> 1. Presence of AMS 2. AMS-C and AMS-R scores 3. Retinal photography 4. Biochemical and physiological measurements </p>	
Notes	<ol style="list-style-type: none"> 1. Trial Registration: Not stated 2. Funder: US Army Research Institute of Environmental Medicine 3. Role of funder: Technical assistance 4. A priori sample size estimation: No 5. Conducted: Not stated 6. Declared conflicts of interest: Not reported 	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "The treatment order was randomly assessed" (Page 684) Insufficient information to score this item as low or high risk of bias
Allocation concealment (selection bias)	Unclear risk	Insufficient information to score this item as low or high risk of bias

Johnson 1984 (Continued)

Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Incomplete outcome data (attrition bias) All outcomes	High risk	4 participants (33%) were lost to follow-up
Selective reporting (reporting bias)	Low risk	Reporting bias was not detected
Other bias	Unclear risk	It is unclear if previous events of HAI (specifically in phase 1) affected the probability of new events in second phase of cross-over trials

Kayser 2008

Methods	<ol style="list-style-type: none"> 1. Design: Parallel design (3 arms: 2 randomized and 1 open arm) 2. Country: Tanzania (Mount Kilimanjaro) 3. Multisite: No 4. International: No 5. Treatment duration: 5 days 6. Follow-up: 6 days 7. Rate of ascent: 2725 metres/day 1; 1055 metres/day 2; 720 metres/day 3; 960 metres/day 4 8. Final altitude reached: 5896 metres 9. AMS scale: Lake Louise Symptom Score (LLSS) and physician assessment 10. Randomization unit: patient 11. Analysis unit: patient
Participants	<p>93 potential participants (non-acclimatized, altitude-naïve, attempting a fast climb up Mount Kilimanjaro)</p> <p>Exclusion criteria: not reported. 44 participants chose prevention with acetazolamide</p> <p>Randomized to:</p> <p>Calcium carbasalate 15 (48.4%)</p> <p>Placebo 16 (51.6 %)</p> <p>No participants randomized were excluded</p> <p>18 participants lost to follow-up, refusing to participate in any data collection</p> <p>Main characteristics of participants:</p> <p>Age mean (SD): No reported</p> <p>History of AMS: Not stated</p>
Interventions	<p>Intervention:</p> <ol style="list-style-type: none"> 1. acetazolamide 500 mg, oral for 5 days 2. calcium carbasalate 380 mg, 380 mg/day oral, for 5 days

Kayser 2008 (Continued)

	3. Control: placebo	
Outcomes	<p>This trial did not specify by primary or secondary outcomes</p> <ol style="list-style-type: none"> 1. Prevention failure: Headache and LLS score ≥ 3; Headache and LLS + clinical score ≥ 4; Headache and LLS + clinical + functional score ≥ 4 2. HACE: Severe ataxia, vomiting, decreased consciousness 3. Disease-free fast climb experience 	
Notes	<ol style="list-style-type: none"> 1. Trial Registration: Not stated 2. Funder: Dutch tabloid Magazine 3. Role of funder: Provide medical assistance for its readers in the organization of a climb of Mount Kilimanjaro 4. A priori sample size estimation: No 5. Conducted: Not stated 6. Declared conflicts of interest: No 	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "The subjects who agreed to participate in the trial were randomized into two groups stratified for age and sex" (Page 16)
Allocation concealment (selection bias)	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Incomplete outcome data (attrition bias) All outcomes	Low risk	No participants were reported as lost to follow-up
Selective reporting (reporting bias)	High risk	Participant-important outcomes, such as adverse events, were not reported
Other bias	Low risk	No other biases were identified

Ke 2013

Methods	<ol style="list-style-type: none"> 1. Design: Prospective randomized study 2. Country: Nepal 3. Multisite: No 4. International: No 5. Treatment duration: 4 days 6. Follow-up: unclear 7. Rate of ascent: none 8. Final altitude: 3658 metres 9. AMS scale: Lake Louise Score
Participants	<ol style="list-style-type: none"> 1. 28 healthy lowland young men (14 - 22 years old) with no altitude experiences (> 2500 metres) in the preceding 2 years <p>Randomized into 3 groups: Acetazolamide group (n = 9, 32%) Gingko biloba (n = 10, 36%) Placebo (n = 9, 32%)</p> <p>Participants received 3-day pretreatment and 1-day treatment</p> <p>Main characteristics of participants: Age (mean): 19.2 (range 14 - 22 years old) Percentage/number of women/men: 28 men</p>
Interventions	<ol style="list-style-type: none"> 1. Acetazolamide 125 mg twice daily 2. Gingko biloba 120 mg twice daily. 3. Placebo
Outcomes	<ol style="list-style-type: none"> 1. The primary outcome was pulmonary artery systolic pressure (PASP) to hypoxia on the first day 2. Secondary outcomes included: AMS, arterial oxygen saturation (SaO₂), mean artery pressure (MAP), heart rate (HR), and spirometry parameters (FVC, FEV1%, PEF) to hypoxia
Notes	<ol style="list-style-type: none"> 1. Trial Registration: not stated 2. Funder: National Key Technology R&D Program (Grant 2009BAI85B04); National Nature Science Foundation of China (Grant 81172621); and Program for Changjiang Scholars and Innovative Research Team in University 3. A priori sample size estimation: No 4. Conducted: Not stated 6. Declared conflicts of interest: Yes. None declared

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "The participants were randomized into three groups according to random numbers generated by using a software package with nine in the acetazolamide group, ten in the gingko biloba group and nine in the placebo group" (Page 163)

Ke 2013 (Continued)

Allocation concealment (selection bias)	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Quote: “(...) and placebo (provided by the Institute of Pharmaceuticals of the Fourth Military Medical University) were packaged in visually identical capsules at the Institute of Pharmaceuticals of the Fourth Military Medical University (...)” (Page 163)
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Incomplete outcome data (attrition bias) All outcomes	Low risk	No participants were reported as lost to follow-up
Selective reporting (reporting bias)	Low risk	Reporting bias was not detected
Other bias	Low risk	No other biases were identified

Küpper 2008

Methods	<ol style="list-style-type: none"> 1. Design: randomized, doubled-blind, placebo-controlled trial 2. Country: Italy 3. Multisite: No 4. International: No 5. Follow-up: 8 days 6. Treatment duration: 2 days 7. Intention-to-treat: No 8. Follow-up: 24 hours 9. Rate of ascent: first 5 days at 1000 metres, 3440 metres ascent partial, then maximum height of 4560 10. Final altitude reached: 4559 metres 11. AMS scale: Lake Louise AMS questionnaire (LLS)
Participants	<ol style="list-style-type: none"> 1. 24 healthy men eligible. 4 excluded or refused to participate; the reasons for exclusion were sleep disorders, heart disease history, previous episodes of cerebral oedema or high altitude pulmonary 20 participants randomized to receive either 300 mg slow-release theophylline tablets (n = 10) or an identical-appearing placebo (n = 10) Participants lost to follow-up, 1 in the theophylline group and 2 in the placebo group, were unable to ascend to Margherita hut due to adverse weather conditions Main characteristics of participants: Number/Percentage of men: 100% Percentage/number History of AMS: None of the subjects had a history of AMS

Interventions	<ol style="list-style-type: none"> 1. Theophylline group (intervention): 300 mg slow-release tablets, 1 tablet each day at 8 p.m. during 5 days prior to ascent and 2 days 1 night during ascent 2. Placebo group (control): 300 mg identical-appearing placebo tablets, 1 tablet each day at 8 p.m. during 5 days prior to ascent and 2 days 1 night during ascent
Outcomes	<p>This study does not establish primary or secondary outcomes</p> <ol style="list-style-type: none"> 1. Incidence of AMS (AMS-C score \geq 4) 2. Scores of AMS 3. Theophylline levels 4. Sleep hypoxaemia and breathing pattern 5. Polysomnographic parameters
Notes	<ol style="list-style-type: none"> 1. Trial Registration: Not stated 2. Funder: "This investigation was supported by an unrestricted grant of 3M Pharmaceuticals Inc., Neuss, Germany. 3M Pharmaceuticals Inc. also provided the study medication and placebo. Respiroics Inc., Pittsburgh, PA, USA, provided logistic support (sleep recorders and laptops during study duration and helicopter flights for transport of this material). The Margherita hut research lab is supported by several European universities, the Italian Alpine Club, and structural and research funds of the European Union" 3. Role of funder: Not stated 4. A priori sample size estimation: No 5. Conducted: Unclear 6. Declared conflicts of interest: yes. Page 312

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "Participants were randomized (random allocation; see Figure 1) to receive either 300 mg slow-release theophylline tablets (Unilair 300; 3M Pharmaceuticals Inc., Neuss, Germany) or an identical-appearing placebo" (Page 308) Insufficient information to score this item as low or high risk of bias
Allocation concealment (selection bias)	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias

Küpper 2008 (Continued)

Incomplete outcome data (attrition bias) All outcomes	Low risk	3 participants (15%) were lost to follow-up
Selective reporting (reporting bias)	High risk	Participant-important outcomes, such as adverse events, were not reported
Other bias	Unclear risk	Possible industry bias

Larson 1982a

Methods	<ol style="list-style-type: none"> 1. Design: Parallel (2 arms) 2. Country: USA 3. Multisite: No 4. International: No 5. Treatment duration: 24 hours 6. Follow-up: Until 48 hours 7. Rate of ascent: Unclear 8. Final altitude reached: 4394 metres 9. AMS scale: GHAQ modified
Participants	<p>64 participants enrolled (volunteers who normally resided at or near sea level, all in good general health and none had ascended to higher than 3000 metres for at least 4 weeks before participating)</p> <p>Randomized to:</p> <p>Acetazolamide (n = 31, 48.4%)</p> <p>Placebo (n = 33, 51.6%)</p> <p>7 participants lost to follow-up</p> <p>2 participants in acetazolamide group and 3 in placebo did not leave base camp because of excessive fatigue or inadequate clothing; 5 participants in placebo group did not reach the summit, but were included in analysis because they reached at least 3000 metres</p> <p>Main characteristics of participants:</p> <p>Age: range 21 - 48 years</p> <p>Percentage of men: 54 (84.3%) and women:10 (15.3%)</p> <p>Age (mean, SD): Acetazolamide = 28.7 (0.9); Placebo = 29.2 (1)</p> <p>Percentage of men: Acetazolamide= 87.1%; Placebo = 81.8%</p> <p>Pulse rate, beats per minute (mean, SD): Acetazolamide= 65.1 (1.8); Placebo = 64.0 (1.8)</p> <p>There is not enough information on the 6 climbers who ascended twice (cross-over arm)</p>
Interventions	<ol style="list-style-type: none"> 1. Acetazolamide group (intervention): Acetazolamide tablets 250 mg every 8 hours, beginning 1 day before ascent 2. Placebo group (control): Placebo tablets every 8 hours, beginning 1 day before ascent <p>Cointerventions: Not reported</p>
Outcomes	<p>This trial did not specify by primary or secondary outcomes</p> <ol style="list-style-type: none"> 1. AMS assessment (GHAQ scores) at sea level, 1600 metres, 3000 metres, 4394 metres (summit) or high point attained above base camp 2. Spirometric data: resting minute ventilation, expired vital capacity and peak flow, at

Larson 1982a (Continued)

	sea level, 1600 metres, 3000 metres and or near the summit, after resting for at least 10 minutes	
Notes	1. Trial Registration: Not stated 2. Funder: "The acetazolamide (Diamox) and placebo used in this study were provided by Darrel Leichty, Bellevue, Wash, who is a product representative of Lederle Laboratories, Division of American Cyanamid Company, Wayne, NJ" (Page 332) 3. Role of funder: Not stated 4. A priori sample size estimation: No 5. Conducted: Not stated 6. Declared conflicts of interest: Not reported	
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "Using a random numbers table and in a double-blind fashion" (Page 329)
Allocation concealment (selection bias)	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Quote: "Packets containing tablets and data collection forms were prepared by persons not directly involved with the study (..)" (Page 329)
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Incomplete outcome data (attrition bias) All outcomes	Low risk	Around 10% of participants were lost to follow-up
Selective reporting (reporting bias)	High risk	Participant-important outcomes, such as adverse events, were not reported
Other bias	Unclear risk	Possible industry bias

Larson 1982b

Methods	<ol style="list-style-type: none"> 1. Design: Cross-over study 2. Country: USA 3. Multisite: No 4. International: No 5. Treatment duration: Unclear 6. Follow-up: Unclear 7. Rate of ascent: Unclear 8. Final altitude reached: 4394 metres 9. AMS scale: GHAQ modified 	
Participants	<p>6 participants enrolled (volunteers who normally resided at or near sea level, all in good general health and none had ascended to higher than 3000 metres for at least 4 weeks before participating) . Approximately 1 year between the 2 climbs No participants lost to follow-up Main characteristics of participants: No information was provided for these participants</p>	
Interventions	<ol style="list-style-type: none"> 1. Acetazolamide group (intervention): Acetazolamide tablets 250 mg every 8 hours, beginning 1e day before ascent 2. Placebo group (control): Placebo tablets every 8 hours, beginning 1 day before ascent <p>Cointerventions: None reported</p>	
Outcomes	<p>This trial did not specify by primary or secondary outcomes:</p> <ol style="list-style-type: none"> 1. AMS assessment (GHAQ scores) at sea level, 1600 metres, 3000 metres, 4394 metres (summit) or high point attained above base camp 2. Spirometric data: resting minute ventilation, expired vital capacity and peak flow, at sea level, 1600 metres, 3000 metres and or near the summit, after resting for at least 10 minutes 	
Notes	<ol style="list-style-type: none"> 1. Trial Registration: Not stated 2. Funder: "The acetazolamide (Diamox) and placebo used in this study were provided by Darrel Leichthy, Bellevue, Wash, who is a product representative of Lederle Laboratories, Division of American Cyanamid Company, Wayne, NJ" (Page 332) 3. Role of funder: Not stated 4. A priori sample size estimation: No 5. Conducted: Not stated 6. Declared conflicts of interest: Not reported 	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "Using a random numbers table and in a double-blind fashion" (Page 329)
Allocation concealment (selection bias)	Unclear risk	Insufficient information to score this item as low or high risk of bias

Larson 1982b (Continued)

Blinding of participants and personnel (performance bias) All outcomes	Low risk	Quote: “Packets containing tablets and data collection forms were prepared by persons not directly involved with the study (. . .)” (Page 329)
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Incomplete outcome data (attrition bias) All outcomes	Low risk	Around 10% of participants were lost to follow-up
Selective reporting (reporting bias)	High risk	Participant-important outcomes, such as adverse events, were not reported
Other bias	Unclear risk	Possible industry bias. It is unclear if previous events of HAI (specifically in phase 1) affected the probability of new events in second phase of cross-over trials

Lipman 2012

Methods	<ol style="list-style-type: none"> 1. Design: Parallel design (2 arms) 2. Country: USA 3. Multisite: No 4. International: No 5. Treatment duration: 1 day 6. Follow-up: 1 day 7. Rate of ascent: unclear. Aprox 2305 - 2356 metres every 6 hours 8. Final altitude reached: 3810 metres 9. AMS scale: Lake Louise Questionnaire Acute Mountain Sickness Score (LLQ)
Participants	<p>89 participants were recruited through a variety of e-mail list-serves with both local and national distribution, as well as posted advertisements in northern and southern California</p> <p>Randomized to: Placebo 42 Ibuprofeno 44</p> <p>2 participants were excluded post hoc for meeting acute mountain sickness criteria at baseline, and 1 for receiving diuretic medication during the study</p> <p>No participants were lost to follow-up</p> <p>Main characteristics of participants: Age: Placebo 34.8 (13.2), Ibuprofen 38.4 (14.5) Percentage/number of women/men: Placebo 14 women (33.3%), 28 men; Ibuprofen 14 women (31.8%), 40 men Percentage/number History of AMS: Placebo 5 (11.9%), Ibuprofeno 2 (4.6%) Percentage/number Type of HAI reported: Unclear History of headaches: Placebo 2 (4.8%), Ibuprofeno 5 (11.4%)</p>

Interventions	1. Ibuprofen: 600 mg 4 doses of medication at baseline, 3545 metres, 3810 metres and the next morning after descending 2. Placebo: same regimen
Outcomes	1. Primary outcome measures: Incidence and severity of AMS as calculated on the Lake Louise Questionnaire score 2. Secondary outcome measures: headache severity by visual analogue scale and peripheral oxygen saturation by fingertip pulse oximetry (SpO ₂) from baseline
Notes	1. Trial Registration: "Not stated" 2. Funder, role of funder: "This research was made possible by a Research grant from the Division of Emergency Medicine, Stanford University School of Medicine and financial support from the American Alpine Club" 3. A priori sample size estimation: Yes (page 486) 4. Conducted: July and August 2010 5. Declared conflicts of interest: Yes (page 489)

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "Participants were randomized to visually identical commercial-grade ibuprofen 600 mg or placebo, using a computer-generated random sequence, with the randomization code unavailable to administrators and participants" (Page 485)
Allocation concealment (selection bias)	Low risk	Quote: "Participants were randomized to visually identical commercial-grade ibuprofen 600 mg or placebo, using a computer-generated random sequence, with the randomization code unavailable to administrators and participants" (Page 485)
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Incomplete outcome data (attrition bias) All outcomes	Low risk	No participants were lost to follow-up
Selective reporting (reporting bias)	High risk	Participant-important outcomes, such as adverse events, were not reported

Other bias	Low risk	No other biases were identified
------------	----------	---------------------------------

Luks 2007

Methods	<ol style="list-style-type: none"> 1. Design: Randomized, doubled-blind cross-over trial 2. Country: USA 3. Multisite: No 4. International: No 5. Treatment duration: 4 days 6. Follow-up: Until symptoms of AMS became intolerable to the participant or they reached the maximum study duration of 8 hours. Washout time of 2 weeks between 2 observations 8. Rate of ascent (m/h): 158 metres to 3900 metres simulated in a chamber with normobaric hypoxia 9. Final altitude reached: 3900 metres 10 AMS scale: Lake Louise Acute Mountain Sickness scoring survey
Participants	<p>Number enrolled unclear (“Healthy volunteers between the ages of 18 and 55” Page 134)</p> <p>Potential volunteers were excluded from the study if they had chronic pulmonary, cardiac, renal or liver disease, if they had a history of allergies or were already taking anti-inflammatory corticosteroids or medications inhibiting leukotriene synthesis or blocking receptor binding or if they had recently been at high altitude (more than a day at an elevation of 1500 m or higher in the preceding 2 weeks)</p> <p>Randomized to:</p> <p>Montelukast group (n = 10)</p> <p>Placebo group (n = 10)</p> <p>1 participant randomized was excluded because they completed 1 session, but did not return for the second session, because of severe symptoms during the first testing session</p> <p>1 participant lost to follow-up</p> <p>Main characteristics of participants:</p> <p>Age: 24 to 41</p> <p>4 men, 6 women</p> <p>Percentage/number History of AMS: Not stated</p>
Interventions	<ol style="list-style-type: none"> 1. Montelukast group (intervention): 10 mg tablet (Singulair, Merck and Co.) daily for 4 days 2. Placebo group (control): Similar-appearing placebo tablet 3. Co-interventions: for 15 minutes each hour, participants rode a stationary bicycle at a moderate pace in order to simulate the hiking or other physical activity someone might undertake at high altitude
Outcomes	<ol style="list-style-type: none"> 1. Primary outcome measure: Lake Louise Acute Mountain Sickness score at the end of the testing session 2. Secondary outcome measures: Score on the headache component of the Lake Louise scale, the length of time participants were able to remain in the chamber, their average heart rate and arterial blood oxygen saturations throughout their chamber exposure, and the pre- and post-exposure urinary leukotriene E4 concentrations

Luks 2007 (Continued)

Notes	1. Trial Registration: Not stated 2. Funding: Merck Research Laboratories, West Point, Pennsylvania, supported this study 3. Role of sponsor: Not stated 4. A priori sample size estimation: No 5. Conducted: unclear 6. Declared conflicts of interest: Yes (Page 137) “The authors have no other financial support or conflicts of interest to disclose regarding this study”	
Risk of bias		
Bias	Authors’ judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote “ was determined by the flip of a coin...” (Page 132) Insufficient information to score this item as low or high risk of bias
Allocation concealment (selection bias)	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Quote “..neither the subject nor the investigator were aware of the assignment for a particular testing session” (Page 132)
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Incomplete outcome data (attrition bias) All outcomes	High risk	2 participants (20%) were excluded from further analyses
Selective reporting (reporting bias)	High risk	Participant-important outcomes, such as adverse events, were not reported
Other bias	Unclear risk	Possible industry bias. It is unclear if previous events of HAI (specifically in phase 1) affected the probability of new events in second phase of cross-over trials

Methods	<ol style="list-style-type: none"> 1. Design: Randomized, doubled-blind, placebo-controlled trial 2. Countries: Italy, Switzerland 2. Multisite: Yes 3. International: Yes 4. Treatment duration: 3 days 5. Follow-up: 48 hours 6. Rate of ascent: ascended from 1100 metres to 3200 metres by cable car, taking about 1½ hours. Continued by foot to 3600 metres, where they slept overnight, and continued the next morning to 4559 metres in about 4 hours 7. Final altitude reached: 4559 metres 8. AMS scale: Clinical examination by Lake Louise scoring protocol
Participants	<p>29 participants enrolled (mountaineers with a previous history of HAPE)</p> <p>Randomized to:</p> <p>Placebo group (n = 9)</p> <p>Tadalafil group (n = 10)</p> <p>Dexamethasone group (n = 10)</p> <p>2 participants in the Tadalafil group were withdrawn from the study because they developed severe AMS on the evening of arrival at 4559 metres</p> <p>No participants were lost to follow-up</p> <p>Main characteristics of participants:</p> <p>Age (Mean/SD): Placebo group 41/8; tadalafil group 46/3; dexamethasone group 44/3</p> <p>Number of women/men: Placebo group 2/9; Tadalafil group 1/10; dexamethasone group 1/10</p> <p>History of HAPE: (Interquartile range): Placebo group 1 (1 - 3); tadalafil group 1 (1 - 2); dexamethasone 1 (1 - 2)</p>
Interventions	<ol style="list-style-type: none"> 1. Tadalafil group (intervention): Tadalafil 10 mg orally, twice daily started on the morning of the day before ascent to high altitude and continued until the end of the study 2. Dexamethasone group (intervention): Dexamethasone 8 mg twice daily started on the morning of the day before ascent to high altitude and continued until the end of the study 3. Placebo group (control): White gelatin capsules, identical in appearance, containing placebo, twice daily started on the morning of the day before ascent to high altitude and continued until the end of the study
Outcomes	<ol style="list-style-type: none"> 1. Primary outcome: Development of HAPE 2. Secondary outcomes: Incidence of AMS
Notes	<ol style="list-style-type: none"> 1. Trial Registration: Clinical Trials gov identifier: NCT00274430 2. Funder: The Hartmann-Müller Foundation, the Pierluigi Crivelli Foundation, and the Anna Feddersen-Wagner Funds (Switzerland) 3. Role of funder: "The funding sources did not influence the study design; the collection, analysis, or interpretation of the data; or the writing of the manuscript and its submission for publication" 4. A priori sample size estimation: Yes. The group was not able to recruit 54 participants and decided to perform the study after 29 participants had been enrolled 5. Conducted: Not reported 6. Declared conflicts of interest: None disclosed

Maggiorini 2006 (Continued)

<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote "...assigned to individual participants according to a computer-generated list" (Page 498)
Allocation concealment (selection bias)	Low risk	Quote: "Before the study, the pharmacist at the University Hospital Zurich packaged the medication into numbered bottles, which were assigned to individual participants according to a computer-generated list" (Page 498)
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of outcome assessment (detection bias) All outcomes	Low risk	quote: "Two physicians who were blinded to treatment assignment performed clinical examinations according to a predefined checklist in the mornings" (Page 498)
Incomplete outcome data (attrition bias) All outcomes	Low risk	No participants were excluded at follow-up
Selective reporting (reporting bias)	High risk	Participant-important outcomes, such as adverse events, were not reported
Other bias	Low risk	No other biases were identified

Mirakhlmov 1993

Methods	<ol style="list-style-type: none"> 1. Design: Parallel (2 arms) 2. Country: Kirguistán 3. Multisite: No 4. International: No 5. Treatment duration: 1 day 6. Follow-up: Unclear 7. Rate of ascent: Unclear 8. Final altitude reached: 3200 metres 9. AMS scale: No
Participants	<p>16 participants with bronchial asthma were recruited</p> <p>Randomized (single-blinded) into 2 groups:</p> <p>Control group (n = 8, 50%)</p> <p>Intervention group (n = 8, 50%)</p>

	<p>No participants randomized were excluded from the study</p> <p>No participants lost to follow-up:</p> <p>Main characteristics of participants:</p> <p>Age (range): 22 - 49 years</p> <p>Age (mean \pm SD): Intervention group: 34 \pm 3; and Control group: 32 \pm 3</p> <p>Number of men/women: 6 men (37.5%), 10 women (62.5%)</p> <p>Almost all participants had daily bouts of breathlessness, which were relieved by inhaled beta2-agonist</p> <p>5 participants were treated with small doses of prednisolone</p>
Interventions	<p>1. Control group: Anti-asthmatic treatment (control group)</p> <p>2. Intervention group: Anti-asthmatic treatment plus acetazolamide 250 mg twice at day</p>
Outcomes	<p>Outcomes were not classified as primary or secondary</p> <p>1. Severity of nocturnal hypoxaemia in asthmatic participants after the ascent to 3200 metres</p> <p>2. Frequency and severity of AMS and of nocturnal hypoxaemia</p> <p>3. Acclimatization to altitude by repeated overnight oximetry</p>
Notes	<p>1. Trial Registration: Not stated</p> <p>2. Funding: Not stated</p> <p>3. Role of sponsor: Not stated</p> <p>4. A priori sample size estimation: No</p> <p>5. Conducted: not stated</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote "...after the initial investigations, patients were randomly divided.." (Page 537)
Allocation concealment (selection bias)	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Incomplete outcome data (attrition bias) All outcomes	Low risk	No participants were lost to follow-up
Selective reporting (reporting bias)	High risk	Participant-important outcomes, such as adverse events, were not reported

Mirrakhlmov 1993 (Continued)

Other bias	Unclear risk	No other biases were identified
------------	--------------	---------------------------------

Montgomery 1989

Methods	<ol style="list-style-type: none"> 1. Design: Double-blind, randomized study 2. Country: Colorado (Snowmass, Steamboat Springs) USA 3. Multisite: Yes 4. International: No 5. Treatment duration: Unclear 6. Follow-up: 5/6 days 7. Rate of ascent: Unclear 8. Final altitude reached: 2700 and 2050 metres 9. AMS scale: AMS score unclear
Participants	<p>73 persons, mainly health professionals, mostly physicians were recruited and randomized to receive:</p> <p>Dexamethasone (n = 38, 52%)</p> <p>Placebo (n = 35, 48%)</p> <p>No participants were lost to follow-up or excluded</p> <p>Participant characteristics:</p> <p>Placebo n = 35 (14 women, 21 men), age 37.9 ± 7.8 years</p> <p>Dexamethasone n = 38 (10 women, 28 men), age 35.8 ± 6.5 years</p>
Interventions	<p>4 mg of dexamethasone acetate or an identical-appearing placebo every 6 hours for 6 doses</p> <p>Drug administration began within 3 hours after arrival at the ski resorts</p>
Outcomes	AMS symptoms and incidence
Notes	<ol style="list-style-type: none"> 1. Trial Registration: Not stated 2. Funder: Merck Sharpe & Dohme 3. Role of funder: To provided the dexamethasone and the placebo 4. A priori sample size estimation: No 5. Conducted: January 1986 and February 1987 6. Declared conflicts of interest: Not reported

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote ".... was randomized..." (Page 735) Insufficient information to score this item as low or high risk of bias
Allocation concealment (selection bias)	Unclear risk	Insufficient information to score this item as low or high risk of bias

Montgomery 1989 (Continued)

Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Incomplete outcome data (attrition bias) All outcomes	Low risk	No participants were lost to follow-up
Selective reporting (reporting bias)	High risk	Participant-important outcomes, such as adverse events, were not reported
Other bias	Unclear risk	Possible industry bias

Moraga 2007

Methods	<ol style="list-style-type: none"> 1. Design: Randomized, open-label, placebo-controlled trial 2. Country: Chile 3. Multisite: No 4. International: Yes 5. Treatment duration: 4 days 6. Follow-up: 4 days 7. Rate of ascent: Began 0830 hours from Antofagasta (sea level) via highway. Arrival at Calama (2400 metres) at 1230 hours was followed by a 1-hour stop, and arrival at Ollagüe was at 1700 hours. Travel time was approximately 8½ hours 8. Final altitude reached: 3696 metres 9. AMS scale: Lake Louise Questionnaire
Participants	<p>50 participants enrolled (students from the Medical College at the University of Antofagasta voluntarily consented to participate in the study). 13 students were excluded for having previous experience with high altitude. 2 were evaluated by physicians and were excluded for having incidents of seizure and recent pneumonia</p> <p>36 participants randomized to:</p> <p>Ginkgo biloba (12, 33%) Acetazolamide (12, 33%) Placebo (12, 33%)</p> <p>No participants were excluded No participants were lost to follow-up</p> <p>Main characteristics of participants: Age (median/mean- Percentiles 5/95, SD): Placebo 22.2 ± 1.1 Acetazolamide 23.3 ± 1.2 Ginkgo biloba 22.1 ± 2.9</p> <p>Percentage/number of women/men: all men Percentage/number History of AMS: None</p>

Interventions	<p>1. Ginkgo biloba group (intervention): Ginkgo biloba extract Egb761 80 mg/12 hours. Administration route unspecified. At sea level a month before ascending to high altitude for 3 days, at high altitude 24 hours before ascending and continued for 3 days</p> <p>2. Placebo group (control): Administration route unspecified. At sea level a month before ascending to high altitude for 3 days, at high altitude 24 hours before ascending and continued for 3 days</p> <p>3. Acetazolamide group (control): Acetazolamide 250 mg/12 hours. Administration route unspecified. At sea level a month before ascending to high altitude for 3 days, at high altitude 24 hours before ascending and continued for 3 days</p>
Outcomes	Primary outcome was assessment of AMS through the Lake Louise Questionnaire measurement at sea level and at 3696 metres
Notes	<p>1. Trial Registration: Not stated</p> <p>2. Funding: Grant PEI-1332 Project given by the Investigation Unit at the University of Antofagasta, Chile</p> <p>3. Role of sponsor: Not stated</p> <p>4. A priori sample size estimation: No</p> <p>5. Conducted: Not stated</p> <p>6. Declared conflicts of interest: Not reported</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote "randomization was computer generated" (Page 252)
Allocation concealment (selection bias)	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Incomplete outcome data (attrition bias) All outcomes	Low risk	No participants were lost to follow-up
Selective reporting (reporting bias)	High risk	Participant-important outcomes, such as adverse events, were not reported
Other bias	Low risk	No other biases were identified

Muza 2004

Methods	<ol style="list-style-type: none"> 1. Design: Randomized, double-blind, placebo-controlled cross-over trial 2. Country: USA 3. Multisite: No 4. International: No 5. Treatment duration: 2 days 6. Follow-up: 24 hours during each test phase 7. Rate of ascent : 45 mmHg/minute 8. Final altitude reached: 4300 metres 9. AMS scale: ESQ-C score and the Lake Louise AMS Scoring System (LLS) 	
Participants	<p>12 participants enrolled (volunteers lifelong low-altitude residents and had no exposure to altitudes greater than 1000 metres for at least 6 months immediately preceding the study. All were US Army personnel who participated in regular physical training and were of average fitness. All volunteers received medical examinations, and none was found to have any condition that would warrant exclusion from the study)</p> <p>1 participant excluded. No reason given</p> <p>Randomized to:</p> <p>Montelukast (n = 11)</p> <p>Placebo (n = 11)</p> <p>No participants randomized were excluded or lost to follow-up</p> <p>Main characteristics of participants:</p> <p>Age 24 ± 4 years</p> <p>9 men, 2 women</p> <p>Percentage/number History of AMS: None</p> <p>Percentage/number Type of HAI reported: None</p>	
Interventions	<ol style="list-style-type: none"> 1. Intervention group: Montelukast 10 mg was given orally at 08:00 at beginning of a test phase and the second 10 mg dose was given about 24 hours later, just prior to decompressing the chamber to simulated altitude 2. Placebo group (control): An identical-appearing tablet containing lactose was ingested on the same schedule during its corresponding test phase 	
Outcomes	<p>This trial did not specify by primary or secondary outcomes</p> <ol style="list-style-type: none"> 1. AMS assessed by ESQ-C score and LLS score 2. Specific ventilatory, cardiovascular, body fluid, and other physiologic parameters indicative of the early acclimatization process 3. Markers of inflammation and hypoxic stress 	
Notes	<ol style="list-style-type: none"> 1. Trial Registration: Not stated 2. Funding: This investigation was supported by the U.S. Army Medical Research and Materiel Command. Additional support was received from Merck & Co 3. Role of funder: Not stated 4. A priori sample size estimation: No 5. Conducted: received for review in July 2002 6. Declared conflicts of interest: Yes, none reported 	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement

Muza 2004 (Continued)

Random sequence generation (selection bias)	Unclear risk	Quote: “randomized controlled trial” (Page 413) Insufficient information to score this item as low or high risk of bias
Allocation concealment (selection bias)	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Incomplete outcome data (attrition bias) All outcomes	Low risk	1 participant (8.3%) was excluded from analyses
Selective reporting (reporting bias)	High risk	Participant-important outcomes, such as adverse events, were not reported
Other bias	Unclear risk	It is unclear if previous events of HAI (specifically in phase 1) affected the probability of new events in second phase of cross-over trials

PACE 2006

Methods	<ol style="list-style-type: none"> 1. Design: Parallel (3 arms) 2. Country: Nepal 3. Multisite: No 4. International: No 5. Treatment duration: 6 days 6. Follow-up: 24 hours 7. Rate of ascent: Average = 354 metres/day 8. Final altitude reached: 4928 metres 9. AMS scale: Lake Louise questionnaire
Participants	<p>222 participants enrolled (healthy non-Nepali participants between 18 and 65 years of age with no acute infections who had not slept higher than 2700 metres or taken acetazolamide within the last 2 weeks)</p> <p>Randomized to:</p> <p>250 mg acetazolamide group (74, 33.3%)</p> <p>750-mg group (82, 37%)</p> <p>Placebo (66, 29.7%)</p> <p>18 participants lost to follow-up (12%). Reasons not provided</p> <p>Main characteristics of participants:</p>

	<p>Age (mean, SD): placebo = 38, 11.4; 250 mg acetazolamide group = 36.8, 11; 750 mg acetazolamide group = 38.9, 12.6</p> <p>Percentage of men: placebo = 69.5%; 250 mg acetazolamide group = 65.7%; 750 mg acetazolamide group = 60.3%</p> <p>Percentage of History of severe altitude illness: Placebo = 11.9%; 250 mg acetazolamide group = 4.5%; 750 mg acetazolamide group = 11.5%</p> <p>Baseline oxygen saturation (mean, SD): Placebo = 90.9, 2.8; 250 mg acetazolamide group = 91.4, 2.8; 750 mg acetazolamide group = 91.4, 3</p>
Interventions	<p>1. 250 group (intervention): 125 mg oral twice a day for 6 days</p> <p>2. 750 group (intervention): 375 mg oral twice a day for 6 days</p> <p>3. Placebo group (control): placebo capsules oral twice day for 6 days</p> <p>Co-interventions: Not reported</p>
Outcomes	<p>Primary outcomes</p> <p>1. Composite incidence and severity of AMS as measured by the LLQ (AMS = 3+ points on LLQ; severe AMS = 5+ points on LLQ)</p> <p>Secondary outcomes</p> <p>1. Composite headache incidence and severity</p> <p>2. Oxygen saturation decrease from baseline to midpoint and endpoint as measured by resting pulse oximetry</p>
Notes	<p>1. Trial Registration: Not stated</p> <p>2. Funder: These studies were supported by grant 3200-0092.8 5 from the Swiss National Science Foundation</p> <p>3. Role of funder: Not stated</p> <p>4. A priori sample size estimation: Yes</p> <p>5. Conducted: October - November 2003</p> <p>6. Declared conflicts of interest: Not reported</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "Random treatment group assignment codes were prepared by Deurali-Janata and placed in sealed opaque envelopes" (Page 19) Insufficient information to score this item as low or high risk of bias
Allocation concealment (selection bias)	Low risk	Quote: "Random treatment group assignment codes were prepared by Deurali-Janata and placed in sealed opaque envelopes" (Page 19)
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Quote: "The placebo substance was visually identical to the acetazolamide, and both placebo and drug were packed in identical

PACE 2006 (Continued)

		capsules” Page 19 Quote: “(...) in sealed opaque envelopes unavailable to the study administrators who enrolled the patients” (Page 19)
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Incomplete outcome data (attrition bias) All outcomes	Low risk	12% of participants lost to follow-up
Selective reporting (reporting bias)	Low risk	Reporting bias was not detected
Other bias	Unclear risk	Possible industry bias. Quote: “Commercial pharmaceutical-grade acetazolamide was purchased from Wyeth Pharmaceuticals and placed in capsules by Deurali-Janata Pharmaceuticals at their processing plant in Katmandu, Nepal” (Page 19)

Parati 2013

Methods	<ol style="list-style-type: none"> 1. Design: Parallel (2 arms) 2. Country: Italy 3. Multisite: No 4. International: No 5. Treatment duration: 5 days 6. Follow-up: 2 days 7. Rate of ascent (m/h): 4559/28 hours 8. Final altitude reached: 4559 metres 9. AMS scale: Lake Louise Score
Participants	<p>44 participants enrolled (healthy lowlanders without known cardiovascular disease, no chronic cardiovascular therapy, no history of severe mountain sickness, no recent exposure to altitudes > 2000 metres, and no contraindications to acetazolamide)</p> <p>Randomized to:</p> <p>Acetazolamide group (n = 22). 3 participants not analysed</p> <p>Placebo group (n = 22). 2 participants not analysed</p> <p>Main characteristics of participants:</p> <p>Age (median): acetazolamide group 35.6 ± 7.1; Placebo group 37.0 ± 9.5</p> <p>Men: acetazolamide group n = 9; Placebo group n = 10</p> <p>History of AMS: None</p> <p>Type of HAI reported: None</p>
Interventions	<ol style="list-style-type: none"> 1. Acetazolamide group (intervention): acetazolamide 250 mg every 12 hours for 3 days at sea level and continued for 48 hours at high altitude 2. Placebo group (control): tablets every 12 hours for 3 days at sea level and continued for 48 hours at high altitude

Outcomes	1. Primary outcome Central blood pressure, pulse wave velocity 2. Secondary outcome Arterial oxygen saturation Acute Mountain Sickness	
Notes	1. Trial Registration: EudraCT 2010-019986-27 2. Funder: Ministry of Health. IRCCS istituto auxologico italiano 3. Role of funder: Not stated 4. A priori sample size estimation: Yes 5. Conducted: Not stated 6. Declared conflicts of interest: Yes	
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "Subjects were randomly assigned to receive PL or AC, 250 mg" (Page 760) Insufficient information to assess as low or high risk of bias for this item
Allocation concealment (selection bias)	Unclear risk	Insufficient information to assess as low or high risk of bias for this item
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Insufficient information to assess as low or high risk of bias for this item
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to assess as low or high risk of bias for this item
Incomplete outcome data (attrition bias) All outcomes	Low risk	5 participants (11%) were excluded from final analysis
Selective reporting (reporting bias)	High risk	Participant-important outcomes, such as adverse events, were not reported
Other bias	Low risk	No other biases were detected

Methods	<ol style="list-style-type: none"> 1. Design: Parallel (4 arms) 2. Country: Nepal 3. Multisite: No 4. International: No 5. Treatment duration: 2 days 6. Follow-up: Unclear 7. Rate of ascent: Unclear 8. Final altitude reached: 4928 metres 9. AMS scale: Lake Louise score
Participants	<p>614 trekkers were enrolled. They were healthy non-Nepali men and women aged 18 - 65 years travelling directly between the baseline villages of Pheriche or Dingboche (4280 metres and 4358 metres respectively) and the end point in Lobuje (4928 metres)</p> <p>Participants were excluded if they had acute mountain sickness, signs and symptoms of a substantial acute infection, had slept above 4500 metres, had taken ginkgo or acetazolamide within 2 weeks before enrolment, had any known cardiac, pulmonary, or other chronic disease that would render them at increased risk of altitude illness</p> <p>Randomized to:</p> <p>Placebo group (n = 151, 24.5%)</p> <p>Ginkgo group (n = 157, 25.5%)</p> <p>Acetazolamide group (n = 152, 24.7%)</p> <p>Combined acetazolamide and ginkgo group (n = 154, 25%)</p> <p>No participants randomized were excluded from analysis</p> <p>Participants lost to follow-up: 127 (20.7%), uniformly distributed between groups</p> <p>Main characteristics of participants:</p> <p>Age (mean, SD):</p> <p>Placebo group: 36.4, 10.8</p> <p>Acetazolamide group: 36.4, 11</p> <p>Ginkgo group: 36.7, 10.5</p> <p>Combined acetazolamide and ginkgo group: 36.7, 11.4</p> <p>Number of men, %:</p> <p>Placebo group: 88, 74%</p> <p>Acetazolamide group: 79, 67%</p> <p>Ginkgo group: 83, 67%</p> <p>Combined acetazolamide and ginkgo group: 88, 70%</p>
Interventions	<ol style="list-style-type: none"> 1. Ginkgo 120 mg twice daily 2. Acetazolamide 250 mg twice daily 3. Combined ginkgo 120 mg and acetazolamide 250 mg twice daily 4. Placebo twice daily
Outcomes	<p>Primary outcome measure:</p> <ol style="list-style-type: none"> 1. Incidence and severity of acute mountain sickness at the study end point as judged by the Lake Louise scoring system <p>Secondary end points:</p> <ol style="list-style-type: none"> 1. Incidence and severity of headache 2. End point pulse oximetry

Notes	1. Trial Registration: Not stated 2. Funder: Pharmaton provided financial support for study expenses 3. Role of funder: Financial support, manufactured Ginko extract 4. A priori sample size estimation: Yes 5. Conducted: between 6 October and 24 November 2002 7. Declared conflicts of interest: Yes	
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote "the randomisation code was computer generated by Deurali-Janta Pharmaceuticals (Kathmandu, Nepal) and held by an independent physician" (Page 2)
Allocation concealment (selection bias)	Low risk	Quote "the randomisation code was computer generated by Deurali-Janta Pharmaceuticals (Kathmandu, Nepal) and held by an independent physician" (Page 2)
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Incomplete outcome data (attrition bias) All outcomes	High risk	Quote: "The 127 participants (20.7%) lost to follow up..." (Page 2)
Selective reporting (reporting bias)	Low risk	Reporting bias was not detected.
Other bias	Unclear risk	Possible industry bias: The sponsor manufactured the Ginko extract used

Rock 1987

Methods	<ol style="list-style-type: none"> 1. Design: Paralell (2 arms) 2. Country: USA 3. Multisite: No 4. International: No 5. Treatment duration: 96 hours 6. Follow-up: 6 days 7. Rate of ascent: 708.3. Travel by helicopter in < 6 hours 9. Final altitude reached: 4300 metres 10. AMS scale: ESQ-C, ESQ-R, Hackett score, Jhonson Score 	
Participants	<p>16 men enrolled ((volunteers; lifelong sea level residents without exposure to altitudes > 1000 metres for at least 6 months prior to their participation)</p> <p>Exclusion criteria: Any illness or medical contraindication to altitude exposure or to dexamethasone administration</p> <p>Randomized to:</p> <p>Control group (9, 56%)</p> <p>Intervention group (7, 44%)</p> <p>1 participant randomized was excluded from the control group for chest pain</p> <p>No participants lost to follow-up</p> <p>Main characteristics of participants:</p> <p>Age (range): 16 - 26 years</p> <p>Number of men/women: 100% men</p>	
Interventions	<ol style="list-style-type: none"> 1. Treatment group (intervention): 4 mg dexamethasone orally every 6 hours for 48 hours at sea level and 48 hours after arrival at high altitude 2. Control group (control): identically-appearing placebo orally with the same schedule 	
Outcomes	<p>This trial did not specify by primary or secondary outcomes</p> <ol style="list-style-type: none"> 1. AMS symptoms by several scales 2. Haematocrit and haemoglobin 	
Notes	<ol style="list-style-type: none"> 1. Trial Registration: Not stated 2. Funder: Not stated 3. Role of funder: Not stated 4. A priori sample size estimation: No 5. Conducted: Not stated 6. Declared conflicts of interest: Not reported 	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	<p>Quote: "Subjects had been assigned at random to either a treatment or control.." (Page 669)</p> <p>Insufficient information to score this item as low or high risk of bias</p>

Rock 1987 (Continued)

Allocation concealment (selection bias)	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Quote: “control group followed the same drug altitude schedule, but received an identically appearing placebo” (Page 669) Insufficient information to score this item as low or high risk of bias
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: “At the time of each assessment the physicians were unaware of which treatment the subject was receiving” (Page 669)
Incomplete outcome data (attrition bias) All outcomes	Low risk	1 participant (6.25%) was excluded from further analyses
Selective reporting (reporting bias)	High risk	Participant-important outcomes, such as adverse events, were not reported
Other bias	Unclear risk	No other biases were identified

Rock 1989a

Methods	<ol style="list-style-type: none"> 1. Design: Cross-over. 2. Country: USA 3. Multisite: No 4. International: No 5. Treatment duration: 52 hours 6. Follow-up: Unclear 7. Rate of ascent: 600 metres/minute. Hypobaric chamber 8. Final altitude reached: 4570 metres 9. AMS scale: ESQ, AMS-C, AMS-R, Johnson scale
Participants	<p>30 young, healthy men, lifelong residents at low altitude, without any prolonged exposure to altitudes > 2500 metres in the 6 months immediately preceding the study were randomized. 2 of them were unable to participate and 3 were excluded</p> <p>Exclusion criteria: not stated</p> <p>Randomized to</p> <p>Dexamethasone 0.25 mg (n = 8)</p> <p>Placebo. Each subject served as their own control</p> <p>2 participants randomized were excluded from analysis, because they were unable to participate for personal reasons, prior to the beginning of testing</p> <p>3 participants lost to follow-up: 2 were excluded for viral illness and one withdrew for administrative reasons. The data from these 3 individuals were not included in the analysis</p> <p>Main characteristics of participants:</p> <p>Age (mean, SD): 22.3, 2.4 years</p> <p>Number of men: 100%</p>

Rock 1989a (Continued)

Interventions	1. Dexamethasone 0.25 mg orally every 12 hours 2. Placebo identically-appearing, containing lactose. orally every 12 hours Exposures into the chamber were 3 weeks apart
Outcomes	This trial did not state primary or secondary outcome 1. AMS incidence 2. Physiological variables such as haemoglobin, plasma volume, urine output 3. Cortisol levels
Notes	1. Trial Registration: Not stated. 2. Funder: Not stated 3. Role of funder: Not stated 4. A priori sample size estimation: No 4. Conducted: Not stated 6. Declared conflicts of interest: No

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "30 subjects were assigned at random by an individual not involved in the data collection" (Page 569) Insufficient information to score this item as low or high risk of bias
Allocation concealment (selection bias)	Low risk	Quote: "30 subjects were assigned at random by an individual not involved in the data collection" (Page 569)
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Quote: "Neither the subjects nor the investigators collecting the data were aware of which treatment the subjects were receiving during drug administration and data collection" (Page 569)
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: "The clinical interview was performed by a physician (R.F.L.) who was unaware of the subject's responses on the ESQ at the time of the interview" (Page 569) Quote: "Neither the subjects nor the investigators collecting the data were aware of which treatment the subjects were receiving during drug administration and data collection" (Page 569)
Incomplete outcome data (attrition bias) All outcomes	Low risk	5 participants (16%) were lost to follow-up

Rock 1989a (Continued)

Selective reporting (reporting bias)	High risk	Participant-important outcomes, such as adverse events, were not reported
Other bias	Unclear risk	It is unclear if previous events of HAI (specifically in phase 1) affected the probability of new events in second phase of cross-over trials. Possible industry bias

Rock 1989b

Methods	<ol style="list-style-type: none"> 1. Design: Cross-over. 2. Country: USA 3. Multisite: No 4. International: No 5. Treatment duration: 52 hours 6. Follow-up: Unclear 7. Rate of ascent: 600 metres/minute. Hypobaric chamber 8. Final altitude reached: 4570 metres 9. AMS scale: ESQ, AMS-C, AMS-R, Johnson scale
Participants	<ol style="list-style-type: none"> 1. 30 young, healthy men, lifelong residents at low altitude, without any prolonged exposure to altitudes > 2500 metres in the 6 months immediately preceding the study were randomized. 2 of them were unable to participate and 3 were excluded Exclusion criteria: not stated Dexamethasone 1 mg: 9 participants Placebo. Each participant served as their own control Two participants randomized were excluded from analysis, because they were unable to participate for personal reasons, prior to the beginning of testing 3 participants lost to follow-up: 2 were excluded for viral illness and 1 withdrew for administrative reasons. The data from these 3 individuals were not included in the analysis 4. Main characteristics of patients: Age (mean, SD): 22.3, 2.4 years Number of men: 100%
Interventions	<ol style="list-style-type: none"> 1. Dexamethasone 1 mg orally every 12 hours 2. Placebo identically-appearing, containing lactose, orally every 12 hours
Outcomes	<p>This RCT did not state primary or secondary outcome</p> <ol style="list-style-type: none"> 1. AMS incidence 2. Physiological variables such as haemoglobine, plasma volume, urine output 3. Cortisol levels
Notes	<ol style="list-style-type: none"> 1. Trial Registration: Not stated 2. Funder: Not stated 3. Role of funder: Not stated 4. A priori sample size estimation: No 5. Conducted: Not stated

6. Declared conflicts of interest: No		
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "30 subjects were assigned at random by an individual not involved in the data collection" (Page 569) Insufficient information to score this item as low or high risk of bias
Allocation concealment (selection bias)	Low risk	Quote: "30 subjects were assigned at random by an individual not involved in the data collection" (Page 569)
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Quote: "Neither the subjects nor the investigators collecting the data were aware of which treatment the subjects were receiving during drug administration and data collection" (Page 569)
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: "The clinical interview was performed by a physician (R.F.L.) who was unaware of the subject's responses on the ESQ at the time of the interview" (Page 569) Quote: "Neither the subjects nor the investigators collecting the data were aware of which treatment the subjects were receiving during drug administration and data collection" (Page 569)
Incomplete outcome data (attrition bias) All outcomes	Low risk	5 participants (16%) were lost to follow-up
Selective reporting (reporting bias)	High risk	Participant-important outcomes, such as adverse events, were not reported
Other bias	Unclear risk	It is unclear if previous events of HAI (specifically in phase 1) affected the probability of new events in second phase of cross-over trials. Possible industry bias

Rock 1989c

Methods	<ol style="list-style-type: none"> 1. Design: Cross-over. 2. Country: USA 3. Multisite: No 4. International: No 5. Treatment duration: 52 hours 6. Follow-up: Unclear 7. Rate of ascent: 600 metres/minute. Hypobaric chamber 8. Final altitude reached: 4570 metres 9. AMS scale: ESQ, AMS-C, AMS-R, Johnson scale 	
Participants	<ol style="list-style-type: none"> 1. 30 young, healthy men, lifelong residents at low altitude, without any prolonged exposure to altitudes > 2500 m in the 6 months immediately preceding the study were randomized. 2 of them were unable to participate and 3 were excluded <p>Exclusion criteria: not stated</p> <p>Dexamethasone 4 mg: 8 participants</p> <p>Placebo. Each participant served as their own control</p> <ol style="list-style-type: none"> 2 participants randomized were excluded from analysis, because they were unable to participate for personal reasons, prior to the beginning of testing 3 participants lost to follow-up: 2 were excluded for viral illness and 1e withdrew for administrative reasons. The data from these 3 individuals were not included in the analysis <p>Main characteristics of participants:</p> <p>Age (mean, SD): 22.3, 2.4 years</p> <p>Number of men: 100%</p>	
Interventions	<ol style="list-style-type: none"> 1. Dexamethasone 4 mg orally every 12 hours 2. Placebo identically-appearing, containing lactose, orally every 12 hours <p>Exposures into the chamber were 3 weeks apart</p>	
Outcomes	<p>This trial did not state primary or secondary outcome</p> <ol style="list-style-type: none"> 1. AMS incidence 2. Physiological variables 3. Cortisol levels 	
Notes	<ol style="list-style-type: none"> 1. Trial Registration: Not stated 2. Funder: Not stated 3. Role of funder: Not stated 4. A priori sample size estimation: No 5. Conducted: Not stated 6. Declared conflicts of interest: No 	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "30 subjects were assigned at random by an individual not involved in the data collection" (Page 569) Insufficient information to score this item as low or high risk of bias

Rock 1989c (Continued)

Allocation concealment (selection bias)	Low risk	Quote: “30 subjects were assigned at random by an individual not involved in the data collection” (Page 569)
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Quote: “Neither the subjects nor the investigators collecting the data were aware of which treatment the subjects were receiving during drug administration and data collection” (Page 569)
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Quote: “The clinical interview was performed by a physician (R.F.L.) who was unaware of the subject’s responses on the ESQ at the time of the interview” (Page 569) Quote: “Neither the subjects nor the investigators collecting the data were aware of which treatment the subjects were receiving during drug administration and data collection” (Page 569)
Incomplete outcome data (attrition bias) All outcomes	Low risk	5 participants (16%) were lost to follow-up
Selective reporting (reporting bias)	High risk	Participant-important outcomes, such as adverse events, were not reported
Other bias	Unclear risk	It is unclear if previous events of HAI (specifically in phase 1) affected the probability of new events in second phase of cross-over trials. Possible industry bias

Sartori 2002

Methods	<ol style="list-style-type: none"> 1. Design: parallel study (2 arms) 2. Country: Italy 3. Multisite: No 4. International: No 5. Treatment duration: 3 days 6. Follow-up: Unclear 7. Rate of ascent: 155.8 metres/hour 8. Final altitude reached: 4559 metres 9. AMS scale: Lake Louise AMS scoring
Participants	37 participants out of 51 with a previous event of HAPE (at least 1 radiographically-documented episode of high-altitude pulmonary oedema within the previous 4 years), were randomized to: Salmeterol group (n = 18, 48.6%)

	Placebo group (n = 19, 51.4%) No participants randomized were excluded from analysis or lost to follow-up Main characteristics of participants: Age (mean, SD): Salmeterol group: 49.6 ± 10.2 Placebo group: 46 ± 12.6 Percentage of women/men: Salmeterol group 5/13 Placebo group 4/15 History of AMS (number of previous episodes): Salmeterol group 2.4 ± 1 Placebo group 1.9 ± 1.1	
Interventions	1. Salmeterol group (intervention): 125 µg salmeterol every 12 hours with pressurized metred-dose inhaler 2. Placebo group (control): inhaled placebo pressurized metred dose inhaler every 12 hours Both groups started on the morning of the day before began the ascent and continued until the end of the study Co-interventions: Not reported	
Outcomes	This trial did not specify by primary or secondary outcomes 1. Incidence of HAPE 2. Lake Louise Score 3. Systolic pulmonary-artery pressure (by echocardiography) 4. SaO ₂ , PaO ₂ , PaCO ₂	
Notes	1. Trial Registration: Not stated 2. Funder: Swiss National Science Foundation (grants 32.46797.96 and 3238-051157.97), the Placide Nicod Foundation, the Emma Muschamp Foundation, and the International Olympic Committee 3. Role of funder: Not stated 4. A priori sample size estimation: No 5. Conducted: Not stated 6. Declared conflicts of interest: No	
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "... were randomly assigned to inhale either..." (Page 1632) Insufficient information to score this item as low or high risk of bias
Allocation concealment (selection bias)	Unclear risk	Insufficient information to score this item as low or high risk of bias

Sartori 2002 (Continued)

Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Incomplete outcome data (attrition bias) All outcomes	Low risk	No participants were lost to follow-up
Selective reporting (reporting bias)	High risk	Participant-important outcomes, such as adverse events, were not reported
Other bias	Low risk	No other biases were identified

SPACE 2011

Methods	<ol style="list-style-type: none"> 1. Design: Parallel (3 arms) 2. Country: Nepal 3. Multisite: No 4. International: No 5. Treatment duration: 30 hours - 4 days 6. Follow-up: Unclear 7. Rate of ascent: Unclear 8. Final altitude reached: 5000 metres 9. AMS scale: Lake Louise score
Participants	<p>311 participants enrolled (healthy men and women between 18 and 65 years without AMS or any concurrent illness and not taking acetazolamide)</p> <p>Exclusion criteria: Mild AMS (more than 1 mild symptom on the LLS); significantly depressed oxygen saturation (< 75%); pregnancy or those who could not exclude the possibility of being pregnant or have missed menses by over 7 days; history of allergy to acetazolamide or other sulfa drugs; individuals who were on ACE inhibitors (e.g. enalapril) or other diuretics (e.g. amiloride or triamterene); individuals who had spent 24 hours at an altitude of 4500 metres (14,000 feet) within the last 9 days; individuals known to have taken any of the following in the prior 2 days: acetazolamide (Diamox), steroids (dexamethasone, prednisone), theophylline, or diuretics (furosemide); individuals failing to provide informed consent at the study enrolment site at Pheriche</p> <p>Randomized to:</p> <ul style="list-style-type: none"> 114 Spironolactone, 36.6% 118 Acetazolamide, 37.9% 79 Placebo, 25.4% <p>25 participants randomized (8%, uniformly distributed) were excluded from analysis because they violated the protocol:</p> <ul style="list-style-type: none"> Acetazolamide group (8, 7,7 %) Spironolactone group (10, 9,8%) Placebo group (7, 9,8%)

	<p>Participants lost to follow-up: Acetazolamide group: n = 15, 12% Spironolactone group: n = 12, 10.5% Placebo group: n = 8, 10% Main characteristics of participants: Age (mean, SD): Acetazolamide group 37, 12.2 Spironolactone group 37.7, 12 Placebo group 39.4, 12.1 Number of men, %: Acetazolamide group 59 (62.1%) Spironolactone group 67 (62.8%) Placebo group 46 (71.9%)</p>
Interventions	<ol style="list-style-type: none"> 1. Acetazolamide group (intervention): acetazolamide 250 mg twice a day orally for 4 days 2. Spironolactone group (intervention): Spironolactone 50 mg twice a day orally for 4 days 3. Placebo group (control): placebo twice a day orally for 4 days
Outcomes	<p>Primary outcome. Incidence and severity of AMS Secondary outcome: Incidence of headache together with severity of AMS SpO₂</p>
Notes	<ol style="list-style-type: none"> 1. Trial Registration: ISRCTN77054547 2. Funder: Wellcome Trust, UK 3. Role of funder: Financial support 4. A priori sample size estimation: no 5. Conducted: October 6 and November 24, 2007 6. Declared conflicts of interest: no

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote : "randomization of spironolactone, acetazolamide, and placebo was conducted by Deurali-Janta Pharmaceuticals Pvt. Ltd" (Page 17) Insufficient information to score this item as low or high risk of bias
Allocation concealment (selection bias)	Unclear risk	Quote : "randomization of spironolactone, acetazolamide, and placebo was conducted by Deurali-Janta Pharmaceuticals Pvt. Ltd" (Page 17) Quote: "Three sealed master lists of the

SPACE 2011 (Continued)

		randomization code were held by the manufacturer, an independent clinician at the Nepal International Clinic in Katmandu, and an independent clinician at the aid post in Pheriche (study enrollment location)” (Page 17)
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Incomplete outcome data (attrition bias) All outcomes	Low risk	Around 10 - 12% of participants were lost to follow-up
Selective reporting (reporting bias)	High risk	Participant-important outcomes, such as adverse events, were not reported
Other bias	Low risk	No other biases were identified

Subudhi 2011

Methods	<ol style="list-style-type: none"> 1. Design: Cross-over design (3 arms) 2. Country: USA 3. Multisite: No 4. International: No 5. Treatment duration: 2 days 6. Follow-up: Unclear 7. Final altitude reached: simulated altitude of 4875 metres 8. AMS scale: Lake Louise Score
Participants	<p>29 healthy volunteers who had resided at 1650 metres for at least 1 year were screened. All had to accept each treatment Acetazolamide 250 mg Dexamethazone 4 mg Placebo</p> <p>Exclusion criteria: recent (1 month) exposure to altitudes above 2500 metres; medical conditions affected by hypoxia, or poor aerobic fitness</p> <p>9 participants (31%) randomized dropped out of the study “due to the large time commitment required to obtain an additional trial” (Page 1220). They were excluded from the analysis</p> <p>Participants lost to follow-up: None stated</p> <p>Main characteristics of participants: Age (mean, SD): age not stated Number of men, %: 16, 80%</p>

Interventions	<ol style="list-style-type: none"> 1. Acetazolamide 250 mg every 8 hours 2. Dexamethazone 4 mg every 8 hours 3. Placebo every 8 hours
Outcomes	<p>This trial does not state primary or secondary outcomes</p> <ol style="list-style-type: none"> 1. Physiological cardiopulmonary variables: heart rate, SpO₂, pulmonary function 2. Cerebral haemodynamic variables: Cerebral blood flow (doppler), critical closing pressure; resistance area product; cerebral vasomotor reactivity to CO₂; cerebrovascular conductance index 3. AMS score self-reported
Notes	<ol style="list-style-type: none"> 1. Trial Registration: Not reported 2. Funder: National Heart, Lung, and Blood Institute, Marren Foundation and the Altitude Research Center 3. Role of funder: Financial support 4. A priori sample size estimation: Not stated 5. Conducted: Unclear 6. Declared conflicts of interest: Yes

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote "Using a randomized, double-blind, placebo controlled, crossover design, we evaluated..." (Page 1220) Insufficient information to score this item as low or high risk of bias
Allocation concealment (selection bias)	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Incomplete outcome data (attrition bias) All outcomes	High risk	31% (9/29) of participants were lost to follow-up
Selective reporting (reporting bias)	High risk	Participant-important outcomes, such as adverse events, were not reported
Other bias	Unclear risk	It is unclear if previous events of HAI (specifically in phase 1) affected the prob-

ability of new events in second phase of cross-over trials. Possible industry bias

Van Patot 2008

Methods	<ol style="list-style-type: none"> 1. Design: Parallel (2 arms) 2. Country: USA 3. Multisite: No 4. International: No 5. Treatment duration: 4 days 6. Follow-up: Unclear 7. Rate of ascent: Unclear 8. Final altitude reached: 4300 metres 9. AMS scale: Lake Louise score and ESQ AMS-C
Participants	<p>44 participants who resided between 1400 and 1600 metres were randomized to: Acetazolamide n = 22, 50% Placebo n = 22, 50%</p> <p>Exclusion criteria: Pregnancy; history of cardiac/pulmonary disease (except asthma); alcohol consumption within 24 hours prior to ascent; current viral illness; if they had been above 2000 metres for more than 1 day in the preceding 2 weeks</p> <p>No participants randomized were excluded from analysis</p> <p>Participants lost to follow-up: None</p> <p>Main characteristics of participants: Age (years): Mean (SD): Acetazolamide: 22.9 (5.37) Placebo: 23.7 (6.29) Sex (% men): 56% (18/33) Acetazolamide 52% Placebo 43%</p>
Interventions	<ol style="list-style-type: none"> 1. Acetazolamide 125 mg twice a day for 3 days prior to ascent and for 24 hours while at high altitude 2. Placebo (lactulosa) twice a day for 3 days prior to ascent and for 24 hours while at high altitude
Outcomes	<p>Primary outcome:</p> <ol style="list-style-type: none"> 1. Incidence and severity of AMS based on the AMS-C score and Lake Louise Symptom score <p>Secondary outcome:</p> <ol style="list-style-type: none"> 1. Oxygen saturation and heart rate
Notes	<ol style="list-style-type: none"> 1. Trial Registration: Not stated 2. Funder: Technical Sourcing International, the Wilderness Medicine Society, and the American Academy of Family Physicians Foundation 3. Role of funder: Financial support 4. A priori sample size estimation: No 5. Conducted: Not stated 6. Declared conflicts of interest: Yes

Van Patot 2008 (Continued)

<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote "...randomized to either acetazolamide or placebo treatments using a random-number assignment program" (Page 290)
Allocation concealment (selection bias)	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Incomplete outcome data (attrition bias) All outcomes	Low risk	No participants were lost to follow-up
Selective reporting (reporting bias)	High risk	Participant-important outcomes, such as adverse events, were not reported
Other bias	Low risk	No other biases were identified

Wang 2013

Methods	<ol style="list-style-type: none"> 1. Design: Prospective intervention study 2. Country: China 3. Multisite: No 4. International: No 5. Treatment duration: 4 days 6. Follow-up: Unclear 7. Rate of ascent: None 8. Final altitude: 3651 metres 9. AMS scale: Lake Louise Score
Participants	<p>21 healthy young men (22 - 26 years old) with the following characteristics were recruited: altitude of permanent residence less than 900 metres; no high-altitude exposures (\geq 2500 metres) in the preceding 2 years; no tobacco or recreational drug use; not taking medications that might affect cognitive function or carbonic anhydrase activity; no chronic or genetic diseases; being willing to participate in the study and take the medicine provided; no history of allergy to sulfonamides</p> <p>Randomized to: Acetazolamide group (n = 11, 52.3%)</p>

	Placebo (n = 10, 47.6%) Main characteristics of participants: Age (mean): 19.2 (range 14 - 22 years old) Percentage/number of women/men: 21 men (100%)
Interventions	1. Acetazolamide 125 mg twice daily, for 4 days 2. Placebo twice daily for 4 days
Outcomes	Outcome were not classified as primary or secondary. 1. AMS at high altitude 2. Effects of acute high-altitude exposure on neuropsychological performance 3. Effects of acetazolamide on neuropsychological performance
Notes	1. Trial Registration: not stated 2. Funder: "This study was sponsored by the National Key Technology R&D Program (2009BAI85B04), the National Nature Science Foundation of China (81172621), and the Program for Changjiang Scholars and Innovative Research Team in University (PC-SIRT)" 3. A priori sample size estimation: No 4. Conducted: Not stated 5. Declared conflicts of interest: Yes. None declared

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "Twenty-one volunteers were randomized into the acetazolamide group (n = 11) and the placebo group (n = 10)" (Page 29) Insufficient information to score this item as low or high risk of bias
Allocation concealment (selection bias)	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Quote: "Both performers and subjects were blind to treatment assignment during the trial" (Page 29)
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Incomplete outcome data (attrition bias) All outcomes	Low risk	No participants were lost to follow-up
Selective reporting (reporting bias)	High risk	Participant-important outcomes, such as adverse events, were not reported

Wang 2013 (Continued)

Other bias	Low risk	No other biases were identified
------------	----------	---------------------------------

Wright 1983

Methods	<ol style="list-style-type: none"> 1. Design: Parallel (2 arms) 2. Country: Kenya 3. Multisite: No. 4. International: No. 5. Treatment duration: 18 days 6. Follow-up: 10 days 7. Rate of ascent: Unclear 8. Final altitude reached: 4985 (14 participants) or 5188 metres (6 participants) 9. AMS scale: Standard series of questions, clinical assessment 	
Participants	<p>20 participants enrolled (normally resident at less than 200 metres, none had travelled to high altitude within the previous 6 months)</p> <p>Exclusion criteria: not stated</p> <p>Randomized to:</p> <p>Acetazolamide group (10, 50%)</p> <p>Methazolomide (10, 50%)</p> <p>None of the participants randomized were excluded from analysis</p> <p>No participants lost to follow-up</p> <p>Main characteristics of participants not stated</p> <p>Age (years): mean 36, range 22 - 54</p> <p>Number of men, %: 19, 95%</p>	
Interventions	<ol style="list-style-type: none"> 1. Acetazolamide group (intervention): 2 capsules of 250 mg of acetazolamide + inactive capsule daily 8 days before ascent and until the end of observation period (10 days) 2. Methazolomide group (control): 2 capsules of 50 mg of methazolamide + inactive capsule for the first 5 days and 3 capsules of 50 mg for the remaining 10 days 	
Outcomes	<p>This RCT did not specify by primary or secondary outcomes</p> <ol style="list-style-type: none"> 1. Clinical assessment of AMS 2. Blood gas measurements. PaO₂, SaO₂, PaCO₂ 3. Paraesthesia 	
Notes	<ol style="list-style-type: none"> 1. Trial Registration: Not stated 2. Funder: Lederle Laboratories, the Arthur Thompson Trust Fund, the West Midlands Regional Health Authority, and others (Page 621) 3. Role of funder: Financial support 4. A priori sample size estimation: No 5. Conducted: Not stated 6. Declared conflicts of interest: Not reported 	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement

Wright 1983 (Continued)

Random sequence generation (selection bias)	Unclear risk	Quote “ randomly allocated...” (Page 620) Insufficient information to score this item as low or high risk of bias
Allocation concealment (selection bias)	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Insufficient information to score this item as low or high risk of bias
Incomplete outcome data (attrition bias) All outcomes	Low risk	No participants were lost to follow-up
Selective reporting (reporting bias)	High risk	Participant-important outcomes, such as adverse events, were not reported
Other bias	Unclear risk	Possible industry bias

Wright 2004

Methods	<ol style="list-style-type: none"> 1. Design: Parallel (4 arms) 2. Country: Nepal 3. Multisite: No 4. International: No 5. Treatment duration: Unclear 6. Follow-up: Unclear 7. Rate of ascent: Unclear 8. Final altitude reached: 5200 metres 9. AMS scale: Lake Louise self-reporting AMS questionnaire
Participants	<p>24 participants enrolled (no information provided) Exclusion criteria: not stated Randomized to: Medroxyprogesterone group (6, 25%). Acetazolamide group (6, 25%). Acetazolamide + medroxyprogesterone group (6, 25%). Placebo group (6, 25%)</p> <p>1 participant randomized to acetazolamide was excluded from analysis, because he descended with an unrelated illness No participants lost to follow-up Main characteristics of participants not provided Age (years): range 22 - 65 years Number of men, %: 92%</p>

Wright 2004 (Continued)

Interventions	<ol style="list-style-type: none"> 1. Medroxyprogesterone group (intervention): 3 tablets of 10 mg twice daily 2. Acetazolamide group (intervention): 250 mg twice daily + placebo (3 tablets twice daily) 3. Acetazolamide + medroxyprogesterone group (intervention): 250 mg twice daily + 3 tablets of 10 mg twice daily 4. Placebo group (control): 3 tablets of 50 mg twice daily
Outcomes	<p>This trial did not specify by primary or secondary outcomes</p> <ol style="list-style-type: none"> 1. AMS incidence using LLS 2. AMS symptoms 3. Blood gases
Notes	<ol style="list-style-type: none"> 1. Trial Registration: Not stated 2. Funder: The Wellcome Trust, the Arthur Thompson Trust, the Mount Everest Foundation, Ciba Corning Diagnostics UK and Upjohn Ltd (Page 30) 3. Role of funder: Not stated 4. A priori sample size estimation: No 5. Conducted: Not stated 6. Declared conflicts of interest: Not reported

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "Study medications were randomized via computer-generated code" (Page 237)
Allocation concealment (selection bias)	Low risk	Insufficient information to score this item as low or high risk of bias
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Incomplete outcome data (attrition bias) All outcomes	Low risk	22.7% of participants were lost to follow-up and not include in final analysis
Selective reporting (reporting bias)	High risk	Participant-important outcomes, such as adverse events, were not reported
Other bias	Unclear risk	Possible industry bias

Methods	<ol style="list-style-type: none"> 1. Design: Parallel (4 arms) 2. Country: USA 3. Multisite: No 4. International: No 5. Treatment duration: 4 days 6. Follow-up: Unclear 7. Rate of ascent: Unclear 8. Final altitude reached: 4050 metres 9. AMS scale: ESQ 	
Participants	<p>32 participants enrolled (novice backpackers having no previous history of AMS and no recent travel to high altitudes)</p> <p>Exclusion criteria: Ongoing cardiopulmonary issues; Glucose intolerance or diabetes mellitus</p> <p>Randomized to:</p> <ul style="list-style-type: none"> Dexamethasone group (n = 9) Acetazolamide (n = 7) Dexamethasone + acetazolamide group (n = 8) Placebo group (n = 8) <p>Main characteristics of participants:</p> <ul style="list-style-type: none"> Age (median): 18 - 49 years for all groups Number of women/men: 12 women/20 men History of AMS: None 	
Interventions	<ol style="list-style-type: none"> 1. Dexamethasone group: dexamethasone acetate 4 mg orally every 6 hours for 96 hours 2. Acetazolamide group :Acetazolamide 250 mg twice a day oral for 96 hours 3. Placebo group: 2 vials of unmarked medications, 1 of which was taken twice a day and the other 4 times a day for 96 hours 4. Dexamethasone + acetazolamide group: Dexamethasone acetate 4 mg oral every 6 hours and acetazolamide 250 mg twice a day orally for 96 hours 	
Outcomes	<p>This trial did not specify by primary or secondary outcomes</p> <ol style="list-style-type: none"> 1. Incidence of AMS in recreational climbers to moderate altitudes 2. Prophylactic benefit of the 2 drugs 3. Safety profile of administering dexamethasone and acetazolamide under conditions of moderate altitudes and physical exertion 	
Notes	<ol style="list-style-type: none"> 1. Trial Registration: Not stated 2. Funder: Not stated 3. Role of funder: Not stated 4. A priori sample size estimation: No 5. Conducted: Not reported 6. Declared conflicts of interest: No 	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement

Zell 1988 (Continued)

Random sequence generation (selection bias)	Unclear risk	Quote "Participants were randomly assigned ..." (Page 542) Insufficient information to score this item as low or high risk of bias
Allocation concealment (selection bias)	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Insufficient information to score this item as low or high risk of bias
Incomplete outcome data (attrition bias) All outcomes	Low risk	No participants were lost to follow-up
Selective reporting (reporting bias)	Low risk	Reporting bias was not detected
Other bias	Low risk	No other biases were identified

Zheng 2014

Methods	<ol style="list-style-type: none"> 1. Design: Double-blind randomized controlled trial 2. Country: China 3. Multisite: No 4. International: No 5. Treatment duration: 5 days 6. Follow-up: Unclear 7. Rate of ascent: None 8. Final altitude: 3900 metres 9. AMS scale: Lake Louise Scoring System (LLS)
Participants	<p>138 healthy young men, lowland resident, were recruited Randomized into 3 groups: Budesonide group (n= 46; 33.3%) Dexamethasone (n= 46; 33.3%) Placebo (n= 46; 33.3%)</p> <p>2. Loss to follow-up: Before intervention, 10 participants were lost to follow-up due to personal reasons (4, 3, and 3 in the budesonide, dexamethasone, and placebo groups, respectively) During intervention, 4 participants in the dexamethasone group encountered adverse reactions and discontinued medication before receiving any examination at altitude 124 participants completed the trial, whose data were included in analyses Main characteristics of participants: Age (mean): 20.3 years (range 18 - 35 years old)</p>

	Percentage/number of women/men: 100% men
Interventions	1. Budesonide group: oral starch tablets + inhalation of budesonide (200 µg twice a day) 2. Dexamethasone group: empty inhalers + dexamethasone tablets (4 mg twice a day) 3. Placebo group received both inhaled and oral placebos
Outcomes	1. Primary outcome measure was the incidence of AMS at altitude 2. Secondary outcome measures: Incidence of AMS in severe form, its severity reflected by Lake Louise Scoring System (LLS) score, heart rate, SpO ₂ , spirometric parameters, sleep quality assessed by questionnaires, and adverse reactions related to the investigational drugs
Notes	1. Trial Registration: not stated 2. Funder: This study was supported by the Special Health Research Project, Ministry of Health of P.R. China (grant No. 201002012) 3. A priori sample size estimation: Yes 4. Conducted: Not stated 5. Declared conflicts of interest: Yes. None declared

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "An independent physician randomly assigned the subjects to three groups: the budesonide, dexamethasone, and placebo groups, using a computer-generated random number list with an allocation ratio of 1:1:1" (Page 1002)
Allocation concealment (selection bias)	Low risk	Quote: "An independent physician randomly assigned the subjects to three groups: the budesonide, dexamethasone, and placebo groups, using a computer-generated random number list with an allocation ratio of 1:1:1" (Page 1002)
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Quote: "Empty inhalers could not be distinguished from budesonide inhalers by vision or feel. Starch tablets were similar to dexamethasone in shape, size, and color" (Page 1004) "The subjects, researchers, and other physicians were blinded" (Page 1004)
Blinding of outcome assessment (detection bias) All outcomes	Low risk	"The subjects, researchers, and other physicians were blinded" (Page 1004)

Zheng 2014 (Continued)

Incomplete outcome data (attrition bias) All outcomes	Low risk	Lost to follow-up : 4/42 in budesonide group (9.5%), 3/39 in dexamethasone group (7.7%), 3/43 in the placebo group (7%)
Selective reporting (reporting bias)	Low risk	Reporting bias was not detected
Other bias	Low risk	No other biases were identified

ACTH = Adrenocorticotrophic hormone; am = Ante meridiem/Before noon; AMS = Acute Mountain Sickness; AMS-C = Acute Mountain Sickness score- cerebral subscale ; AMS-R = Acute Mountain Sickness score- respiratory subscale; BP = Blood pressure; ESQ scores = Environmental Symptom Questionnaire; FVC = Forced vital capacity; g/dL = grams/decilitre; GHAQ = Generalized High Altitude Questionnaire; HACE = High altitude cerebral oedema; HAH = High altitude headache; HAI = High altitude illness; HAPE = High altitude pulmonary oedema; ITT = Intention-to-treat; IV = Intravenous; kg = Kilograms; LLQ = Lake Louise questionnaire; LLS = Lake Louise Scoring System; MAP = Mean artery pressure; mg = milligrams; NSAIDs = Nonsteroidal anti-inflammatory drugs; PASP = Pulmonary Artery Systolic Pressure; PEF = Peak expiratory flow; pm = post meridiem: After noon; PH = degree of acidity or alkalinity of a solution; RCT = randomized controlled trial; SD = Standard deviation; SE = Standard error; SEM = standard error of the mean; VAS = Visual analogue scale.

Characteristics of excluded studies [ordered by study ID]

Study	Reason for exclusion
ACME-1 2006	The study is focused on treatment of high altitude illness
Agostoni 2013	This study is not focused on prevention of high altitude illness
Bartsch 1993	The study is focused on treatment of high altitude illness
Bilo 2015	This study is not focused on prevention of high altitude illness
Bloch 2009	Non-randomized clinical trial
Broome 1994	The study is focused on treatment of high altitude illness
Bärtsch 1994	The study is focused on treatment of high altitude illness
Cain 1966	This study is not focused on prevention of high altitude illness
Debevec 2015	This study is not focused on prevention of high altitude illness
Dumont 1999	This study is not focused on prevention of high altitude illness

(Continued)

Forster 1982	This study is not focused on prevention of high altitude illness
Forward 1968	This study is not focused on prevention of high altitude illness
Fulco 2011	This study is not focused on prevention of high altitude illness
Gertsch 2002	This study is not focused on prevention of high altitude illness
Gray 1971	The study is focused on treatment of high altitude illness
Harris 2003	The study is focused on treatment of high altitude illness
Johnson 1988	This study is not focused on prevention of high altitude illness
Jonk 2007	This study is not focused on prevention of high altitude illness
Kotwal 2015	This study is not focused on prevention of high altitude illness
Lalande 2009	This study is not focused on prevention of high altitude illness
Lawley 2012	The study is focused on treatment of high altitude illness
Levine 1989	This study is not focused on prevention of high altitude illness
Liu 2013	This study is not focused on prevention of high altitude illness
Mairer 2012	This study is not focused on prevention of high altitude illness
McIntosh 1986	This study is not focused on prevention of high altitude illness
Purkayastha 1995	This study is not focused on prevention of high altitude illness
Reinhart 1994	This study is not focused on prevention of high altitude illness
Sandoval 2000	This study is not focused on prevention of high altitude illness
Scalzo 2015	This study is not focused on prevention of high altitude illness
Serra 2001	This study is not focused on prevention of high altitude illness
Siebenmann 2011	This study is not focused on prevention of high altitude illness
Singh 1969	The study is focused on treatment of high altitude illness
Solís 1984	This study is not focused on prevention of high altitude illness

(Continued)

Suh 2015	Non-randomized clinical trial
Teppema 2007	This study is not focused on prevention of high altitude illness
Vuyk 2006	This study is not focused on prevention of high altitude illness
White 1984	This study is not focused on prevention of high altitude illness
Wright 1988	This study is not focused on prevention of high altitude illness

Characteristics of studies awaiting assessment [ordered by study ID]

[Dugas 1995](#)

Methods	Double-blind randomized study
Participants	20 healthy volunteers received 5 mg of isradipine (n = 10) or placebo (n = 10) for 8 days. After 5 days of treatment in normoxia, the participants were rapidly transported to an altitude of 4350 m
Interventions	Israpadine (calcium channel blocker) and placebo
Outcomes	AMS symptom score, haemodynamic parameters and renal function
Notes	Full text not available (January 2016)

[Ellsworth 1987](#)

Methods	Double-blind randomized study
Participants	47 climbers participated in this double-blind, randomized trial comparing acetazolamide 250 mg, dexamethasone 4 mg, and placebo every 8 hours as prophylaxis for acute mountain sickness during rapid, active ascent of Mount Rainier (elevation 4392 metres). 42 participants (89.4 %) achieved the summit in an average of 34½ hours after leaving sea level
Interventions	Acetazolamide 250 mg, dexamethasone 4 mg, and placebo every 8 hours
Outcomes	Acute mountain sickness, symptoms reported
Notes	Full text not available (January 2016)

Furian 2016

Methods	Double-blind randomized placebo-controlled trial
Participants	112 COPD patients were studied in Bishkek (760 m), Kyrgyz Republic, after travelling for 6 hours to Tuja Ashu clinic (3200 m) and staying there for 3 days
Interventions	Participants received dexamethasone (2 x 4 mg/d) or placebo before ascent and during stay at 3200 metres
Outcomes	Cumulative incidence of 1 of the following: AMS (AMSc environmental symptom cerebral score ≥ 0.7), severe hypoxaemia (SpO ₂ < 75% for > 30 mins) or discomfort requiring descent to low altitude
Notes	Full text not available (January 2017)

Hefzi 2014

Methods	Double-blind placebo-controlled trial
Participants	29 participants were assigned to a treatment group (14) receiving 800 IU vitamin E, 1000 mg vitamin C, 200,000 IU vitamin A, and 600 mg N-acetylcystein daily, starting 2 months prior to the expedition, or to a placebo group (15)
Interventions	Vitamin group and placebo
Outcomes	AMS scores, Levels of endothelial microparticles
Notes	Full text not available (January 2016)

Kasic 1991

Methods	Randomized study
Participants	24 people who presented with acute mountain sickness
Interventions	A simulated descent of 1432 m (4600 ft) was attained by placing the participants in a fabric hypobaric chamber and pressurizing the chamber to 120 mmHg above ambient pressure. Participants were randomly assigned to either the hypobaric treatment or treatment with 4 litres of oxygen given by facemask; both treatments lasted for 2 hours
Outcomes	Mean arterial oxygen saturation (SaO ₂), symptoms of acute mountain sickness
Notes	Full text not available (January 2016)

Lee 2011

Methods	Randomized trial
Participants	19 adolescents aged 13 - 18 years attempting an ascent of Mount Kalapatar (5500 m)
Interventions	Acetazolamide, methazolamide.
Outcomes	Incidence of AMS, oxygen saturation and pulse rate
Notes	Full text not available (January 2017)

Pun 2014

Methods	Prospective double-blind placebo-controlled randomized trial
Participants	358 pilgrims were recruited at Dhunche (1950 metres) and followed up at Chandanbari (3350 m), and up to the sacred lake Gosaikunda. Most of these pilgrims ascended from Dhunche to the lake in 2 - 3 days
Interventions	Low-dose acetazolamide (125 mg) and placebo
Outcomes	Lake Louise score (LLS) for AMS measurement, arterial oxygen saturation (SpO ₂) and heart rate
Notes	Full text not available (January 2016)

Roncin 1996

Methods	Randomized trial
Participants	44 participants were enrolled in a study of the preventive effect of Ginko biloba extract (EGb 761) on acute mountain sickness (AMS) and vasomotor changes of the extremities during a Himalayan expedition
Interventions	Ginko biloba extract (EGb 761) 160 mg and placebo
Outcomes	ESQ score and the cold gradient measured by photoplethysmography
Notes	Full text not available (January 2016)

Swenson 1997

Methods	Randomized trial
Participants	19 healthy volunteers were assessed, who ingested in randomized order both a high carbohydrate (68% CHO) or normal carbohydrate (45% CHO) diet for 4 days. On the 4th day, participants were exposed to 8 hours of 10% normobaric oxygen
Interventions	High carbohydrate (68% CHO) or normal carbohydrate (45% CHO) diet for 4 days

Swenson 1997 (Continued)

Outcomes	Lake Louise Consensus Questionnaire, interleukins 1 beta, 6 and 8 (IL-1 beta, IL-6, IL-8) and tumour necrosis factor alpha (TNF-alpha)
Notes	Full text not available (January 2016)

Utz 1970

Methods	None known
Participants	None known
Interventions	None known
Outcomes	None known
Notes	Full text not available (January 2016)

Wang 1998

Methods	Randomized trial
Participants	65 men
Interventions	Conventional therapy group received oxygen, intravenous furosemide, aminophylline and dexamethasone; nifedipine group received oral nifedipine (10 mg, three times a day) in addition to conventional therapy; and participants in the nitric oxide group received nitric oxide (10 ppm) inhalation for 30 mins, in addition to oral nifedipine
Outcomes	Pulmonary rales on auscultation and shadows on chest radiograph
Notes	Full text not available (January 2016)

Xiangjun 2014

Methods	Randomized trial
Participants	80 healthy young male plain residents (17 - 33 years old)
Interventions	Inhalation of budesonide (200 µg twice a day), procaterol tablet (25 µg twice a day), inhalation of budesonide/fomoterol (160 µg/4.5 µg, twice a day) or placebo (1 tablet, twice a day)
Outcomes	Lake Louis AMS questionnaire, blood pressure, heart rate, and oxygen saturation
Notes	Full text not available (January 2017)

AMS: Acute Mountain Sickness; CHO: Carbohydrate; EGb 761: Extract of Ginkgo biloba 761; ESQ: Environmental Symptom Questionnaire; HR: Heart rate; IL: Interleukine; LLS: Lake Louise score; mg: milligrams; min: minutes; ppm: parts per million; TNF: Tumor necrosis factor.

Characteristics of ongoing studies *[ordered by study ID]*

ChiCTR-TRC-13003319

Trial name or title	Oral zolpidem for improving sleep and then prevention of acute mountain sickness: a single centre, randomized, double-blind, controlled, prospective trial
Methods	Interventional
Participants	<p>Inclusion criteria:</p> <ol style="list-style-type: none"> 1. Aged between and including 18 and 35 years 2. People rapidly ascending to high altitude. The gender ratio depends on actual situation 3. There is no history of plateau for a long time exposure 4. Before assessment, all participants must be voluntary and sign a written informed consent <p>Exclusion criteria:</p> <ol style="list-style-type: none"> 1. Recent history of taking sleeping pills 2. Engaged in specialized sports training 3. Participants cannot take the drugs in our trial because of allergic history or other reasons 4. Participants with bad compliance 5. Participants with serious illnesses, e.g. sleep apnoea 6. Recent history of upper respiratory tract infection 7. The driver 8. Participants with psychological or neurological disorder, and other conditions which are not appropriate for our trial <p>Gender: both</p>
Interventions	<p>Experimental: Oral zolpidem (10 mg, qd, oral)</p> <p>Control: Oral placebo, the same dosage as oral zolpidem</p>
Outcomes	Lake Louise Score
Starting date	30 June 2013
Contact information	Huang Lan
Notes	Recruiting

ChiCTR-TRC-13003590

Trial name or title	The meaning of intravenous iron supplementation in acute mountain sickness: a randomized, double-blinded, placebo-controlled trial
Methods	Interventional

ChiCTR-TRC-13003590 (Continued)

Participants	<p>Inclusion criteria:</p> <ol style="list-style-type: none"> 1. Healthy people ready to travel from Beijing to Tibet by air 2. Participants knowing the aim of the study and giving informed consent <p>Exclusion criteria:</p> <ol style="list-style-type: none"> 1. Not finishing the procedure 2. Coronary heart disease, uncontrolled hypertension and other severe diseases 3. Anaemia, especially iron deficiency anaemia <p>Age minimum: 18 years old Age maximum: 65 years old Gender: Both</p>
Interventions	<p>Intervention group: Intravenous iron 200 mg Control: Placebo</p>
Outcomes	Serum iron; Lake Louise score
Starting date	30 July 2013
Contact information	Ren Xuewen
Notes	Recruiting

NCT00886912

Trial name or title	Prevention of acute mountain sickness by intermittent hypoxic training
Methods	Interventional
Participants	<p>Inclusion criteria:</p> <ol style="list-style-type: none"> 1. Healthy 2. Non-smoker 3. Endurance training minimum twice a week <p>Exclusion criteria:</p> <ol style="list-style-type: none"> 1. Any diseases 2. Previous exposure to altitudes higher than 2000 metres (last 6 weeks) <p>Age minimum: 18 years old Age maximum: 55 years old Gender: Both</p>
Interventions	<ol style="list-style-type: none"> 1. Hypoxia 2. Normoxia
Outcomes	<p>Incidence of acute mountain sickness (time frame: after 20 hours at 4559 metres) Severity of acute mountain sickness (time frame: after 20 hours at 4559 metres)</p>
Starting date	June 2008
Contact information	Kai Schommer, MD

NCT00886912 (Continued)

Notes	Recruiting
-------	------------

NCT01606527

Trial name or title	Prospective, double-blind, randomized, placebo-controlled trial of ibuprofen versus placebo for prevention of neurologic forms of altitude sickness
Methods	Evaluating ibuprofen versus placebo for the prevention of neurological forms of altitude illness, including high altitude headache (HAH), acute mountain sickness (AMS), high altitude cerebral edema (HACE) and High Altitude Anxiety
Participants	The study will take place in the spring and summer of 2012 at the Marine Corps Mountain Warfare Training Center in the Eastern Sierras near Bridgeport, California. US Marines from near sea level will participate in battalion-level training exercises at between 8500 and 11,500 feet, where some altitude illness is expected
Interventions	Ibuprofen 600 mg orally three times daily Placebo, same schedule
Outcomes	Change in the incidence of AMS as measured on the Lake Louise AMS Questionnaire Change in High Altitude Headache measured by a visual analogue scale (VAS) Change in cognitive performance as measured by King-Devick Change in the presence of anxiety and somatic symptoms using the BSI-12 screening tool Change in the oxygen concentration using pulse oximetry Change in hydration status as measured by urine specific gravity Change in HAH incidence and severity as measured on the Lake Louise AMS Questionnaire Change in cognitive performance as measured by the Quickstick Change in the presence of anxiety and somatic symptoms using the GAD-2 screening tool Incidence of severe AMS as measured by a score of 6 or more on the Lake Louise AMS Questionnaire
Starting date	July 2012
Contact information	Jeffrey Gertsch MD, Naval Health Research Center
Notes	The recruitment status of this study is unknown. The completion date has passed and the status has not been verified in more than 2 years

NCT01682551

Trial name or title	Evaluation of the prevention and treatment effects of Chinese medicine on high altitude illness
Methods	Interventional
Participants	Inclusion criteria: 1. Healthy adults Exclusion criteria: 1. Chronic disease: cardiovascular disease, psychological disease, anaemia, migraine 2. Long-term use of the following materials: Chinese herbs, steroid, antibiotics

NCT01682551 (Continued)

	3. Altitude acclimation: have been to mountain over 2000 metres in the past month 4. Pregnancy Age minimum: 20 years Age maximum: 70 years Gender: Both
Interventions	Drug: acetazolamide Drug: Chinese Medicine
Outcomes	Incidence of acute mountain sickness will be measured by the Lake Louise Self Report (Lake Louise Score = 4 with headache) (time frame: the Lake Louise Score will be measured at 12 pm of the second day after hiking to determine the onset of AMS) Arterial oxygen saturation (time frame: before and after the hiking) Blood pressure (time frame: before and after the hiking) Heart rate (time frame: before and after the hiking)
Starting date	September 2012
Contact information	Not stated
Notes	Not yet recruiting

NCT01794078

Trial name or title	A randomized, 4-sequence, double-blind study to test the safety of combined dosing with aminophylline and ambrisentan in exercising healthy human volunteers at simulated high altitude
Methods	Interventional
Participants	Inclusion criteria: 1. Written informed consent to participate in the study prior to undergoing any screening procedures. The participant will be given a signed and dated copy of the informed consent 2. Participants must be healthy non-smoking (for 6 months or longer at start of Cycle 1) adult male and female volunteers; at least 18 through 50 years at screening, with a BMI of 18 - 33 kg/m ² and weighing at least 143 pounds. (65 kg). Participants' health status will be determined by medical history, physical examination, vital signs, ECG, blood chemistry, haematology, and urinalysis performed at screening 3. Be willing to fast for a minimum of 2 hours prior to screening 4. Be willing to abstain from alcohol and xanthine-containing food and beverages from 48 hours before check-in for each study day 5. Women who are of non-childbearing potential must be: a) Surgically sterile (removal of both ovaries and/ or uterus at least 12 months prior to dosing) and with an FSH level at screening of 40 m IU/mL b) Naturally postmenopausal (spontaneous cessation of menses) for at least 24 consecutive months prior to dosing on Day 1, and with an FSH level at screening of 40 m IU/mL 6. Women of child-bearing potential must have a negative serum or urine pregnancy test at screening, during the study, and must agree to avoid pregnancy during study and for 3 months after the last dose of study drug. Pregnancy is tested at screening, during check-in of each testing cycle, during the follow-up visit, and at any given point if deemed necessary by the physician or designate. During treatment, women of child-

bearing potential must use 2 acceptable methods of contraception at the same time unless she has had a documented tubal sterilization or chooses to use a Copper T 380A IUD or LNG 20 IUS, in which case no additional contraception is required. Abstinence is not considered a form of contraception. Medically acceptable contraceptives include: (1) documented surgical sterilization (such as a hysterectomy), (2) barrier methods (such as a condom or diaphragm) used with a spermicide, or (3) an intrauterine device (IUD) or intrauterine system (IUS)

7. Male participants must agree to take all necessary measures to avoid causing pregnancy in their sexual partners during the study and for 3 months after the last dose of study drug. Medically acceptable contraceptives include: (1) surgical sterilization (such as a vasectomy), or (2) a condom used with a spermicidal. Contraceptive measures such as Plan B (TM), sold for emergency use after unprotected sex, are not acceptable methods for routine use

8. Agree not to donate blood, platelets, or any other blood components 30 days, or plasma 90 days, prior to consenting and for 1 month after the last dose

9. Male participants must agree not to donate sperm during the study and for 12 weeks after the last dose

Exclusion criteria:

1. People with laboratory results outside the normal range, if considered clinically significant by the physician or delegate. In addition, they must have a haemoglobin concentration of 12.0 g/dL

2. A mental capacity that is limited to the extent that the person cannot provide legal consent or understand information regarding the side effects of the study drug

3. Currently abusing drugs or alcohol or with a history of drug or alcohol abuse within the past 2 years

4. Unwillingness or unable to comply with the protocol, or to co-operate fully with the physician and site personnel

5. Use of any of the following:

a) Any concomitant medication including oral contraceptive hormones. People who have received any prescribed or non-prescribed (over-the-counter) systemic medication, topical medications, or herbal supplements within 14 days from Day 1. St. John's Wort (hypericin) must not have been taken for at least 30 days prior to Cycle 1, Day 1

b) Any drugs, foods or substances known to be strong inhibitors or strong inducers of CYP enzymes (also known as cytochrome P450 enzymes)

6. Clinically significant ECG abnormality, in the opinion of the physician or delegate

7. Vital signs or clinically significant laboratory values at the screening visit that in the opinion of the physician or delegate would make the person an inappropriate candidate for the study

8. A VO₂ max value of less than 42 mL/kg/minute, as determined during exercise testing at screening. This value represents an educated estimate, and may be changed, to include new information, at the discretion of the physician

9. A history of, or otherwise indicated predisposition for, claustrophobia, i.e. the fear of closed, narrow spaces (because of the limited size of the high altitude chamber)

10. A history of "undeserved" altitude sickness, i.e. altitude sickness at only moderate altitude. This would consist of altitude-related headaches, dizziness, or nausea during plane rides, or when travelling to moderately elevated locations of less than 2743.2 metres/9000 ft

11. Has taken any other investigational drug during the 30 days prior to the screening visit or is currently participating in another investigational drug clinical trial

12. Made any significant donation or had a significant loss of blood within 30, or donated plasma within 90 days of consenting

13. Receipt of a transfusion or any blood products within 90 days prior to start of Cycle 1

14. History or manifestation of clinically significant neurological, gastrointestinal, renal, hepatic, cardiovascular, psychological, pulmonary, metabolic, endocrine, haematologic or other medical disorders. For the purpose of the study, individual fitness and health are more important than family history of disease burden as

NCT01794078 (Continued)

	<p>a criterion for participation. For example, an individual may have significant family history of cardiovascular disease; however, the individual's active lifestyle makes a manifestation of such disease at a young age unlikely. To account for such expected variation, the ultimate decision whether to exclude or include an individual based on family history or manifestation of disease will be made by the physician. The physician may choose to use physiological assessments, such as, e.g. ECG, blood pressure, and VO² max fitness level as an aid for decision-making</p> <p>15. Any condition that might interfere</p> <p>Age minimum: 18 years old</p> <p>Age maximum: 50 years old</p> <p>Gender: Both</p>
Interventions	<p>Drug: Ambrisentan 5 mg</p> <p>Drug: Aminophylline 400 mg</p>
Outcomes	<p>The safety of combined or single-dose aminophylline and ambrisentan at simulated altitude in exercising adults (time frame: Safety endpoints will be measured during simulated high altitude (Cycle 2) at least 22 days post-screening)</p> <p>The safety of combined or single-dose aminophylline and ambrisentan at simulated high altitude in resting adults (time frame: Safety endpoints will be measured during an episode of simulated high altitude (Cycle 1) , at least 7 days post-screening)</p>
Starting date	September 2013
Contact information	Claude A Piantadosi, MD
Notes	Active, not recruiting

NCT01993667

Trial name or title	Acetazolamide for the prevention of high altitude illness: a comparison of dosing
Methods	Interventional
Participants	<p>Inclusion criteria:</p> <ol style="list-style-type: none"> 1. 18 years or older 2. English or Indian speaking 3. Mountaineers or trekkers who plan to climb Mount McKinley or trek to Base Camp on Mount Everest <p>Exclusion criteria:</p> <ol style="list-style-type: none"> 1. Low sodium and/potassium blood serum levels 2. Kidney disease or dysfunction 3. Liver disease, dysfunction, or cirrhosis 4. Suprarenal gland failure or dysfunction 5. Hyperchloremic acidoses 6. Angle-closure glaucoma 7. Taking high-dose aspirin (over 325 mg/day) 8. Any reaction to sulfa drugs or acetazolamide 9. Pregnant or lactating women

NCT01993667 (Continued)

Interventions	Drug: Acetazolamide
Outcomes	Prevention of acute mountain sickness as measured by the Lake Louise Score (time frame: 1 year) Side effect profile of acetazolamide (time frame: 1 year)
Starting date	March 2012
Contact information	Scott McIntosh, MD
Notes	Recruiting

NCT02244437

Trial name or title	Ibuprofen versus acetaminophen in the prevention of acute mountain sickness: A double-blind, randomized controlled trial
Methods	Interventional
Participants	<p>Inclusion criteria: Healthy adults between the ages of 18 and 65, men or women, non-Nepali, without AMS or any concurrent illness, and not already taking NSAIDs and acetazolamide or any other drug for the prevention of altitude illness</p> <p>Exclusion criteria: Individuals not meeting inclusion criteria, including mild AMS (more than one mild symptom on the Lake Louise Questionnaire) or significantly depressed oxygen saturation (< 75%); women known to be pregnant, cannot exclude the possibility of being pregnant, or have missed menses by over 7 days; individuals who have spent 24 hours at an altitude of 4500 metres/14,000 feet within the last 9 days; anyone known to have taken any of the following in the last 2 days: acetazolamide (Diamox®), steroids (dexamethasone, prednisone), theophylline, or diuretics (Lasix®); individuals who have a known intracranial space-occupying lesion or a history of elevated intracranial pressure. (i.e. tumours, hydrocephalus, etc)</p> <p>Age minimum: 18 years old Age maximum: 65 years old Gender: Both</p>
Interventions	Drug: Acetaminophen Drug: Ibuprofen
Outcomes	<p>Diagnosis of Acute Mountain Sickness (AMS) (Time Frame: Upon reaching 5000 metres altitude (Lobuche) of Nepal Himalaya)</p> <p>Blood Oxygen Saturation (SPO₂) (time frame: Upon reaching 5000 metres altitude (Lobuche) of Nepal Himalaya)</p> <p>Heart Rate (HR) (time frame: Upon reaching 5000 metres altitude (Lobuche) of Nepal Himalaya)</p> <p>High Altitude Headache (HAH) (time frame: Upon reaching 5000 metres altitude (Lobuche) of Nepal Himalaya)</p>
Starting date	October 2014
Contact information	Nicholas C Kanaan, MD

NCT02244437 (Continued)

Notes	Active, not recruiting
-------	------------------------

NCT02450968

Trial name or title	Dexamethasone for prophylaxis of acute mountain sickness in people with chronic obstructive pulmonary disease travelling to altitude
Methods	Interventional
Participants	<p>Inclusion criteria:</p> <ol style="list-style-type: none"> 1. Chronic obstructive pulmonary disease (COPD), GOLD criteria grade 1 - 2 2. Living at low altitude (< 800 metres) <p>Exclusion criteria:</p> <ol style="list-style-type: none"> 1. COPD exacerbation 2. Severe COPD, GOLD grade 3 or 4 3. Arterial oxygen saturation < 92% at low altitude (< 800 metres) 4. Diabetes, uncontrolled cardiovascular disease such as systemic arterial hypertension, coronary artery disease; previous stroke; pneumothorax in the last 2 months 5. Untreated or symptomatic peptic ulcer disease, glaucoma, obstructive sleep apnoea 6. Internal, neurologic or psychiatric disease that interfere with protocol compliance including current heavy smoking (> 20 cigarettes a day) 7. Pregnant or nursing women <p>Age minimum: 20 years old Age maximum: 75 years old Gender: Both</p>
Interventions	<p>Drug: Dexamethasone</p> <p>Drug: Placebo</p>
Outcomes	<p>Acute mountain sickness, cumulative incidence (time frame: day 3 at 3200 metres)</p> <p>6 minutes walk distance (time frame: Day 2 at 3200 metres)</p> <p>Acute mountain sickness, severity (time frame: day 1, day 2, day 3 at 3200 metres)</p> <p>Arterial blood gases (time frame: Day 2 at 3200 metres)</p> <p>Perceived exertion (time frame: Day 2 at 3200 metres)</p>
Starting date	May 2015
Contact information	Talant M Sooronbaev, MD
Notes	Recruiting

NCT02604173

Trial name or title	A randomized controlled trial of altitude sickness prevention and efficacy of comparative treatments
Methods	Interventional
Participants	<p>Inclusion criteria:</p> <ol style="list-style-type: none"> 1. Men and women 2. Sea level-dwelling hikers 3. Between ages 18 and 65 <p>Exclusion criteria:</p> <ol style="list-style-type: none"> 1. History of allergy to acetazolamide or budesonide (or other corticosteroids) 2. Taken NSAIDs, acetazolamide, or corticosteroids in the week prior to study enrolment 3. Hazardous medical conditions which preclude the ability to moderately hike to high altitude, including: sickle cell anaemia, asthma, or COPD, severe anaemia, or severe coronary arterial disease 4. Pregnancy or suspected pregnancy 5. Participants under 18 years of age or more than 65 6. Sleep above 4000 elevation in the preceding 1 week 7. History of asthma or COPD 8. Current symptoms of an acute upper respiratory illness 9. Unable to complete a moderately strenuous hike at high altitude <p>Gender: Both</p>
Interventions	<p>Drug: Acetazolamide</p> <p>Drug: Budesonide</p> <p>Drug: Placebo</p>
Outcomes	<p>Oxygen saturation (time frame: 24 hours)</p> <p>Pulmonary function testing - FEV1 (Time frame: 24 hours)</p> <p>Pulmonary function testing - FVC (time frame: 24 hours)</p> <p>Pulmonary function testing - PEFr (Time frame: 24 hours)</p>
Starting date	August 2016
Contact information	Grant S Lipman, MD
Notes	Not yet recruiting

NCT02811016

Trial name or title	Effect of inhaled budesonide on the incidence and severity of acute mountain sickness at 4559 metres
Methods	Not stated
Participants	51 healthy volunteers
Interventions	<p>Budesonide 200 µg inhaled at 7:00 a.m. and 7 p.m.</p> <p>Budesonide 800 µg inhaled at 7:00 a.m. and 7 p.m.</p> <p>Placebo Inhalation at 7:00 a.m. and 7 p.m.</p>

NCT02811016 (Continued)

Outcomes	Assessment of incidence and severity of acute mountain sickness by use of 2 internationally standardized and well-established questionnaires Venous (and capillary) blood drawings Transthoracic echocardiography for assessing pulmonary artery systolic pressure
Starting date	June 2016
Contact information	Marc Berger, Salzburger Landeskliniken
Notes	This study has been completed.

NCT02941510

Trial name or title	Inhaled budesonide for altitude illness prevention
Methods	Not stated
Participants	Participants will be recruited from the Denver community and prescreened for eligibility via phone. 100 participants, after consenting, will have baseline data and blood collected and will begin budesonide therapy 72 hours prior to being taken from Denver to Pikes Peak, where they will be observed at altitude for 18 hours. Participants will have the opportunity to withdraw consent at any time and will be monitored continuously by physician-researchers
Interventions	Budenoside, placebo
Outcomes	Primary outcome measures: <ul style="list-style-type: none"> • Changes in inflammation • Incidence of Acute Mountain Sickness (AMS) • Changes in gene regulation
Starting date	April 2017
Contact information	University of Colorado, Denver
Notes	This study is not yet open for participant recruitment.

AMS: Acute Mountain Sickness; BMI: Body mass index; COPD: Chronic obstructive pulmonary disease ; CYP: cytochrome P450 enzymes; dL: decilitre; ECG: electrocardiogram; FEV1: forced expiratory volume in 1 second; FSH: Follicle-stimulating hormone; ft: feet; FVC: forced expiratory vital capacity; GOLD: Global Initiative for Chronic Obstructive Lung Disease criteria,; HAH: High altitude headache; HR: hear rate; kg: kilograms; IUD: Intrauterine device; IUS: Intrauterine system; LNG 20: levonorgestrel 20 µg/day; ml: millilitres; Mg: milligrams; NSAIDs: Nonsteroidal anti-inflammatory drugs ; OTC: over-the-counter; PEFr: peak expiratory flow rate ; qd: every day; TM: Morning-after pill; VO²: maximal oxygen consumption.

DATA AND ANALYSES

Comparison 1. Carbonic anhydrase inhibitors: acetazolamide versus placebo

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Incidence of acute mountain sickness	16	2301	Risk Ratio (M-H, Random, 95% CI)	0.47 [0.39, 0.56]
1.1 Acetazolamide 250 - 255 mg	4	855	Risk Ratio (M-H, Random, 95% CI)	0.60 [0.39, 0.94]
1.2 Acetazolamide 500 mg	8	1111	Risk Ratio (M-H, Random, 95% CI)	0.48 [0.38, 0.61]
1.3 Acetazolamide 750 mg	2	80	Risk Ratio (M-H, Random, 95% CI)	0.33 [0.18, 0.62]
1.4 Other combinations	2	255	Risk Ratio (M-H, Random, 95% CI)	0.30 [0.17, 0.55]
2 Incidence of high altitude pulmonary oedema	7	1138	Risk Ratio (M-H, Random, 95% CI)	0.0 [0.0, 0.0]
3 Incidence of high altitude cerebral oedema	6	1126	Risk Ratio (M-H, Random, 95% CI)	0.32 [0.01, 7.48]
4 Incidence of adverse events: Paraesthesia	5	789	Risk Ratio (M-H, Random, 95% CI)	5.53 [2.81, 10.88]
4.1 Acetazolamide 250 mg	1	197	Risk Ratio (M-H, Random, 95% CI)	12.63 [4.02, 39.64]
4.2 Acetazolamide 500 mg	3	370	Risk Ratio (M-H, Random, 95% CI)	6.72 [3.94, 11.46]
4.3 Acetazolamide 750 mg	1	222	Risk Ratio (M-H, Random, 95% CI)	3.09 [2.00, 4.78]
5 Differences in HAI/AMS scores	6		Std. Mean Difference (Random, 95% CI)	Subtotals only
5.1 acetazolamide 250 mg	3		Std. Mean Difference (Random, 95% CI)	0.19 [0.01, 0.37]
5.2 acetazolamide 500 mg	4		Std. Mean Difference (Random, 95% CI)	-0.57 [-1.20, 0.07]

Comparison 2. Steroids: budesonide vs. placebo

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Incidence of acute mountain sickness	2	132	Risk Ratio (M-H, Random, 95% CI)	0.37 [0.23, 0.61]

Comparison 3. Steroids: dexamethasone vs. placebo

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Incidence of acute mountain sickness	4	176	Risk Ratio (M-H, Random, 95% CI)	0.60 [0.36, 1.00]
2 Differences in HAI/AMS scores	3	50	Std. Mean Difference (IV, Random, 95% CI)	-0.46 [-1.21, 0.29]

Comparison 4. Calcium modulators: nifedipine vs. placebo

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Differences in HAI/AMS scores	2	48	Std. Mean Difference (IV, Random, 95% CI)	-0.56 [-1.85, 0.74]

Comparison 5. NSAIDs and other analgesic: aspirin vs. placebo

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Incidence of AMS	2	60	Risk Ratio (M-H, Random, 95% CI)	0.35 [0.06, 1.95]

Comparison 6. NSAIDs and other analgesic: ibuprofen vs. placebo


Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Incidence of acute mountain sickness	3	598	Risk Ratio (M-H, Random, 95% CI)	0.64 [0.49, 0.82]

Analysis 1.1. Comparison 1 Carbonic anhydrase inhibitors: acetazolamide versus placebo, Outcome 1 Incidence of acute mountain sickness.


Review: Interventions for preventing high altitude illness: Part 1. Commonly-used classes of drugs

Comparison: 1 Carbonic anhydrase inhibitors: acetazolamide versus placebo

Outcome: 1 Incidence of acute mountain sickness



(Continued ...)




Analysis 1.2. Comparison 1 Carbonic anhydrase inhibitors: acetazolamide versus placebo, Outcome 2 Incidence of high altitude pulmonary oedema.

Review: Interventions for preventing high altitude illness: Part I. Commonly-used classes of drugs

Comparison: 1 Carbonic anhydrase inhibitors: acetazolamide versus placebo

Outcome: 2 Incidence of high altitude pulmonary oedema




Analysis 1.3. Comparison 1 Carbonic anhydrase inhibitors: acetazolamide versus placebo, Outcome 3 Incidence of high altitude cerebral oedema.

Review: Interventions for preventing high altitude illness: Part I. Commonly-used classes of drugs

Comparison: 1 Carbonic anhydrase inhibitors: acetazolamide versus placebo

Outcome: 3 Incidence of high altitude cerebral oedema




Analysis 1.4. Comparison 1 Carbonic anhydrase inhibitors: acetazolamide versus placebo, Outcome 4 Incidence of adverse events: Paraesthesia.

Review: Interventions for preventing high altitude illness: Part 1. Commonly-used classes of drugs

Comparison: 1 Carbonic anhydrase inhibitors: acetazolamide versus placebo

Outcome: 4 Incidence of adverse events: Paraesthesia




Analysis 1.5. Comparison 1 Carbonic anhydrase inhibitors: acetazolamide versus placebo, Outcome 5 Differences in HAI/AMS scores.

Review: Interventions for preventing high altitude illness: Part 1. Commonly-used classes of drugs

Comparison: 1 Carbonic anhydrase inhibitors: acetazolamide versus placebo

Outcome: 5 Differences in HAI/AMS scores




Analysis 2.1. Comparison 2 Steroids: budesonide vs. placebo, Outcome 1 Incidence of acute mountain sickness.

Review: Interventions for preventing high altitude illness: Part 1. Commonly-used classes of drugs

Comparison: 2 Steroids: budesonide vs. placebo

Outcome: 1 Incidence of acute mountain sickness




Analysis 3.1. Comparison 3 Steroids: dexamethasone vs. placebo, Outcome 1 Incidence of acute mountain sickness.

Review: Interventions for preventing high altitude illness: Part 1. Commonly-used classes of drugs

Comparison: 3 Steroids: dexamethasone vs. placebo

Outcome: 1 Incidence of acute mountain sickness




Analysis 3.2. Comparison 3 Steroids: dexamethasone vs. placebo, Outcome 2 Differences in HAI/AMS scores.

Review: Interventions for preventing high altitude illness: Part 1. Commonly-used classes of drugs

Comparison: 3 Steroids: dexamethasone vs. placebo

Outcome: 2 Differences in HAI/AMS scores




Analysis 4.1. Comparison 4 Calcium modulators: nifedipine vs. placebo, Outcome 1 Differences in HAI/AMS scores.

Review: Interventions for preventing high altitude illness: Part 1. Commonly-used classes of drugs

Comparison: 4 Calcium modulators: nifedipine vs. placebo

Outcome: 1 Differences in HAI/AMS scores




Analysis 5.1. Comparison 5 NSAIDs and other analgesic: aspirin vs. placebo, Outcome 1 Incidence of AMS.

Review: Interventions for preventing high altitude illness: Part 1. Commonly-used classes of drugs

Comparison: 5 NSAIDs and other analgesic: aspirin vs. placebo

Outcome: 1 Incidence of AMS




Analysis 6.1. Comparison 6 NSAIDs and other analgesic: ibuprofen vs. placebo, Outcome 1 Incidence of acute mountain sickness.

Review: Interventions for preventing high altitude illness: Part 1. Commonly-used classes of drugs

Comparison: 6 NSAIDs and other analgesic: ibuprofen vs. placebo

Outcome: 1 Incidence of acute mountain sickness



ADDITIONAL TABLES

Table 1. Main characteristics of included studies

Study	High mountain	Men (%)	In-creased risk of AMS, HAPE or HACE	Country	Admin-istration timing	Trekking	Final al-titude (mts)	Differ-ence be-tween the end-point and the baseline altitude (mts)	Dura-tion of ascent	Defini-tion de AMS	Conflict of interest

Table 1. Main characteristics of included studies (Continued)

Anony- mous 1981	Yes	100	No	Ecuador	3 days	No (Car)	5000	2225	5 days	No defi- nition was pro- vided	No
AS- CENT 2012	Yes	72.4	No	Nepal	unclear	Yes	4928	648	Unclear	Lake Louise AMS score \geq 3 with headache	No
Ban- deret 1977	Yes	54.2	No	USA	2 days	No (Car)	4300	4100	5 hours	No defi- nition was pro- vided	No
Bartsch 1991	Yes	95.2	Previous episodes of HAPE	Italy	4 days	No (Car)	4559	3429	1 day	No defi- nition was pro- vided	No
Basnyat 2003	Yes	67.1	No	Nepal	2-3 days	Yes	4937	2937	2-3 days	Lake Louise AMS score= headache + 1 symp- tom	Yes
Basnyat 2008	Yes	626	No	Nepal	max 4 dias	Yes	5000	750	36-96 hours	Lake Louise AMS score \geq 3 with headache	Yes
Basu 2002a	Yes	100	No	India	2 days	Yes	3450	3230	3 days	No defi- nition was pro- vided	No
Basu 2002b	Yes	100	No	Nepal	2 days	No (Flight)	3450	3230	Unclear	Lake Louise AMS score	No

Table 1. Main characteristics of included studies (Continued)

Bates 2011	Yes	58	No	Chile	4-5 days		5200	Unclear		Lake Louise AMS score ≥ 3	No
Baumgartner 2003	No	100	No	Switzerland	7 days	No applicable	4559	4069	13 minutes	ESQ= AMS-C SCORE > 70	No
Bernhard 1994	Yes	65.2	40% subjects with previous AMS mild or moderate	Bolivia	4 days	No (Car)	5334	1645	2 hours	Modified ESQ= 3 cerebral symptoms, one with intensity ≥ 2	Yes
Bernhard 1998	Yes	69.2	50% of the subjects had previously visited high altitudes and had experienced mild to moderate AMS	Bolivia	4 days	No (Car)	5334	1645	2 hours	Modified ESQ= 3 cerebral symptoms, one with intensity ≥ 2	No
Bradwell 1986	Yes	90.4	No	Nepal	3 days	Yes	4846	3546	10 days	No definition was provided	No
Burki 1992	Yes	Unclear	No	Pakistan	2 days	No (Car)	4450	3932	8 hours	No definition was provided	No
Burtscher 2001	Yes	64	History of headache	Unclear	2 hours	No (combination)	3480	2880	Unclear	Headache scoring	No

Table 1. Main characteristics of included studies (Continued)

Burtscher 2014	Yes	Unclear	History of AMS	Italy	10 hours	No (combination)	3800	3200	Less than a day by car up to 3480, and 2.8 to 3 hours climbing from there to 3800m	Lake Louise AMS score ≥ 3	Yes
Burtscher 1998	Yes	58.6	History of headache	Unclear	1 hour	Unclear	3480	2880	Unclear	Headache scoring	Yes
Carlsten 2004	Yes	62.6	No	Nepal	2 hours	No (Flight)	3630	3630	7-8 hours	Lake Louise AMS score ≥ 4	Yes
Chen 2015	Yes	Unclear	No	China	3 days	No (Flight)	3700	3200	2.5 hour	Lake Louise AMS score ≥ 3	No
Chow 2005	Yes	57.8	No	USA	5 days	No (Car)	3800	2570	2 hours	Lake Louise AMS score ≥ 5	No
Ellsworth 1991	Yes	61.1	No	USA	1 day	No (combination)	4392	3262	1 day	Modified ESQ = AMS-C > 0,7 + AMS-R > 0,6	No
Faull 2015	Yes	70	Unclear	Italy	3 days	No (Cable-cars or train)	3459	3309	Unclear	No definition was provided	No
Fischer 2000a	No	100	No	Germany	3 days	No applicable	4500	4500	30 min	No definition was provided	No

Table 1. Main characteristics of included studies (Continued)

Fischer 2000b	Yes	100	No	Switzerland	3 days	No (Cable-cars or train)	3454	3454	3 hours	No definition was provided	No
Fischer 2004	No	100	No	Germany	3 days	No applicable	4500	4500	15 minutes	ESQ-C score >0,5 or Lake Louise AMS score >3	No
Fulco 2006	No	83.3	No	USA	1 days	No applicable	4300	4300	Unclear	Modified ESQ= AMS-C >0,7 + AMS-R >0,6	No
Greene 1981	Yes	91.6	No	Nepal	2 days	Yes	5895	3895	5 days	No definition was provided	No
Hackett 1976	Yes	71	No	Nepal	4 days	Yes	4243	803	3-4 days	Questionnaire clinical >2	No
Hackett 1988	Yes	100	No	USA	1 hour	No (Flight)	4400	4400	1 hour	AMS Score >2 or Modified ESQ= AMS-C >0,7 + AMS-R >0,6	No
HEAT 2010	Yes	70.5	No	Nepal	1 day	Yes	4928	648	Unclear	No definition was provided	Yes
Hillenbrand	Yes	100	Unclear	Nepal	Unclear	Yes	4930	1490	7 days	Lake Louise	Yes

Table 1. Main characteristics of included studies (Continued)

2006										AMS score ≥ 3 with headache	
Hochapfel 1986	Yes	61,00	No	India	5 days	Yes	5500	2100	9 days	No definition was provided	No
Hohenhaus 1994	Yes	86,00	susceptibility to AMS	Italy	3 days	No (combination)	4559	4069	22 hours	Score clinical proposed at the International Hypoxia symposium+ Do you feel ill? = Yes	Yes
Hussain 2001	Yes	100	No	Pakistan	1 day	No (combination)	4578	4063	1 day	ESQ score ≥ 6	No
Jain 1986	Yes	100	No	USA	1 day	Unclear	3500	3300	Unclear	No definition was provided	No
Johnson 1984	No	100	No	USA	1 day	No applicable	4570	4570	Unclear	Modified ESQ = AMS-C $> 0,7$ + AMS-R $> 0,6$	No
Kayser 2008	Yes	unclear	No		1 day	No (combination)	5896	5896	7 days	Lake Louise AMS score ≥ 3 with headache	No

Table 1. Main characteristics of included studies (Continued)

Ke 2013	Yes	100	No	China	3 days	No (Flight)	3658	Unclear	3 hours	Presence of headache and at least one of the symptoms of nausea or vomiting, fatigue, dizziness, or difficulty sleeping, and a total score of at least 3,	Yes
Küpper 2008	Yes	100	No	Italia	5 days	Yes	4559	4559	2 days	Lake Louise AMS score ≥ 4	No
Larson 1982a	Yes	unclear	No	USA	1 day	Yes	4394	3094	2 days	GHAQ = Headache moderate or more and/or nausea moderate or more	No
Larson 1982b	Yes	84.3	No	USA	1 day	Yes	4394	3094	2 days	GHAQ = Headache moderate or more and/or nausea	No

Table 1. Main characteristics of included studies (Continued)

											moderate or more	
Lipman 2012	Yes	67.4	No	USA	6 hours	No (combination)	3810	2570	12 hours	Lake Louise AMS score ≥ 3 with headache	Yes	
Luks 2007	No	unclear	No	USA	4 days	No applicable	3900	2490	Unclear	No definition was provided	Yes	
Maggiorini 2006	Yes	86.2	History of HAPE	Italia	1 day	No (combination)	4559	4069	2 days	Lake Louise AMS score ≥ 4	Yes	
Mir-rakhlmov 1993	Yes	Unclear	Patients with asthma	Kir-guistán	2 days	No (Car)	3200	2440	4 hours	No definition was provided	No	
Montgomery 1989	Yes	74	No	USA	1,5 days	Unclear	2700	2700	Unclear	AMS score clinical = 3 or more symptoms with a grade 2 or greater	No	
Moraga 2007	Yes	100	No	Chile	3 days	No (Cable-cars or train)	3696	3696	8,5 hours	AMS score clinical ≥ 3 or 1 symptom = 3	No	
Muza 2004 Defl	No	unclear	No	USA	1 hour	No applicable	4300	4300	Unclear	Lake Louise AMS score ≥ 3	Yes	

Table 1. Main characteristics of included studies (Continued)

PACE 2006	Yes	60 to 69	No	Nepal	6 days	Yes	4928	1488	Unclear	Lake Louise AMS score \geq 3	No
Parati 2013	Yes	95	No	Italy	3 days	No (combination)	4559	4437	<28 hours	Lake Louise AMS score \geq 3	Yes
PHAIT 2004	Yes	70 to 74	No	Nepal	2 days	Yes	4928	648	Unclear	Lake Louise AMS score \geq 3 with headache	Yes
Rock 1987	Yes	44	No	USA	2 days	No (Flight)	4300	4300	6 hours	Modified ESQ= AMS-C>0,7 + AMS-R>0,6	No
Rock 1989a	No	100	No	USA	12 hours	No applicable	4570	4570	Unclear	Johnson Score \geq 1	No
Rock 1989b	No	100	No	USA	12 hours	No applicable	4570	4570	Unclear	Johnson Score \geq 1	No
Rock 1989c	No	100	No	USA	12 hours	No applicable	4570	4570	Unclear	Johnson Score \geq 1	No
Sartori 2002	Yes	unclear	susceptible to HAPE	Italy	<6 hours	No (combination)	4559	3429	22 hours	No definition was provided	No
SPACE 2011	Yes	62 to 72	No	Nepal	Unclear	Yes	5000	700	30 hours-4 days	Lake Louise AMS score= headache + 1 symptom	No

Table 1. Main characteristics of included studies (Continued)

Subudhi 2011	No	80	No	USA	1 day	No applicable	4875	3225	1 day	Lake Louise AMS score ≥ 3	Yes
Van Patot 2008	Yes	43 to 52	No	USA	3 days	No (Car)	4300	2700	Unclear	ESQ AMS-C Score ≥ 0 , 7 + Lake Louise AMS score ≥ 3 with headache	Yes
Wang 2013	Yes	44 to 62	No	Bolivia	3 days	No (Flight)	3561	3159	3 hours	No definition was provided	Yes
Wright 1983	Yes	95	Previous severe AMS= 6	Kenia	8 days	No (combination)	4790	3527	3 days	No definition was provided	No
Wright 2004	Yes	92	No	Nepal	Unclear	No (Car)	4680	4680	3 days	Lake Louise AMS score ≥ 3	No
Zell 1988	Yes	62 to 72	No	Nepal	2 days	No (combination)	4050	2710	3 days	No definition was provided	No
Zheng 2014	Yes	100	No	China	1 day	No (Car)	3900	3500	5 days	LLS includes 5 self-reporting symptoms: headache, gastrointestinal symptoms, fatigue/	No

Table 1. Main characteristics of included studies (Continued)

											weak- ness, dizzi- ness/ light- headed- ness and diffi- culty in sleeping. Each symp- tom is scores 0- 3
--	--	--	--	--	--	--	--	--	--	--	--

APPENDICES

Appendix I. Risk categories for acute mountain sickness

Risk categories	Description
Low	Individuals with no prior history of altitude illness and ascending to ≤ 2800 m/9186 feet.
Low	Individuals taking ≥ 2 days to arrive at 2500 - 3000 m/8202 - 9842 feet with subsequent increases in sleeping elevation < 500 m by day/1640 feet by day
Moderate	Individuals with prior history of AMS and ascending to 2500 - 2800 m (8202 - 9186 feet) in 1 day
Moderate	No history of AMS and ascending to > 2800 m (9186 feet) in 1 day
Moderate	All individuals ascending > 500 m/d (1640 feet) (increase in sleeping elevation) at altitudes above 3000 m/9842 feet
High	History of AMS and ascending to ≥ 2800 m/9186 feet in 1 day
High	All individuals with a prior history of HAPE or HACE

(Continued)

High	All individuals ascending to > 3500 m/11482 feet in 1 day
High	All individuals ascending > 500 m/1640 feet/d increase in sleeping elevation above > 3500 m/11482 feet
High	Very rapid ascents (e.g. Mount Kilimanjaro)

Appendix 2. Medical terms glossary

Term	Definition	Source
Anorexia	The lack or loss of appetite accompanied by an aversion to food and the inability to eat	www.ncbi.nlm.nih.gov/mesh
Ataxia	Impairment of the ability to perform smoothly coordinated voluntary movements	www.ncbi.nlm.nih.gov/mesh
Brian herniation	Protrusion of tissue, structure, or part of an organ through the bone, muscular tissue, or the membrane by which it is normally contained	www.ncbi.nlm.nih.gov/mesh
Dyspnoea	Difficult or laboured breathing.	www.ncbi.nlm.nih.gov/mesh
Dizziness	An imprecise term which may refer to a sense of spatial disorientation, motion of the environment, or light-headedness	www.ncbi.nlm.nih.gov/mesh
Endothelium	A layer of epithelium that lines the heart, blood vessels (endothelium vascular), lymph vessels (endothelium lymphatic), and the serous cavities of the body	www.ncbi.nlm.nih.gov/mesh
Fatigue	The state of weariness following a period of exertion, mental or physical, characterized by a decreased capacity for work and reduced efficiency to respond to stimuli	www.ncbi.nlm.nih.gov/mesh
Hallucination	Subjectively experienced sensations in the absence of an appropriate stimulus, but which are regarded by the individual as real	www.ncbi.nlm.nih.gov/mesh
Headache	The symptom of pain in the cranial region.	www.ncbi.nlm.nih.gov/mesh
Hypoxia	A disorder characterized by a reduction of oxygen in the blood	www.ncbi.nlm.nih.gov/mesh

(Continued)

Insomnia	Disorders characterized by impairment of the ability to initiate or maintain sleep	www.ncbi.nlm.nih.gov/mesh
Lightheadedness	See dizziness.	www.ncbi.nlm.nih.gov/mesh
Nausea	An unpleasant sensation in the stomach usually accompanied by the urge to vomit	www.ncbi.nlm.nih.gov/mesh
Pulmonary oedema	Excessive accumulation of extravascular fluid in the lung, an indication of a serious underlying disease or disorder. Pulmonary oedema prevents efficient pulmonary gas exchange in the pulmonary alveoli, and can be life-threatening	www.ncbi.nlm.nih.gov/mesh
Pulmonary alveoli	Small polyhedral outpouchings along the walls of the alveolar sacs, alveolar ducts and terminal bronchioles through the walls of which gas exchange between alveolar air and pulmonary capillary blood takes place	www.ncbi.nlm.nih.gov/mesh
Seizures	Clinical or subclinical disturbances of cortical function due to a sudden, abnormal, excessive, and disorganized discharge of brain cells. Clinical manifestations include abnormal motor, sensory and psychic phenomena	www.ncbi.nlm.nih.gov/mesh

Appendix 3. The most frequent adverse events of the pharmacological interventions

Drug	Description and contraindications	Adverse events	Source
Acetazolamide	Acetazolamide, an inhibitor of the enzyme carbonic anhydrase Hy-persensitivity to acetazolamide or any excipients in the formulation. Since acetazolamide is a sulphonamide derivative, cross sensitivity between acetazolamide, sulphonamide and other sulphonamide derivatives is possible. Acetazolamide therapy is contraindicated in situations in which sodium and/or potassium blood serum levels are depressed, in cases of	Adverse reactions, occurring most often early in therapy, include paraesthesias, particularly a “tingling” feeling in the extremities, hearing dysfunction or tinnitus, loss of appetite, taste alteration and gastrointestinal disturbances such as nausea, vomiting and diarrhoea; polyuria, and occasional instances of drowsiness and confusion	DailyMed

(Continued)

	marked kidney and liver disease or dysfunction, in suprarenal gland failure, and in hyperchloraemic acidoses. It is contraindicated in patients with cirrhosis because of the risk of development of hepatic encephalopathy		
Aspirin	it is a nonsteroidal anti-inflammatory drug.	Reye's syndrome (a rare but serious illness). Stomach bleeding	DailyMed
Bosentan	It is an endothelin receptor antagonist indicated for the treatment of pulmonary arterial hypertension Pregnancy, pre-existing hepatic impairment.	Elevations of liver aminotransferases (ALT, AST) and liver failure. Early liver injury may preclude future use as disease progresses Respiratory tract infection and anaemia	DailyMed
Dexamethasone	Glucocorticoids, naturally occurring and synthetic, are adrenocortical steroids that are readily absorbed from the gastrointestinal tract. Glucocorticoids cause varied metabolic effects. In addition, they modify the body's immune responses to diverse stimuli. Naturally occurring glucocorticoids (hydrocortisone and cortisone), which also have sodium-retaining properties, are used as replacement therapy in adrenocortical deficiency states. Their synthetic analog including dexamethasone are primarily used for their anti-inflammatory effects in disorders of many organ systems Contraindicated in systemic fungal infections.	Several adverse events (e.g. hyperglycaemia, fluid retention, hypokalaemic alkalosis, potassium loss, sodium retention)	DailyMed
Gabapentin	Gabapentin is an anticonvulsant. Gabapentin is contraindicated in patients who have demonstrated hypersensitivity to the drug or its ingredients	Somnolence, dizziness, ataxia, fatigue, and nystagmus	DailyMed

(Continued)

Ginkgo biloba	This homeopathic product has not been evaluated by the Food and Drug Administration for safety or efficacy. FDA is not aware of scientific evidence to support homeopathy as effective	-	DailyMed
Methazolamide	Methazolamide is a potent inhibitor of carbonic anhydrase. Methazolamide therapy is contraindicated in situations in which sodium and/or potassium serum levels are depressed, in cases of marked kidney or liver disease or dysfunction, in adrenal gland failure, and in hyperchloraemic acidoses. In patients with cirrhosis, use may precipitate the development of hepatic encephalopathy	Adverse reactions, occurring most often early in therapy, include paraesthesias, particularly a “tingling” feeling in the extremities; hearing dysfunction or tinnitus; fatigue; malaise; loss of appetite; taste alteration; gastrointestinal disturbances such as nausea, vomiting, and diarrhoea; polyuria; and occasional instances of drowsiness and confusion	DailyMed
Nifedipine	It is a calcium channel blocker. Nifedipine must not be used in cases of cardiogenic shock. It is contraindicated in patients with a known hypersensitivity to any component of the tablet	Headache, flushing/heat sensation, dizziness, fatigue/asthenia, nausea	DailyMed
Phenytoin	Phenytoin sodium is an antiepileptic drug. Phenytoin is contraindicated in those patients who are hypersensitive to phenytoin or other hydantoin	Central Nervous System (the most common manifestations encountered with phenytoin therapy are referable to this system and are usually dose-related. These include nystagmus, ataxia, slurred speech, decreased coordination, and mental confusion), Gastrointestinal System (nausea, vomiting, constipation, toxic hepatitis, and liver damage)	DailyMed
Salmeterol	Long-acting beta2-adrenergic agonist Contraindicated in patients with asthma. It should be used with caution in patients with cardiovascular disorders, espe-	It increases the risk of asthma-related death. Excessive beta-adrenergic stimulation has been associated with seizures, angina, hypertension or hypotension, tachycardia with rates up to	DailyMed

(Continued)

	cially coronary insufficiency, cardiac arrhythmias, and hypertension	200 beats/min, arrhythmias, nervousness, headache, tremor, palpitation, nausea, dizziness, fatigue, malaise, and insomnia	
Selective inhibitor of phosphodiesterase type 5 (taladafil and sildenafil)	It was shown to potentiate the hypotensive effects of nitrates, and its administration to patients who are using organic nitrates, either regularly and/or intermittently, in any form is therefore contraindicated	Headache and flushing	DailyMed
Spirolactone	Aldactone oral tablets contain 25 mg, 50 mg, or 100 mg of the aldosterone antagonist spironolactone Aldactone is contraindicated for patients with anuria, acute renal insufficiency, significant impairment of renal excretory function, or hyperkalaemia	Gynecomastia and hyperkalaemia	DailyMed
Sumatriptan	Sumatriptan is an agonist for a vascular 5-hydroxytryptamine ₁ receptor subtype. It should not be given to patients with history, symptoms, or signs of Ischaemic cardiac, cerebrovascular, or peripheral vascular syndromes	Serious cardiac events, including some that have been fatal. These events are extremely rare and most have been reported in patients with risk factors predictive of CAD. Events reported have included coronary artery vasospasm, transient myocardial ischemias, myocardial infarction, ventricular tachycardia, and ventricular fibrillation	DailyMed
Theophylline	Theophylline is classified as a methylxanthine. Theophylline should be used with extreme caution in patients with the following clinical conditions due to the increased risk of exacerbation of the concurrent condition: active peptic ulcer disease, seizure disorders and cardiac arrhythmias (not including bradyarrhythmias)	Nausea, vomiting, headache, and insomnia	DailyMed

Appendix 4. MEDLINE (Ovid SP) search strategy

1. Brain edema/ or Pulmonary edema/ or ((edema or oedema) adj3 (high altitude or cerebral or pulmonary)).mp. or ((mountain or high altitude) adj3 (sickness or illness)).mp. or high altitude.ti,ab.
2. Exp Primary Prevention/ or exp Drug Therapy/ or (drug therap* or prevent* or acclimati?ation or nifedipine or dexamethasone or taladafil or sildenafil or theophylline or salmeterol or acetazolamide or aspirin or sumatriptan or gabapentin or phenytoin or magnesium or ginkgo biloba or ascorbic acid or alpha-tocopherol acetate or alpha-lipoic acid or beta-carotene or selenium or zinc or bosentan or calcium channel blocker* or of phosphodiesterase type or nonsteroidal anti-inflammatory drug* or steroid* or glucocorticosteroid* or corticosteroid* or non-selective phosphodiesterase inhibitor* or carbonic anhydrase inhibitor* or beta agonist* or 5-HT1 receptor agonist* or N-methyl-D-aspartate antagonist* or antioxidant* or vitamin* or mineral* or endothelin antagonist*).mp.
3. ((randomized controlled trial or controlled clinical trial).pt. or randomized.ab. or placebo.ab. or clinical trials as topic.sh. or randomly.ab. or trial.ti.) not (animals not (humans and animals)).sh.
4. 1 and 2 and 3

Appendix 5. Embase (Ovid SP) search strategy

1. 'brain edema'/exp
2. 'lung edema'/exp
3. (edema OR oedema) NEXT/3 ('high altitude' OR 'altitude' OR 'cerebral' OR 'pulmonary')
4. ('mountain' OR 'high altitude') NEXT/3 ('sickness' OR 'diseases' OR 'illness')
5. #1 OR #2 OR #3 OR #4
6. 'primary prevention'/exp
7. 'drug therapy'
8. 'drug therap**'
9. 'therap**'
10. 'prevent**'
11. 'acclimati?ation'
12. 'nifedipine'
13. 'dexamethasone'
14. 'tadalafil'
15. 'sildenafil'
16. 'theophylline'
17. 'salmeterol'
18. 'acetazolamide'
19. 'acetylsalicylic acid'
20. 'aspirin'
21. 'sumatriptan'
22. 'gabapentin'
23. 'phenytoin'
24. 'magnesium'
25. 'ginkgo biloba'
26. 'ascorbic acid'
27. 'alpha tocopherol'
28. 'alpha-tocopherol acetate'
29. 'alpha-lipoic acid'
30. 'beta carotene'
31. 'selenium'
32. 'zinc'
33. 'bosentan'
34. 'calcium channel blocker**'
35. 'phosphodiesterase type'
36. 'nonsteroidal anti-inflammatory drug**'

37. steroid*
38. glucocorticosteroid*
39. 'non-selective phosphodiesterase inhibitor*'
40. 'carbonic anhydrase inhibitor*'
41. 'beta agonist*'
42. '5-ht1 receptor agonist*'
43. 'n-methyl-d-aspartate antagonist*'
44. antioxidant*
45. vitamin*
46. mineral*
47. 'endothelin antagonist*'
48. #6 OR #7 OR #8 OR #9 OR #10 OR #11 OR #12 OR #13 OR #14 OR #15 OR #16 OR #17 OR #18 OR #19 OR #20 OR #21 OR #22 OR #23 OR #24 OR #25 OR #26 OR #27 OR #28 OR #29 OR #30 OR #31 OR #32 OR #33 OR #34 OR #35 OR #36 OR #37 OR #38 OR #39 OR #40 OR #41 OR #42 OR #43 OR #44 OR #45 OR #46 OR #47
49. 'randomized controlled trial'
50. 'controlled clinical trial'
51. 'controlled clinical trial (topic)'
52. #49 OR #50 OR #51
53. #5 AND #48 AND #52
54. #53 AND 'human'/de
55. #53 AND 'human'/de AND [embase]/lim NOT [medline]/lim

Appendix 6. CENTRAL search strategy

1. MeSH descriptor: [Brain Edema] explode all trees
2. MeSH descriptor: [Pulmonary Edema] explode all trees
3. (?edema near (high?altitude or cerebral or pulmonary)) or ((mountain or high?altitude) near (sickness or illness)) or high?altitude:ti,ab
4. (#1 or #2 or #3)
5. MeSH descriptor: [Secondary Prevention] explode all trees
6. MeSH descriptor: [Primary Prevention] explode all trees
7. MeSH descriptor: [Drug Therapy] explode all trees
8. (drug therapy or prevent* or acclimatization or nifedipine or dexamethasone or taladafil or sildenafil or theophylline or salmeterol or acetazolamide or aspirin or sumatriptan or gabapentin or phenytoin or magnesium or ginkgo biloba or ascorbic acid or alpha-tocopherol acetate or alpha-lipoic acid or beta-carotene or selenium or zinc or bosentan or calcium channel blockers or selective inhibitor of phosphodiesterase type or nonsteroidal anti-inflammatory drug* or steroid* or glucocorticosteroid* or corticosteroid* or non-selective phosphodiesterase inhibitor* or carbonic anhydrase inhibitor* or beta agonist* or 5-HT1 receptor agonist* or N-methyl-D-aspartate antagonist* or antioxidant* or vitamin* or mineral* or endothelin antagonist*):ti,ab
9. (#5 or #6 or #7 or #8)
10. #4 and #9

Appendix 7. Search strategy for LILACS via BIREME interface

tw:(edema cerebral)) OR (tw:(edema pulmonar)) OR (tw:(edema))AND (tw:(enfermedad de altura)) OR (tw:(high-altitude sickness)) OR (tw:(mal agudo de montaña)) OR (tw:(montaña enfermedad\$)) OR (tw:(mal da montanha\$)) OR (tw:(doença de alta altitude\$)) OR (tw:(mal de altura*))

Appendix 8. WHO International Trials Registry Portal search

high-altitude pulmonary edema
Recruitment Status: All

Appendix 9. Study eligibility screening and data extraction form.

Intervention for preventing high altitude illness
Study Selection, Quality Assessment & Data Extraction Form

First author	Journal/Conference Proceedings etc	Year

Study eligibility

RCT/Quasi/CCT (delete as appropriate)	Relevant participants	Relevant interventions	Relevant outcomes
Yes / No / Unclear	Yes / No / Unclear	Yes / No / Unclear	Yes / No* / Unclear

* Issue relates to selective reporting when authors may have taken measurements for particular outcomes, but not reported these within the paper(s). Reviewers should contact trialists for information on possible non-reported outcomes & reasons for exclusion from publication. Study should be listed in 'Studies awaiting assessment' until clarified. If no clarification is received after three attempts, study should then be excluded.

Do not proceed if any of the above answers are 'No'. If study to be included in 'Excluded studies' section of the review, record below the information to be inserted into 'Table of excluded studies'

Freehand space for comments on study design and treatment:

References to trial

Check other references identified in searches. If there are further references to this trial link the papers now & list below. All references to a trial should be linked under one *Study ID* in RevMan.

Code each paper	Author(s)	Journal/Conference Proceedings etc	Year
	<i>The paper listed above</i>		
	<i>Further papers</i>		

Participants and trial characteristics

Participant characteristics	
	Further details
Age (mean, median, range, etc)	
Sex of participants (numbers / %, etc)	
Country	
Other	
Rate of ascent (m/h)	
Final altitude reached (metres)	
AMS scale	
History of HAI	
Type of HAI reported	

Intervention characteristics

Intervention characteristics	
	Further details
Name	
Doses	
Administration route	
Time to administration	

(Continued)

Duration	
----------	--

If RCT included a combination:

Intervention characteristics	
	Further details
Name	
Doses	
Administration route	
Time to administration	
Duration	

If RCT included acclimatization:

Intervention characteristics	
Rate of ascent (m/h)	Further details

Methodological quality

Allocation of intervention	
State here method used to generate allocation and reasons for grading	Grade (circle)
	Low risk of bias (Random)
	High risk of bias (e.g. alternate)
	Unclear

Concealment of allocation Process used to prevent foreknowledge of group assignment in a RCT, which should be seen as distinct from blinding	
State here method used to conceal allocation and reasons for grading	Grade (circle)
	Low risk of bias
	High risk of bias
	Unclear

Blinding	
Person responsible for participants care	Yes / No
Participant	Yes / No
Outcome assessor	Yes / No
Other (please specify)	Yes / No
Intention-to-treat	
An intention-to-treat analysis is one in which all the participants in a trial are analysed according to the intervention to which they were allocated, whether they received it or not	
All participants entering trial	
15% or fewer excluded	
More than 15% excluded	
Not analysed as 'intention-to-treat'	
Unclear	

Free selective report	
State here method used to generate allocation and reasons for grading	Grade (circle)
	Low risk of bias
	High risk of bias
	Unclear

Were withdrawals described? Yes ? No ? not clear ?

Discuss if appropriate

Data extraction

Outcomes relevant to your review	
Copy and paste from 'Types of outcome measures'	
	Reported in paper (circle)
Incidence of AMS (headache, nausea, insomnia, dizziness, and sleep disorder)	Yes / No
Incidence of HACE.	Yes / No
Incidence of HAPE.	Yes / No
Safety of adverse events	Yes / No
Safety (adverse drug reaction)	Yes / No

For Dichotomous data			
Code of paper	Outcomes	Intervention group (n) n = number of participants, not number of events	Control group (n) n = number of participants, not number of events
A	Incidence of AMS ((headache, nausea, insomnia, dizziness, and sleep disorder)		
	Incidence of HACE.		
	Incidence of HAPE		
	Safety of adverse events		
	Safety (adverse drug reaction)		

Other information which you feel is relevant to the results
 Indicate if: any data were obtained from the primary author; if results were estimated from graphs etc; or calculated by you using a formula (this should be stated and the formula given). In general if results not reported in paper(s) are obtained this should be made clear here to be cited in review

Freehand space for writing actions such as contact with study authors and changes

References to other trials

Did this report include any references to published reports of potentially eligible trials not already identified for this review?		
First author	Journal / Conference	Year of publication

Did this report include any references to unpublished data from potentially eligible trials not already identified for this review? If yes, give list contact name and details

(Continued)

Trial characteristics	
	Further details
Single centre / multicentre	
Country / Countries	
How was participant eligibility defined?	
How many people were randomized?	
Number of participants in each intervention group	
Number of participants who received intended treatment	
Number of participants who were analysed	
Drug treatment(s) used	
Dose / frequency of administration	
Duration of treatment (State weeks / months, etc, if cross-over trial give length of time in each arm)	
Median (range) length of follow-up reported in this paper (state weeks, months or years, or if not stated)	
Time-points when measurements were taken during the study	
Time-points reported in the study	
Time-points you are using in RevMan	
Trial design (e.g. parallel / cross-over*)	
Other	

Appendix 10. Transformation of numerical data- Secondary outcome: Differences in HAI/AMS scores

Study	Original data	Trans-formed data	Original data	Trans-formed data	Original data	Trans-formed data	Original data	Trans-formed data
Bates 2011	Median = 4	Mean = 4	IQR = 2 - 6	SD = 3.19	Median = 6.5	Mean = 6.41	IQR = 5 - 7.75	SD = 2.11
Bernhard 1994	SEM = 3.6	SD = 11.94	SEM = 4.8	SD = 16.63	-	-	-	-
Chow 2005	Median = 2	Mean = 2.25	RANGE = 0 - 5	SD = 1.28	Median = 4	Mean = 5.5	RANGE = 1 - 13	SD = 3.11
Chow 2005	Median = 4	Mean = 4.75	RANGE = 1 - 10	SD = 2.38	-	-	-	-
Hackett 1988	SEM = 0.6	SD = 1.58	SEM = 1.0	SD = 2.82	-	-	-	-
Hillenbrand 2006	Median = 1.0	Mean = 0.83	IQR = 0 - 1.5	SD = 1.12	Median = 1.0	Mean = 0.66	IQR = 0 - 1.0	SD = 0.74
Hohenhaus 1994	SE = 2.88	SD = 7.05	SE = 1.1	SD = 4.12	SE = 1.1	SD = 3.97	-	-
Hussain 2001	SE = 0.33	SD = 1.32	SE = 3.04	SD = 7.45	SE = 1.26	SD = 3.09	SE = 1.58	SD = 3.87
Rock 1989a	SE = 0.14	SD = 0.59	SE = 0.31	SD = 1.24	-	-	-	-
Rock 1989b	SE = 0.37	SD = 1.48	SE = 0.11	SD = 0.47	-	-	-	-
Rock 1989c	SE = 0.5	SD = 1	SE = 0.33	SD = 1.32	-	-	-	-
Wright 1983	SEM = 8	SD = 19.59	SE = 7	SD = 22.14	-	-	-	-

HISTORY

Protocol first published: Issue 4, 2012

Review first published: Issue 6, 2017

Date	Event	Description
17 April 2012	Amended	Contact details updated.

CONTRIBUTIONS OF AUTHORS

Conceiving the review: AMC

Co-ordinating the review: VNE, AMC and IAR

Undertaking manual searches: VNE, DMF, RDM, and IAR

Screening search results: VNE, DMF, RDM and IAR

Organizing retrieval of papers: VNE, DMF, RDM and IAR

Screening retrieved papers against inclusion criteria: VNE, DMF, RDM and IAR

Appraising quality of papers: VNE, DMF, RDM, AGG and IAR

Abstracting data from papers: VNE, DMF, RDM, AGG and IAR

Writing to authors of papers for additional information: Not performed

Providing additional data about papers: VNE, DMF, RDM, AGG and IAR

Obtaining and screening data on unpublished studies: VNE, DMF, RDM, AGG and IAR

Data management for the review: IAR and RDM

Entering data into Review Manager 5 ([RevMan 5.3](#)): IAR and RDM

RevMan statistical data: IAR and RDM

Other statistical analysis not using RevMan: AMC and IAR

Interpretation of data: VNE, DMF, RDM, AGG, AMC and IAR

Statistical inferences: VNE, DMF, RDM, AGG, AMC and IAR

Writing the review: VNE, DMF, RDM, AGG, AMC and IAR

Securing funding for the review: VNE, DMF, RDM, AGG, AMC and IAR

Performing previous work that was the foundation of the present study: Not performed

Guarantor for the review (one author): VNE

Person responsible for reading and checking review before submission: IAR

DECLARATIONS OF INTEREST

Victor H Nieto Estrada: nothing to declare.

Daniel Molano Franco: nothing to declare.

Roger David Medina: nothing to declare.

Alejandro Gonzalez Garay: nothing to declare.

Arturo Marti Carvajal: nothing to declare.

Ingrid Arevalo-Rodriguez: nothing to declare.

SOURCES OF SUPPORT

Internal sources

- Fundacion Universitaria de Ciencias de la Salud, Colombia.
- Methodology Research Unit/Neonatology, Instituto Nacional de Pediatría, Mexico.

Academic.

- Instituto de Evaluación Tecnológica en Salud - IETS, Colombia.

External sources

- Iberoamerican Cochrane Center, Spain.

Academic.

DIFFERENCES BETWEEN PROTOCOL AND REVIEW

Given that the original protocol was published in 2012, several sections needed updating to fulfil the current methodological guidelines for Cochrane Reviews. We made the following changes to the published protocol ([Martí-Carvajal 2012](#)):

1. On the recommendation of the editors of the Cochrane Anaesthesia, Critical and Emergency Group, we split the review into three parts, considering the numerous interventions assessed for HAI prevention. This review is the first part and it focuses on commonly-used agents to prevent this condition. Subsequent reviews will address less commonly-used agents to prevent HAI, and non-pharmacological interventions. This change has implications for the title and scope of this review and for later reviews in this series.
2. We updated the [Background](#) with new references to reflect current evidence about the target condition, as well as the scope of common interventions to prevent HAI.
3. We modified the primary and secondary outcomes presented in the protocol ([Martí-Carvajal 2012](#)) to follow the MECIR guidelines and improve their comprehensibility.
4. For this review, we selected six commonly-used types of intervention to prevent HAI. We will address other interventions in the next two reviews belonging to this series.
5. Despite the fact that the protocol did not include any consideration of unit of analysis issues, we have identified 12 cross-over studies in our searches. We have included them in our review to enhance the full reporting of all available evidence, and we have analysed them separately from the parallel studies.
6. We stated in the protocol that we would contact trial authors in case of missing data or selective reporting. However we were unable to conduct this task, usually due to the year of publication of the trial (most of the publications were performed too long ago and it was not possible to obtain a valid contact address or other means to contact trialists).

7. We have introduced several modifications in the [Dealing with missing data](#) section, in order to clarify the intention-to-treat analysis performed and to present the methods for imputing missing information (mostly related to standard deviations).

8. Under [Data synthesis](#) we added the trial sequential analysis procedure, in order to test the boundary before the required information size was reached.

9. We also made extensive modifications to the [Subgroup analysis and investigation of heterogeneity](#) section, and have selected only three variables to analyse. However, we were unable to find information about the third factor (significant pre-existing disease) in the included trials.